



2-2009

W147-Virginia Buttonweed

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: http://trace.tennessee.edu/utk_agexturf

Recommended Citation

"W147-Virginia Buttonweed," The University of Tennessee Agricultural Extension Service, W147-2/09 09-0161,
http://trace.tennessee.edu/utk_agexturf/5

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Weed, Insect and Disease Control is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Turfgrass by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

TURFGRASS SCIENCE

at the University of Tennessee

Virginia Buttonweed (*Diodia virginiana*)

Greg Breeden, Weed Science Extension Assistant
 James T. Brosnan, Assistant Professor, Turfgrass Weed Science
 Plant Sciences

Introduction

Virginia buttonweed (*Diodia virginiana*) is one of the most difficult-to-control broadleaf weeds in turf. It commonly proliferates in poorly drained areas and can tolerate mowing heights as low as one-half inch. This deep-rooted perennial produces both above- and below-ground flowers. Its prolific seed production, extensive root system and ability to vegetatively reproduce make control extremely difficult.

Virginia Buttonweed Identification

Virginia buttonweed is a spreading perennial with opposite leaves (Figure 1) that often have a mottled, yellow appearance due to a virus that commonly infects the foliage. Stems are occasionally hairy. A unique characteristic of Virginia buttonweed is that it produces both above- and below-ground flowers that self-pollinate. The above-ground flowers are white and star-shaped with four petals (Figure 2). Fruit are often green and elliptical with hairy ridges (Figure 3). Virginia buttonweed produces deep taproots and rhizomes that often start at the nodes. Rhizomes can be found as deep as several feet below the soil sur-

face. Virginia buttonweed can form very dense mats in established turf. Above-ground fruit are buoyant, allowing them to be transported with surface water to other areas. Virginia buttonweed can also reproduce from stem fragments. If not removed, clippings from mowing or weedeating can establish into mature plants. Hand removal of Virginia buttonweed is often ineffective, because any stem fragments or rhizomes left behind can produce a viable plant.



Figure 1. Mat of Virginia buttonweed (*Diodia virginiana*)



Figure 2. Virginia buttonweed (*Diodia virginiana*) flower



Figure 3. Virginia buttonweed (*Diodia virginiana*) fruiting structure

Table 1. Herbicide options for Virginia buttonweed control

Product Name	Active Ingredients	Rate/ac	Turf Species
Confront	triclopyr + clopyralid	1-2 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass, Centipedegrass
Corsair	chlorsulfuron	3 oz.	Kentucky Bluegrass, Fine Fescue, Bermudagrass, Zoysiagrass, Centipedegrass
Escalade II	2,4-D + fluroxypyr + dicamba	2-3 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Manor	metsulfuron	1 oz.	Kentucky Bluegrass, Fine Fescue, Bermudagrass, Zoysiagrass, Centipedegrass
Millennium Ultra 2	2,4-D + clopyralid + dicamba	2-3 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Momentum fx2	2,4-D + triclopyr + fluroxypyr	3-4 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass, Centipedegrass
Monument	trifloxysulfuron	0.56 oz.	Bermudagrass, Zoysiagrass
Powerzone	carfentrazone + MCPA + MCPP + dicamba	2-4 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Speedzone	carfentrazone + 2,4-D + MCPP + dicamba	3-5 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Spotlight	fluroxypyr	0.67-2.5 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass, Centipedegrass
Surge	sulfentrazone + 2,4-D + MCPP + dicamba	3-4 pts.	Kentucky Bluegrass, Fine Fescue, Tall Fescue, Perennial Ryegrass, Bermudagrass, Zoysiagrass
Three-way Selective, Trimec Classic, Trimec Southern, Triplet, Others	2,4-D + MCPP + dicamba	Product-Dependent	Product-Dependent

Virginia Buttonweed Control Options

Option #1 - Sulfonylurea herbicides: Two applications (four weeks apart) of Corsair (chlorosulfuron) at 3 oz/acre, Manor (metsulfuron) at 1 oz/acre or Monument (trifloxysulfuron) at 0.56 oz/acre are the best programs to control Virginia buttonweed utilizing sulfonylurea herbicides. See Table 1 for a complete list of herbicides active against Virginia buttonweed. Beginning applications in May (before Virginia buttonweed begins to flower) will improve control, as will including a non-ionic surfactant at 0.25 percent v/v in the spray solution.

Option # 2 - Phenoxy and Phenoxy-type herbicides: Sequential applications of materials that contain three or more phenoxy (or phenoxy-type) herbicides can be used to control Virginia buttonweed in established turf. Materials containing clopyralid or fluroxypyr in the mixture have performed well in research at the University of Tennessee. Clopyralid is found in such herbicides as Confront (triclopyr and clopyralid) and Millenium Ultra (2,4-D, clopyralid and dicamba), while fluroxypyr can be found in herbicides such as Spotlight (fluroxypyr) and Escalade II (fluroxypyr, 2,4-D and MCPP).

Sequential applications of Confront at 1 or 2 pt/acre, Millenium Ultra at 2.5 pt/acre and Escalade II at 2 or 3 pt/acre are the most effective options for control of Virginia buttonweed with phenoxy or phenoxy-type herbicides. See Table 1 for a complete list of herbicides active against Virginia buttonweed. Multiple applications of Confront will injure bermudagrass at the 2 pt/acre rate, so be sure to use the lower rate of 1 pt/acre.

Final Thoughts

Virginia buttonweed control in established turf can be difficult due to the plant's rhizomatous growth habit, extensive root system, aggressive growth and ability to reproduce vegetatively. Sequential applications of postemergence herbicides will be required to provide effective control.

Always refer to the product label for specific information on proper product use, tank-mix compatibility and turfgrass tolerance.

For more information on turfgrass weed control, visit the University of Tennessee's turfgrass weed science Web site, <http://tennesseeturfgrassweeds.org>

This replaces the original publication of the same name developed by Scott McElroy and Greg Breeden in 2007.

Disclaimer

This publication contains herbicide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the herbicide applicator's responsibility, by law, to read and follow all current label directions for the specific herbicide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.

Visit the UT Extension Web site at
<http://www.utextension.utk.edu/>

W147-2/09 09-0161