

#### University of Tennessee, Knoxville Trace: Tennessee Research and Creative Exchange

Physics and Astronomy Publications and Other Works

Physics and Astronomy

7-2010

# Observation of the first Bs--> J/Psi Phi Event in CMS

Stefan M Spanier University of Tennessee - Knoxville, sspanier@utk.edu

Giordano Cerizza University of Tennessee - Knoxville, gcerizza@utk.edu

Follow this and additional works at: http://trace.tennessee.edu/utk\_physastrpubs Part of the <u>Elementary Particles and Fields and String Theory Commons</u>

**Recommended** Citation

Spanier, Stefan M and Cerizza, Giordano, "Observation of the first Bs--> J/Psi Phi Event in CMS" (2010). *Physics and Astronomy Publications and Other Works*. http://trace.tennessee.edu/utk\_physastrpubs/4

This Presentation is brought to you for free and open access by the Physics and Astronomy at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Physics and Astronomy Publications and Other Works by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

## Observation of the first $B_s \rightarrow J/\psi \phi$ Event at CMS

**Giordano Cerizza**, Stefan Spanier University of Tennessee, Knoxville

for the CMS Collaboration 35<sup>th</sup> International Conference on High Energy Physics Paris, July 22 – 28, 2010 View into the open CMS detector – the **innermost cylinder** houses the silicon detector that measures the tracks of charged particles. The group at UTK contributes to this detector's operation and systematic studies. We also instrument the beam pipe below it with diamond detectors to protect it against adverse beam conditions.



The B<sub>s</sub> particle is produced in proton-proton collision with the Large Hadron Collider (LHC). It is observed in its decay into two particles, J/ $\psi$  and  $\phi$ , that in turn decay into longer lived charged particles that can be detected by sub-detectors of the CMS experiment. The hit positions along the particle trajectories measured by several layers of particle track detectors are input to a track reconstruction algorithm. The reconstructed particle tracks are displayed on the next page.

The tracks in red correspond to the two particles from the sub-sequent decay of the J/ $\psi$  particle – the two particles are muons, particles that are abundant in the cosmic radiation. The yellow tracks correspond to the so-called kaons from the  $\phi$  particle. The red and yellow tracks intersect at the point where the Bs particle decayed – this happens somewhat displaced from the point where the protons collide due to the long lifetime of this B<sub>s</sub> particle. This displaced vertex is a distinct signature and allows to identify it. The green tracks are other by-products of the collision.

The Bs particle and its peculiar decay are of interest for us to study a fundamental symmetry that might explain how the apparent matter antimatter asymmetry occurred during the evolution of the Universe starting from the Big-Bang.



CMS Experiment at LHC, CERN Data recorded: Sun Jul 4 01:33:41 2010 EDT Run/Event: 139364 / 20750462 Lumi section: 20  $B_s \rightarrow J/\psi \phi$  candidate event

View along the proton beam – protons collided at the yellow point in the center. Reconstructed charged particle tracks and hits in the outer muon detectors.

**K**<sup>--</sup>

### View along the proton beam



#### Side-view of the CMS detector



#### Zoom into the production region



The B<sub>s</sub> particle was created on July 4<sup>th</sup>, 2010, in the CMS detector, and found soon thereafter by our graduate student Giordano Cerizza. He was able to measure several parameters that confirm the nature of this decay.

The particle moves with relativistic speed. Therefore its proper lifetime of only 1.5 10<sup>-12</sup>s is dilated so that it travels nearly 5mm in the detector before it decays.

#### Measured parameters:

 $Mass(J/\psi K^+K^-) = 5346.5 MeV/c^2$  $Mass(\mu^{+}\mu^{-}) = 3104.3 MeV/c^{2}$  $Mass(K^{+}K^{-}) = 1018.57 MeV/c^{2}$  $B^{vtx}$  Prob. = 0.78 Proper decay length = 895 ± 50 μm  $ct/\sigma_{ct} = 17.8$  $\beta \gamma = 5.9$  $p(\mu^{\star}) = 15.5 \text{ GeV/c} (global)$  $p(\mu) = 7.6 \text{ GeV/c (global)}$  $p_{\tau}(\mu^{+}) = 2.9 \text{ GeV/c (global)}$  $p_{\tau}(\mu) = 2.1 \text{ GeV/c (global)}$  $p_{\tau}(K^{+}) = 1.4 \text{ GeV/c}$  $p_{\tau}(K) = 2.2 \text{ GeV/c}$