



4-2005

SP341-K Common Stalk Borer in Field Corn

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: http://trace.tennessee.edu/utk_agexcrop

 Part of the [Agronomy and Crop Sciences Commons](#), and the [Entomology Commons](#)

Recommended Citation

"SP341-K Common Stalk Borer in Field Corn," The University of Tennessee Agricultural Extension Service, SP431-K (Rev) 4/05 E12-4615 05-0324, http://trace.tennessee.edu/utk_agexcrop/27

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Insect, Pest and Disease Control - Corn is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Field & Commercial Crops by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Insects

The Common Stalk Borer in Field Corn

Charles R. Patrick, Professor, Entomology and Plant Pathology
Scott D. Stewart, Associate Professor, Entomology and Plant Pathology
Angela Thompson, Assistant Professor, Plant Sciences and Landscape Systems

Introduction

The common stalk borer in Tennessee is reported to attack several crops in addition to field corn. However, the preferred host of the insect is giant ragweed. This preference is due to the largeness of the stem in which the larvae can mature. Fields of corn planted into a no-till situation are preferred by the moths because of the grassy weeds in those fields. The adult moth deposits eggs on grassy weeds in the early spring. Later, the larvae will migrate to any corn planted in the fields.

Description

The adult moth is dark grayish-brown with a number of small white spots on the wing. The hind wings are pale gray-brown. The larva is very small upon hatching, with brownish and white stripes and a dark purple band around the middle of the body. This purple band fades as the larva matures. The larva reaches a length of 1½ inches when fully grown. (See Figure 1).


Life History

Adult female moths deposit their eggs on grasses in the fall, and the eggs overwinter on the grasses. In the spring, the eggs hatch and bore into the stems of the grass. The larva then burrows up and down the stem, feeding on the



Figure 1. The Common Stalk Borer

tissue until it becomes too large for that stem. The larva will then migrate to a larger-stemmed plant and bore into it. If this plant is corn, the larva will mature and pupate inside the corn stalk. There does not appear to be more than one generation per year in Tennessee.



Damage

Damage is done when the larva bores up and down the stalk, eating out the heart of the corn plant. Corn is usually attacked when the plants are less than 2 feet high. The larva may enter the corn plant from the side of the stem or from the top of the plant and move down into the whorl. Damaged plants may be recognized by a dead leaf in the center of the whorl. (See Figure 2).

Control Practices

A thorough cleanup of weeds in the field is essential to prevent the larvae from migrating to corn. As soon as the grassy weeds are killed in a field, the larvae move from them to a larger-stemmed plant, such as corn. One method of control is to burn weeds with an herbicide and treat with an insecticide as the larvae move to the corn. Careful timing is essential to achieve control of the borers before they bore into the corn plants. Once the insects are inside the corn plant, they are difficult to control.



Figure 2. Adult Moth

Photo Credit:

University of Illinois, IPM Web site, by Michael Greifenkamp

Insecticide	<u>Chemical Control</u> Rate/Acre	Preharvest Period
Asana XL 0.67	5.8 – 9.6 oz.	21 days
Baythroid 2	1.6 – 2.6 oz.	21 days
Capture 2	2.1 – 6.4 oz.	30 days
Mustang Max 0.8	2.72 – 4.0 oz.	30 days
Pounce 3.2	4 – 8 oz.	30 days
Warrior 1	2.56 – 3.84 oz.	21 days

Precautionary Statement

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store, or dispose of a pesticide. According to laws regula□

Disclaimer Statement

Pesticides recommended in this publication were registered for the prescribed uses when printed. Pesticide registrations are continuously reviewed. Should registration of a recommended pesticide be canceled, it would no longer be recommended by the University of Tennessee. Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others which may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product.

SP431-K (Rev) 4/05 E12-4615 05-0324

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.
UT Extension provides equal opportunities in programs and employment.