



October 2010

# Selected hormonal and neurotransmitter mechanisms regulating

James L. Sartin

*Auburn University Main Campus*

Joseph A. Daniel

*Berry College*

Brian K. Whitlock

*University of Tennessee - Knoxville, bwhitloc@utk.edu*

Robyn R. Wilborn

*Auburn University Main Campus*

Follow this and additional works at: [http://trace.tennessee.edu/utk\\_largpubs](http://trace.tennessee.edu/utk_largpubs)

 Part of the [Animal Sciences Commons](#), [Endocrinology Commons](#), [Large or Food Animal and Equine Medicine Commons](#), [Other Neuroscience and Neurobiology Commons](#), [Reproductive and Urinary Physiology Commons](#), and the [Veterinary Physiology Commons](#)

## Recommended Citation

Sartin, James L.; Daniel, Joseph A.; Whitlock, Brian K.; and Wilborn, Robyn R., "Selected hormonal and neurotransmitter mechanisms regulating" (2010). *Faculty Publications and Other Works -- Large Animal Clinical Sciences*.  
[http://trace.tennessee.edu/utk\\_largpubs/27](http://trace.tennessee.edu/utk_largpubs/27)

This Article is brought to you for free and open access by the Veterinary Medicine -- Faculty Publications and Other Works at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Faculty Publications and Other Works -- Large Animal Clinical Sciences by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

# Selected hormonal and neurotransmitter mechanisms regulating feed intake in sheep

J. L. Sartin<sup>1†</sup>, J. A. Daniel<sup>2</sup>, B. K. Whitlock<sup>3</sup> and R. R. Wilborn<sup>4</sup>

<sup>1</sup>Department of Anatomy, Physiology & Pharmacology, College of Veterinary Medicine, Auburn University, Auburn, AL 36849, USA; <sup>2</sup>Department of Animal Science, Berry College, Mt. Berry, GA 30149, USA; <sup>3</sup>Department of Large Animal Clinical Sciences, College of Veterinary Medicine, The University of Tennessee, Knoxville, TN 37996, USA; <sup>4</sup>Department of Clinical Sciences, College of Veterinary Medicine, Auburn University, Auburn, AL 36849, USA

(Received 1 March 2010; Accepted 25 May 2010; First published online 22 July 2010)

---

*Appetite control is a major issue in normal growth and in suboptimal growth performance settings. A number of hormones, in particular leptin, activate or inhibit orexigenic or anorexigenic neurotransmitters within the arcuate nucleus of the hypothalamus, where feed intake regulation is integrated. Examples of appetite regulatory neurotransmitters are the stimulatory neurotransmitters neuropeptide Y (NPY), agouti-related protein (AgRP), orexin and melanin-concentrating hormone and the inhibitory neurotransmitter, melanocyte-stimulating hormone (MSH). Examination of messenger RNA (using in situ hybridization and real-time PCR) and proteins (using immunohistochemistry) for these neurotransmitters in ruminants has indicated that physiological regulation occurs in response to fasting for several of these critical genes and proteins, especially AgRP and NPY. Moreover, intracerebroventricular injection of each of the four stimulatory neurotransmitters can increase feed intake in sheep and may also regulate either growth hormone, luteinizing hormone, cortisol or other hormones. In contrast, both leptin and MSH are inhibitory to feed intake in ruminants. Interestingly, the natural melanocortin-4 receptor (MC4R) antagonist, AgRP, as well as NPY can prevent the inhibition of feed intake after injection of endotoxin (to model disease suppression of appetite). Thus, knowledge of the mechanisms regulating feed intake in the hypothalamus may lead to mechanisms to increase feed intake in normal growing animals and prevent the wasting effects of severe disease in animals.*

---

**Keywords:** appetite, neuropeptide Y, proopiomelanocortin, leptin, sheep

## Implications

Appetite control by the brain is a critical component of the normal growth and development process as well as of major importance for reproduction. From an understanding of how these mechanisms function, we can develop specific protocols to support appetite in circumstances where feed intake is compromised. For example, development of a melanocortin-4 receptor antagonist that can cross the blood–brain barrier provides the opportunity to increase feed intake in normal animals, as well as to prevent the reduction in feed intake and perhaps body mass wasting in disease conditions. Other areas to consider might be early lactation or in metabolic disease circumstances. Thus, the study of brain control of appetite could have a major impact on improved growth and minimizing losses due to disease or other external stresses.

## Introduction

Appetite control is a complex process whereby multiple stimulatory and inhibitory inputs are integrated in the hypothalamus to yield an increase or decrease in feed intake. Initial lesioning studies in rodents determined a critical role for the arcuate nucleus (ARC) and ventromedial nucleus (VMN) in the control of appetite. Similar studies of lesions to the paraventricular nucleus (PVN) caused hyperphagia, while lesions of the lateral hypothalamic area (LHA) produced anorexia. These studies have been expanded upon to provide a current concept whereby the ARC contains first-order neurons that in turn activate second-order neurons in the LHA to activate feeding of the PVN to inhibit feed intake (Schwartz *et al.*, 2000). These neural pathways have been extensively described for the rodent species (Schwartz *et al.*, 2000; Valassi *et al.*, 2008), and though less studied, are no less important in ruminants. In the ARC, there resides a grouping of neurons that release either orexigenic or anorexigenic neurotransmitters. The orexigenic neurotransmitters found in the ARC are neuropeptide Y (NPY) and agouti-related protein (AgRP),

---

† E-mail: sartijl@auburn.edu

while the primary anorexigenic neurotransmitter found in the ARC is  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH), a product of the proopiomelanocortin (POMC) gene. The second-order neurons are located in the LHA and consist of neurons containing melanin-concentrating hormone (MCH) and orexin (ORX). These appetite-regulating neurons are in turn impacted by a diverse array of modifiers (both stimulatory and inhibitory) ranging from hormones such as leptin, ghrelin and insulin to inflammatory cytokines to nutritional molecules such as glucose, fatty acids, amino acids and volatile fatty acids to nerve impulses from the liver and intestine (for overviews of the rodent literature, see Schwartz *et al.*, 2000; Valassi *et al.*, 2008).

Following an orexigenic stimulus, NPY and AgRP are released from their axon terminals adjacent to second-order neurons in the LHA to activate neurons containing either MCH or cells containing ORX. These neurotransmitters are in turn released from their axon terminals to activate higher order centers to alter feed intake. In the event that the net activation within the ARC is inhibitory to feed intake,  $\alpha$ -MSH is released in the LHA to decrease activation of MCH and ORX and subsequently reduce feed intake (Figure 1). Though a simplistic description of the neural architecture and the process regulating appetite, this provides a framework to test this model in ruminant species.

The remainder of this review will focus on the hypothalamus and the regulation of these pathways in sheep with attention to attempts to activate orexigenic and anorexigenic pathways to influence feed intake.


Studies in sheep indicated that unlike the rodent models, feeding activated c-Fos expression in the supraoptic nucleus (Chaillou *et al.*, 2000). In addition, underfeeding produced c-Fos expression in the PVN, ARC, VMN and dorsomedial nuclei, indicating the importance of these areas in the sheep for appetite control, similar to data in rats.

### Neuropeptide Y

Beginning with the first-order neurons in this pathway, there is a potent orexigenic neurotransmitter, NPY. There is both immunohistochemical evidence for NPY protein expression

and *in situ* hybridization evidence for NPY gene expression localized to the ovine hypothalamus. In colchicine-treated sheep, NPY neurons were found distributed between two populations of cells within the ARC, median eminence, pituitary stalk, the dorsomedian and dorsocaudal nuclei, and in the periventricular nucleus (PeVN; Polkowska *et al.*, 2006a), but were absent in the magnocellular neurons of the paraventricular and supraoptic nuclei (Chaillou *et al.*, 2002; Chaillou and Tillet, 2005). There is also a rich interconnection between the ARC and VMN in sheep (Qi *et al.*, 2008). In addition, numerous studies have demonstrated expression of NPY mRNA in the ARC (Clarke *et al.*, 2000; Adam *et al.*, 2002; Pilon *et al.*, 2003). Interestingly, *in vitro* autoradiography (Williams *et al.*, 1999) and dual label immunohistochemistry (Iqbal *et al.*, 2001b) indicate that leptin receptors are expressed by NPY neurons, suggesting a mechanism for leptin to reduce feed intake in ruminants.

In order to demonstrate a role for NPY as an appetite regulator in ruminants, the effect of alteration in nutritional stimuli should be reflected by changes in NPY neuron activation. For example, reducing caloric intake by fasting should provide a stimulus to NPY. Indeed, feeding a protein-restricted energy balanced diet (8% v. 18% protein) resulted in an increase in immunohistochemical staining for NPY in the PeVN and ARC (Polkowska and Gladysz, 2001). In other studies, negative energy balance created by a 4-day fast resulted in an increase in NPY gene expression in the ARC (Adam *et al.*, 2002) with similar data obtained in castrate male sheep in low body condition (Archer *et al.*, 2002a). In another study comparing lean ewes to fat ewes, the number of neurons positive for NPY was increased by 225% as well as an increase in NPY-positive cells co-expressing leptin receptors (Kurose *et al.*, 2005). A related study employed a daily 4-h leptin infusion for four consecutive days that was found to reduce the immunoreactive staining density for NPY (Polkowska *et al.*, 2006b). Finally, short-day photoperiod exposure in ovariectomized ewes or in rams reduced the number of cells positive for NPY mRNA (by *in situ* hybridization) as well as the number of silver grains per cell, suggesting reduced NPY expression during a period that corresponds to



**Figure 1** Abbreviated outline of the neural pathways regulating appetite in the hypothalamus. Neuropeptide Y (NPY), Agouti-related protein (AGRP), melanin-concentrating hormone (MCH), Orexin (ORX),  $\alpha$ -melanocyte-stimulating hormone MSH, proopiomelanocortin (POMC). NPY, AgRP, ORX, MCH circles represent stimulatory cells and the POMC circles represent inhibitory cells. Reprinted from Sartin *et al.* (2005).

reduced voluntary feed intake (Clarke *et al.*, 2000, 2003). Similar effects were observed with wethers (Dobbins *et al.*, 2004). By contrast, the effect of fasting to increase NPY expression appeared to be restricted to long-day photoperiod exposed sheep (Archer *et al.*, 2004). Moreover, McShane *et al.* (1992) found that undernutrition elevated cerebrospinal concentrations of NPY, consistent with the gene and protein expression data. In a study of negative energy balance produced as a consequence of lactation, Sorenson *et al.* (2002) found that the negative energy balance of lactation was associated with an increase in NPY expression in the ARC and dorsomedial hypothalamus. Thus, the evidence suggests that NPY is present in the area of the hypothalamus that initiates feed intake and is regulated in a manner consistent with expected changes due to an increase or a reduction in calorie intake. In addition to feed intake regulation, the NPY fasting response may correlate to increased growth hormone (GH) and decreased luteinizing hormone (LH). Fasting is a well-known stimulus to GH that can increase the expression of NPY. In an interesting study linking appetite to the growth axis, growth hormone-releasing hormone was found to colocalize with NPY in the ARC during periods of undernutrition (Iqbal *et al.*, 2005). Moreover, NPY is a known stimulus to GH release in sheep (McMahon *et al.*, 1999; Morrison *et al.*, 2003) and cattle (Thomas *et al.*, 1999) and an inhibitor of LH (Morrison *et al.*, 2003), suggesting that the effects of fasting may be linked to GH and LH by way of the NPY neuron.

Fetal sheep also express NPY within the hypothalamus at least by day 110 of gestation, which led to an interesting hypothesis that fetal nutrition may imprint on developing neural circuitry in the hypothalamus and have serious consequences in later life (Mülh usler *et al.*, 2004). The role of fetal NPY was further illustrated by lateral ventricle injections of NPY into fetal sheep, which resulted in increased swallowing behavior (El-Haddad *et al.*, 2003).

For NPY to be considered an appetite regulator, it must also be capable of stimulating an increase in feed intake in *ad libitum* fed sheep. Indeed, intracerebroventricular (ICV) injection of NPY in sheep resulted in a pronounced increase in feed intake (Miner *et al.*, 1989; Sartin *et al.*, 2001; Wagner *et al.*, 2004; Whitlock *et al.*, 2005). An interesting study of the NPY receptor types mediating feed intake (Clarke *et al.*, 2005) indicated that the major receptor mediating the increased intake due to NPY was the NPY-Y1 receptor, and to a lesser degree, the Y2 receptor. This effect of NPY to increase feed intake was also significant in the presence of rumen distention or propionate infusion (Miner *et al.*, 1990). In a related study, endotoxin was used to inhibit feed intake in sheep (as a model for disease) and was coupled with NPY infusion ICV (McMahon *et al.*, 1999). The infusion of NPY in continually endotoxin-treated sheep normalized feed intake during the period of infusion. More importantly, feed intake remained at normal levels even after the removal of NPY. Therefore, NPY has been characterized as a neurotransmitter of the appetite system that regulated increased feed intake and is in turn regulated by nutritional as well as photoperiod manipulations in sheep.

#### *Agouti-related protein*

AgRP mRNA (Henry *et al.*, 2001a; Adam *et al.*, 2002; Archer *et al.*, 2002a) and AgRP protein (Wagner *et al.*, 2004) were localized to the ARC nucleus in sheep. Cloning of a partial sequence of AgRP mRNA indicated a high homology with bovine AgRP but the comparison of the human message indicated only about 56% homology for the portion cloned (Wagner *et al.*, 2004). Upregulation of AgRP mRNA was observed in chronically underfed ewes (Henry *et al.*, 2001a; Archer *et al.*, 2004 and 2005; Relling *et al.*, 2010) or castrate male sheep fasted for 4 days (Adam *et al.*, 2002). In addition, dual label immunohistochemistry (for AgRP and c-Fos) indicated that fasting for 3 days increased the number of AgRP-positive neurons and the activation of AgRP neurons in the ARC (percentage of AgRP neurons also expressing c-Fos; Wagner *et al.*, 2004). Gene expression for AgRP was also increased in sheep with a low body condition score compared to sheep with a high body condition score (Archer *et al.*, 2004). Interestingly, feed restriction did not alter mRNA expression for the receptor targets of AgRP (Iqbal *et al.*, 2001a), the melanocortin-3 receptor (MC3R) and the MC4R. Similarly, lambs born from ewes fed 160% of metabolizable energy requirements had more body fat and a lower expression of AgRP message (M ulh usler *et al.*, 2006). The infusion of glucose into fetal sheep in late gestation (day 130 to 140) did not affect the expression of AgRP in the fetal ARC, though there were increases in body fat in the glucose-infused sheep (M ulh usler *et al.*, 2005). These studies reflect a functioning appetite regulatory system in the fetus and lend support to the hypothesis that fetal nutrient exposure may have an outcome in postnatal life.

The effects of AgRP on feed intake are thought to be mediated largely via the MC4R, where AgRP inhibits the activity of the receptor to prevent  $\alpha$ -MSH from activating the receptor and inhibiting appetite (Ellacott and Cone, 2004). Thus, AgRP increases appetite by inhibition of a receptor that inhibits appetite (or an inverse agonist). Injection of AgRP into the lateral ventricle of *ad libitum* fed sheep resulted in an increase in cumulative feed intake. The effect was present between 4 and 12 h, but unlike other animal models, the effect on feed intake did not persist to 24 h (Wagner *et al.*, 2004). Examination of whether altered AgRP expression contributed to reduced feed intake after endotoxin injection determined that there were no changes in AgRP expression during the early phase of appetite suppression (Sartin *et al.*, 2008). In another experiment, AgRP injected before endotoxin prevented the inhibition of appetite (Sartin *et al.*, 2008) characteristic of endotoxin treatment. This suggests that MC4R is a major component of the mechanism for inhibition of feed intake in diseased ruminants and suggests a possible means of reversing the negative impact of disease on feeding and perhaps on subsequent body wasting associated with disease (Sartin *et al.*, 2005).

AgRP is a potent neurotransmitter regulating feed intake in sheep. The synthesis of AgRP is regulated by nutritionally relevant stimuli and AgRP can activate feeding in sheep that are satiated. Thus, like NPY, this neurotransmitter is clearly

an appetite regulator in sheep, and perhaps may provide a unique opportunity to manipulate appetite in disease, stress and/or other metabolic problems related to a reduction in feed intake.

#### *Melanin-concentrating hormone*

The expression of MCH is pronounced in the lateral hypothalamic area, based on mRNA expression (Henry *et al.*, 2000) and immunohistochemical data evidence of the MCH protein (Tillet *et al.*, 1996; Chaillou *et al.*, 2003; Whitlock *et al.*, 2005). Cloning of a partial ovine MCH sequence indicates a high homology to other species (Whitlock *et al.*, 2005). While localization is similar to other species, there are some minor differences in localization (Tillet *et al.*, 1996). In sheep, MCH has been found to be localized to the supramammillary nucleus, nucleus medialis thalami and nucleus reunions (Tillet *et al.*, 1996). Of further interest are studies which indicate that MCH neurons from the LHA provide input to the ARC and VMN (Qi *et al.*, 2008). This adds a possible layer of reciprocal control of appetite, metabolic and endocrine control within the hypothalamus.

Comparison of gene expression for MCH in thin *v.* fat ewes (Henry *et al.*, 2000; Anukulkitch *et al.*, 2009) indicated more mRNA expression for MCH in thin ewes (as well as lower insulin, lactate and free fatty acids in thin ewes), while neurotransmitter gene expression in first-order neurons (NPY and POMC) was unaffected. Moreover, leptin receptors were expressed in all MCH neurons in the LHA (Iqbal *et al.*, 2001b), though in the absence of the ARC there was no effect of leptin injection on MCH neuron activation (as measured by *c-Fos* staining; Qi *et al.*, 2010). However, even though the presence of leptin receptors suggests that effects of fasting might regulate MCH neurons, undernutrition did not alter the number of MCH-positive neurons as examined by immunohistochemistry (Chaillou *et al.*, 2003). Similarly, short-term fasting of 3 days also had no effect on MCH mRNA in the whole hypothalamus or on the number of MCH-positive neurons in the LHA or on the percentage of neurons co-expressing *c-Fos* with MCH, suggesting that these MCH neurons were not affected by the nutritional stimulus (Whitlock *et al.*, 2005). These studies contrast with studies in laboratory animals that suggest that fasting increases MCH (Schwartz *et al.*, 2000).

The physiological effects of MCH have also been examined. Parkes (1996) infused MCH into the lateral ventricle of the sheep brain for 24 h. The infusion of MCH increased serum osmolality, urine volume, urine sodium and potassium concentrations. Examination of appropriate endocrine responses showed that MCH increased serum glucose, decreased plasma aldosterone concentrations and had no effect on serum protein, adrenocorticotropin, cortisol, vasopressin, renin, endothelin or atrial natriuretic peptide. These changes suggest that MCH may regulate both diuretic and natriuretic functions in sheep (Parkes, 1996). Finally, MCH injection into the lateral ventricle of the brain increased feed intake, but there was no dose-response effect. A similar increase in feed intake was

found between a single injection of MCH into the ventricle of the brain and a molar-equivalent dose of NPY (Whitlock *et al.*, 2005). However, infusion of MCH did not increase feed intake, though an infusion of NPY produced a significant increase in feed intake, thus indicating that MCH regulates feed intake as predicted by the model and suggests a possible down-regulation of the MCH receptor when confronted with prolonged high levels of MCH. Therefore, MCH appears to be a very potent regulator of appetite in sheep, though evidence for interactions with endocrine or metabolic systems have not been extensively examined in sheep.

#### *Orexin*

ORX expression has been demonstrated in neurons of the LHA (Iqbal *et al.*, 2001c; Qi *et al.*, 2008) as well as the dorsomedial nucleus, zona incerta, perifornical area and anterior hypothalamic area (Iqbal *et al.*, 2001c). The receptor for orexin (OX1R) was localized to the ARC, median eminence, lateral hypothalamic nuclei and ventral portion of the preoptic area (Zhang *et al.*, 2005). Interestingly, ORX neurons originating in the LHA provide input to both the ARC and VMN (Qi *et al.*, 2008). Combined, these studies link ORX to regions of the hypothalamus related to feed intake, endocrine, metabolic and reproductive regulation.

Use of nutritional manipulation by chronic undernutrition *v.* controls indicated no effect on ORX gene expression (Iqbal *et al.*, 2003). However, ORX gene expression was negatively associated with adiposity in sheep selected for fatness or lean body condition (Anukulkitch *et al.*, 2010). ORX gene expression was enhanced in short-day photoperiod when appetite is typically lower (Archer *et al.*, 2002b). Similarly, following sheep throughout periods of seasonal appetite changes again demonstrated a negative correlation of ORX to adiposity (Anukulkitch *et al.*, 2010). Related is the demonstration that 100% of ORX neurons express the leptin receptor, suggesting a means of communicating nutritional status to the appetite centers of the hypothalamus to integrate appetite signals (Iqbal *et al.*, 2001c). Moreover, leptin activation of *c-Fos* in ORX neurons was demonstrated in the absence of the ARC, suggesting a direct action of leptin on second-order neurons in sheep (Qi *et al.*, 2010), which differs slightly with the model proposed for laboratory animals.

ORX is related to the regulation of feed intake and, indeed, injection of ORX B into the lateral ventricles of the sheep brain (but not IV injection) enhanced short-term feed intake (Sartin *et al.*, 2001). There have also been hypotheses of a relationship to hypothalamic control of reproduction, where ORX neurons terminate close to gonadotropin releasing hormone neurons (Iqbal *et al.*, 2001c; Qi *et al.*, 2010). In addition to reproduction, ORX neurons project to the PeVN adjacent to somatostatin neurons (Iqbal *et al.*, 2005), suggesting a possible role in regulating GH. Indeed, ORX A and B receptors are expressed in somatotropes and both molecules activate calcium channels in the pituitary via protein kinase C mechanisms (Xu *et al.*, 2002 and 2003). ORX B will also release GH from isolated pituitary cells. Interestingly,

there was an increase in plasma cortisol, along with an anecdotal report of increased water intake. However, ICV injection of ORX B in sheep had no effect on plasma LH, GH or insulin concentrations (Sartin *et al.*, 2001), though perhaps in steroid-treated or photoperiod-manipulated sheep, there may have been a different outcome. Thus, ORX also appears to be a molecule that cannot only enhance feed intake, but may also integrate appetite control with reproduction and metabolic endocrinology in sheep.

#### *Proopiomelanocortin/α-melanocyte-stimulating hormone*

Based on the appetite model in Figure 1, neurons containing POMC should be localized to the ARC. However, it must be emphasized that POMC expression does not always correlate to POMC neurons regulating appetite. In sheep, *in situ* hybridization studies clearly localize POMC to the ARC (Henry *et al.*, 2000; Clarke *et al.*, 2000; Lincoln *et al.*, 2001; Adam *et al.*, 2002). In addition, immunohistochemical evidence for  $\alpha$ -MSH (product of the POMC gene) was likewise found in the ARC (Sartin *et al.*, 2008). In one study, POMC gene expression in the ARC was found to be unaffected by a short-term (4-day) fast (Adam *et al.*, 2002). In a study of feed intake and adiposity, castrate male sheep with estradiol implants were fed (4 weeks) to maintain body condition or to produce low or high body condition (Archer *et al.*, 2002a). The expression of POMC was similar across all groups. In another study of long-term changes in nutrition, there were no differences in thin *v.* fat ewes in terms of POMC gene expression (Henry *et al.*, 2000). And finally, another 4-week trial found no effects of feed restriction on POMC gene expression (Relling *et al.*, 2010). By contrast, McShane *et al.* (1993) observed a 52% decrease in POMC expression with a long-term feed restriction in ewes (restricted ewes were fed 30% of requirements for 7 weeks), so that with sufficient energy reductions it was possible to impact POMC. Based on these data, it is interesting to note that an effect of fasting to reduce POMC gene expression is a more common finding in laboratory animals as compared to the minimal effects on POMC expression in sheep. This may indicate that the grazing animal has developed a different strategy to regulate the inhibition of appetite compared to intermittent eaters such as the rodent.

Additional studies have examined photoperiod and the interaction between photoperiod and nutritional effects on expression of the POMC gene. Clarke *et al.* (2000) compared variations in POMC gene expression with voluntary feed intake and natural photoperiod. In the ARC, NPY, and feed intake changed in the opposite direction to POMC. In another study contrasting long-day photoperiod and short-day photoperiod, Clarke *et al.* (2003) found that there was a decrease in gene expression for POMC in the medial ARC in long-day photoperiod exposed rams. Anukulkitch *et al.* (2009) also found that POMC gene expression in ewes was reduced during long-day photoperiod exposure, the opposite of that seen with NPY gene expression. Castrate rams supplemented with testosterone were also found to have a reduced POMC gene expression during long-day photoperiods (Hileman *et al.*, 1998).

A photoperiod study conducted for 11 weeks (Archer *et al.*, 2004) found no effect of photoperiod alone on POMC gene expression in castrate male sheep supplemented with estradiol and either *ad libitum* fed or fasted. However, when intact rams were switched from a long- to a short-day photoperiod, POMC gene expression was increased. When this photoperiod strategy was combined with feed restriction, POMC gene expression was reduced in the short-day photoperiod group. Thus, POMC neuron regulation in the sheep requires both the appropriate photoperiod and reproductive steroid background as well as nutritional changes before regulation of POMC neurons was consistently observed. Moreover, POMC is a precursor to multiple products, which makes interpretation of general observations difficult in the absence of knowledge of the protein products coupled to the POMC expression. This may, in part, explain the variable POMC response to fasting observed in this species.

Fetal sheep also have POMC gene expression in the ARC (Mülhäusler *et al.*, 2004) in mid- to late gestation. Glucose infusion to late gestation fetal sheep increased fat mass in the sheep, accompanied by increased POMC gene expression (Mülhäusler *et al.*, 2005). Similar increases in fetal POMC expression were obtained by increased nutrition delivered to the mother, which was related to elevated fetal glucose levels (Adam *et al.*, 2006). These studies imply that maternal nutrition may program later postnatal development within the appetite regulatory pathway. Indeed, increased maternal nutrition (160% of controls) was found to result in increased plasma glucose and body fat after birth and this was accompanied by increased POMC expression (Mülhäusler *et al.*, 2006), thus providing direct evidence for fetal environment having a role in programming of appetite regulatory genes in postnatal life.

Effects of treatments on  $\alpha$ -MSH (protein product of the POMC gene) synthesis or release or effects of treatment with  $\alpha$ -MSH have been less studied in sheep. In one study, injection of a melanocortin agonist, MTII (which activates the same receptor as  $\alpha$ -MSH), upregulated kisspeptin and LH release (Backholer *et al.*, 2009). In another study,  $\alpha$ -MSH immunohistochemistry indicated that with time after endotoxin injection and associated with prolonged reductions (6 h) in feed intake, there is a reduction in  $\alpha$ -MSH staining and a reduction in dual labeling for c-Fos, suggesting that  $\alpha$ -MSH neurons were not being activated under these circumstances (Sartin *et al.*, 2008). This reduced expression of POMC and  $\alpha$ -MSH neuron activation was accompanied by an increase in AgRP gene expression and c-Fos staining. These changes at 6 h are consistent with an increase in appetite stimulation and correlate to the time when sheep first eat following endotoxin injection. While speculative at this time, perhaps these changes may account for the subsequent compensatory feeding that occurs between 24 and 48 h when cumulative feed intake between the controls and endotoxin-treated sheep no longer differs.

The POMC neuron is typically activated with feeding and inhibited by fasting. This is further complemented by the opposing changes in the endogenous MC4R antagonist,

AgRP. An elevation in AgRP inhibits MC4R activity and prevents feed intake inhibition. Thus, the POMC neuron represents the initiation of a potent inhibitory system regulating appetite, though in sheep the POMC response is influenced by photoperiod and steroid exposure.

A summary of neurotransmitters and appetite control in sheep is shown in Table 1.

### Leptin

Leptin is a product of the fat cell and is transported from the blood to the cerebrospinal fluid (CSF) to inhibit NPY and AgRP neurons, which would result in reduced feed intake. Leptin will also activate POMC neurons to inhibit feed intake. In ruminants, leptin gene expression and immunohistochemistry indicate that leptin is synthesized by the fat cell (Dyer *et al.*, 1997; Kumar *et al.*, 1998; Daniel *et al.*, 2003). Circulating leptin concentrations are episodic in nature and are increased by obesity and decreased by fasting, though there were no apparent photoperiod effects on plasma leptin concentrations (Kumar *et al.*, 1998; Marie *et al.*, 2001; Daniel *et al.*, 2002; Ehrhardt *et al.*, 2003; Altmann *et al.*, 2005; Chilliard *et al.*, 2005; Delavaud *et al.*, 2007). Feeding high fat or carbohydrate meals resulted in increased plasma leptin concentrations (Yildiz *et al.*, 2003) suggesting that higher energy content rather than the source of energy was important. After release from the adipocyte, leptin crosses the blood–brain barrier to enter the hypothalamus as indicated by its entry into the CSF compartment (Thomas *et al.*, 2001; Adam *et al.*, 2006), though amounts of leptin entering the CSF were greater in long-day as opposed to short-day photoperiods (Adam *et al.*, 2006). Interestingly, plasma and CSF levels only correlated in long-day photoperiod exposed sheep, suggesting a photoperiod influence on leptin transport that corresponds to reported sensitivity in POMC gene expression to photoperiod (Clarke *et al.*, 2003; Archer *et al.*, 2004; Anukulkitch *et al.*, 2009).

Treatment of sheep with ICV but not IV leptin resulted in an inhibition of feed intake (Henry *et al.*, 1999 and 2001b; Clarke *et al.*, 2001; Morrison *et al.*, 2001 and 2002; Miller *et al.*, 2002 and 2007), though the effect could be related to the time of year and whether the sheep were well fed or fasted (thin body condition; Clarke *et al.*, 2001). Interestingly, some of the leptin receptors in the ARC were found to be localized to NPY neurons (Williams *et al.*, 1999; Kurose *et al.*, 2005)

and leptin decreased NPY gene expression (Henry *et al.*, 1999) and immunostaining (Polkowska *et al.*, 2006b) in the ARC. There was an increase in the number of NPY neurons expressing leptin receptors in lean sheep (Kurose *et al.*, 2005) and leptin receptor gene expression was increased in long-day exposed sheep (Clarke *et al.*, 2003). Thus, the evidence suggests that leptin inhibits feed intake, in part by reducing NPY expression as suggested by the model (Figure 1). Interestingly, unlike studies in rodent models, lesions of the ARC do not fully remove the inhibitory effects of leptin in sheep (Qi *et al.*, 2010). In this model, leptin increased c-Fos expression in the dorsomedial nucleus, VMN and PVN. While lesions of the ARC prevented c-Fos labeling in MCH neurons following leptin treatment, ORX neurons in the lateral hypothalamus and dorsomedial nucleus were still evident. These data provide evidence that all actions of leptin are not mediated exclusively through first-order to second-order neurons in sheep (Qi *et al.*, 2010).

Fetal sheep also have circulating leptin, which is regulated in the fetus (McMillen *et al.*, 2006). With high nutrient intake in the pregnant ewe, there is an expected increase in plasma leptin. However, the increased leptin is not reflected in fetal circulating leptin (Mülhäusler *et al.*, 2006).

Inhibition of feed intake is a hallmark of disease, leading to the hypothesis that an increase in leptin during disease might provide a mechanism for the disease-associated reduction in feed intake. However, in a well-characterized endotoxin injection model of disease in sheep, there was no evidence of an increase in circulating leptin, in spite of significant reductions in feed intake (Soliman *et al.*, 2001; Daniel *et al.*, 2003), though tumor necrosis factor and the endotoxin receptor (CD14) were both found to be expressed in the ovine adipocyte (Daniel *et al.*, 2003) as expected. In parasitized lambs, one study suggested that leptin might be a partial factor in reduced feed intake (Zaralis *et al.*, 2008), though in other studies, plasma leptin concentrations were unchanged by the infection (Greer *et al.*, 2009; Zaralis *et al.*, 2009). However, no studies have examined the effects of disease on CSF leptin levels, which could change by altered uptake of leptin independently of changes in plasma levels. While leptin may have a role in disease suppression of appetite in other species, the evidence is lacking in sheep, suggesting that some other mechanism may be in place to mediate peripheral disease effects on hypothalamic neurons regulating appetite.

**Table 1** Summary of neurotransmitters and appetite control in sheep

Neurotransmitter	Abbreviation	Feed intake effect in sheep	Effect of fasting on gene expression	Leptin receptor/effect of leptin on gene expression
$\alpha$ -melanocyte-stimulating hormone	$\alpha$ -MSH	?	No change or decrease	Yes/increase
Neuropeptide Y	NPY	Increase	Increase	Yes/decrease
Agouti-related protein	AgRP	Increase	Increase	?
Melanin-concentrating hormone	MCH	Increase	No change	Yes/?
Orexin	ORX	Increase	Increase	Yes/?

? = information not available.

## Conclusion

This review did not make an attempt to contrast the sheep and other models of appetite control (for these contrasts, see Clarke, 2008). It should be noted that as a grazing animal with continuous nutrient absorption, the nutritional demands differ with regard to intermittent eaters. Indeed, there are key differences in these animal models, especially with regard to factors such as photoperiod, on the one hand, and GH regulation by NPY on the other. The expression of NPY, AgRP, MCH and ORX was found in the proper locations in the hypothalamus for appetite control. Moreover, first-order neurons (NPY and AgRP) are responsive to altered nutrition and all four of these key neurotransmitters increase feed intake. Thus, in general terms, the simple model for neurotransmitter regulation of appetite in Figure 1 is a reasonable fit to use for framing hypotheses in ruminant species. However, studies such as those providing the indication that leptin inhibition of appetite was not exclusively transmitted through first-order neurons as well as the lowered sensitivity of POMC neurons (unless modified by photoperiod) for activation indicates definitive species differences with laboratory animals and indicates a clear need for further study of appetite in farm animal species.

## Acknowledgments

This work is based on an invited presentation at the *60th Annual Meeting of the European Association for Animal Production* held in Barcelona, Spain during August 2009. This project was supported by National Research Initiative Competitive Grant no. 2004-35206-14136 from the USDA Cooperative State Research, Education and Extension Service.

## References

Adam CL, Findlay PA and Miller DW 2006. Blood-brain leptin transport and appetite and reproductive neuroendocrine responses to intracerebroventricular leptin injection in sheep: influence of photoperiod. *Endocrinology* 147, 4589–4598.

Adam CL, Archer ZA, Findlay PA, Thomas L and Marie M 2002. Hypothalamic gene expression in sheep for cocaine- and amphetamine-regulated transcript, pro-opiomelanocortin, neuropeptide Y, agouti-related peptide and leptin receptor and responses to negative energy balance. *Neuroendocrinology* 75, 250–256.

Altmann M, Sauerwein H and Borell E 2005. Relationship between plasma leptin concentrations and carcass composition in fattening mutton: a comparison with ultrasound results. *Journal of Animal Physiology and Animal Nutrition (Berlin)* 89, 326–330.

Anukulkitch C, Rao A, Dunchea FR and Clarke IJ 2009. A test of the lipostat theory in a seasonal (ovine) model under natural conditions reveals a close relationship between adiposity and melanin concentrating hormone expression. *Domestic Animal Endocrinology* 36, 138–151.

Anukulkitch C, Rao A, Periera A, McEwan J and Clarke IJ 2010. Expression of genes for appetite-regulating peptides in the hypothalamus of genetically selected lean and fat sheep. *Neuroendocrinology* 91, 223–238. (doi: 10.1159/000262283).

Archer ZA, Rhind SM, Findlay PA, Kyle CE, Thomas L, Marie M and Adam CL 2002a. Contrasting effects of different levels of food intake and adiposity on LH secretion and hypothalamic gene expression in sheep. *Journal of Endocrinology* 175, 383–393.

Archer ZA, Findlay PA, Rhind SM, Mercer JG and Adam CL 2002b. Orexin gene expression and regulation by photoperiod in the sheep hypothalamus. *Regulatory Peptides* 104, 41–45.

Archer ZA, Findlay PA, McMillen SR, Rhind SM and Adam CL 2004. Effects of nutritional status and gonadal steroids on expression of appetite-regulatory genes in the hypothalamic arcuate nucleus of sheep. *Journal of Endocrinology* 182, 409–419.

Archer ZA, Rhind SM, Findlay PA, Kyle CE, Barber MC and Adam CL 2005. Hypothalamic responses to peripheral glucose infusion in food-restricted sheep are influenced by photoperiod. *Journal of Endocrinology* 184, 515–525.

Backholer K, Smith J and Clarke IJ 2009. Melanocortins may stimulate reproduction by activating orexin neurons in the dorsomedial hypothalamus and kisspeptin neurons in the preoptic area of the ewe. *Endocrinology* 150, 5488–5497.

Chaillou E and Tillet Y 2005. Nutrition and hypothalamic neuropeptides in sheep: histochemical studies. *Histology and Histopathology* 20, 1209–1225.

Chaillou E, Baumont R, Tramu G and Tillet Y 2000. Effect of feeding on Fos protein expression in sheep hypothalamus with special reference to the supraoptic and paraventricular nuclei: an immunohistochemical study. *European Journal of Neuroscience* 12, 4515–4524.

Chaillou E, Baumont R, Chilliard Y and Tillet Y 2002. Several subpopulations of neuropeptide Y-containing neurons exist in the infundibular nucleus of sheep: an immunohistochemical study of animals on different diets. *Journal of Comparative Neurology* 444, 129–143.

Chaillou E, Baumont R, Fellman D, Tramu G and Tillet Y 2003. Sensitivity of galanin- and melanin concentrating hormone-containing neurons to nutritional status: an immunohistochemical study in the ovariectomized ewe. *Journal of Neuroendocrinology* 15, 459–467.

Chilliard Y, Delavaud C and Bonnet M 2005. Leptin expression in ruminants: nutritional and physiological regulation in relation with energy metabolism. *Domestic Animal Endocrinology* 29, 3–22.

Clarke IJ 2008. Models of 'obesity' in large animals and birds. *Frontiers of Hormone Research (Karger, Basel)* 36, 107–117.

Clarke IJ, Backholer K and Tilbrook AJ 2005. Y2 receptor-selective agonist delays the estrogen-induced luteinizing hormone surge in ovariectomized ewes, but Y1-receptor-selective agonist stimulates voluntary food intake. *Endocrinology* 146, 769–775.

Clarke IJ, Scott CJ, Rao A, Pompolo S and Baker-Gibb ML 2000. Seasonal changes in the expression of neuropeptide Y and pro-opiomelanocortin mRNA in the arcuate nucleus of the ovariectomized ewe: relationship to the seasonal appetite and breeding cycles. *Journal of Neuroendocrinology* 12, 1105–1111.

Clarke IJ, Tilbrook AJ, Turner AI, Doughton BW and Goding JW 2001. Sex, fat and the tilt of the earth: effects of sex and season on the feeding response to centrally administered leptin in sheep. *Endocrinology* 142, 2725–2728.

Clarke IJ, Rao A, Chilliard Y, Delavaud C and Lincoln GA 2003. Photoperiod effects on gene expression for hypothalamic appetite-regulating peptides and food intake in the ram. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 284, R101–R115.

Daniel JA, Whitlock BK, Baker JA, Steele BP, Morrison CD, Keisler DH and Sartin JL 2002. Effect of body fat mass and nutritional status on 24-hour leptin profiles in ewes. *Journal of Animal Science* 80, 1083–1089.

Daniel JA, Elsasser TH, Morrison CD, Keisler DH, Whitlock BK, Steele BP, Pugh D and Sartin JL 2003. Leptin, tumor necrosis factor- $\alpha$  (TNF), and CD14 in ovine adipose tissue and changes in circulating TNF in lean and fat sheep. *Journal of Animal Science* 81, 2590–2599.

Delavaud C, Bocquier F, Baumont R, Chaillou E, Ban-Tokuda T and Chilliard Y 2007. Body fat content and feeding level interact strongly in the short- and medium-term regulation of plasma leptin during underfeeding and re-feeding in adult sheep. *British Journal of Nutrition* 98, 106–115.

Dobbins A, Lubbers LS, Jackson GL, Kuehl DE and Hileman SM 2004. Neuropeptide Y gene expression in male sheep: influence of photoperiod and testosterone. *Neuroendocrinology* 79, 82–89.

Dyer CJ, Simmons JM, Matteri RL and Keisler DH 1997. cDNA cloning and tissue-specific gene expression of ovine leptin, NPY-Y1 receptor, and NPY-Y2 receptor. *Domestic Animal Endocrinology* 14, 295–303.

Ehrhardt RA, Greenwood PL, Bell AW and Boisclair YR 2003. Plasma leptin is regulated predominantly by nutrition in preruminant lambs. *Journal of Nutrition* 133, 4196–4201.

- El-Haddad MA, Ismail Y, Guerra C, Day L and Ross MG 2003. Neuropeptide Y administration into cerebral ventricles stimulates sucrose ingestion in the near-term ovine fetus. *American Journal of Obstetrics and Gynecology* 189, 949–952.
- Ellacott KL and Cone RD 2004. The central melanocortin system and the integration of short- and long-term regulators of energy homeostasis. *Recent Progress in Hormone Research* 59, 395–408.
- Greer AW, Boisclair YR, Stankiewicz M, McNulty RW, Jay NP and Sykes AR 2009. Leptin concentrations and the immune-mediated reduction of feed intake in sheep infected with nematode *Trichostrongylus colubriformis*. *British Journal of Nutrition* 102, 954–957.
- Henry BA, Rao A, Ikenasio BA, Mountjoy KG, Tilbrook AJ and Clarke IJ 2001a. Differential expression of cocaine- and amphetamine-regulated transcript and agouti related-protein in chronically food-restricted sheep. *Brain Research* 918, 40–50.
- Henry BA, Goding JW, Tilbrook AJ, Dunshea FR and Clarke IJ 2001b. Intracerebroventricular infusion of leptin elevates the secretion of luteinizing hormone without affecting food intake in long-term food-restricted sheep, but increases growth hormone irrespective of bodyweight. *Journal of Endocrinology* 168, 67–77.
- Henry BA, Tilbrook AJ, Dunshea FR, Rao A, Blache D, Martin GB and Clarke IJ 2000. Long-term alterations in adiposity affect the expression of melanin-concentrating hormone and enkephalin but not proopiomelanocortin in the hypothalamus of ovariectomized ewes. *Endocrinology* 141, 1506–1514.
- Henry BA, Goding JW, Alexander WS, Tilbrook AJ, Canny BJ, Dunshea FR, Rao A, Mansell A and Clarke IJ 1999. Central administration of leptin to ovariectomized ewes inhibits food intake without affecting the secretion of hormones from the pituitary gland: evidence for a dissociation of effects on appetite and neuroendocrine function. *Endocrinology* 140, 1175–1182.
- Hileman SM, Kuehl DE and Jackson GL 1998. Photoperiod affects the ability of testosterone to alter proopiomelanocortin mRNA, but not luteinizing hormone-releasing hormone mRNA, levels in male sheep. *Journal of Neuroendocrinology* 10, 587–592.
- Iqbal J, Manley TR, Ciofi P and Clarke IJ 2005. Reduction in adiposity affects the extent of afferent projections to growth hormone releasing hormone and somatostatin neurons and the degree of colocalization of neuropeptides in growth hormone-releasing hormone and somatostatin cells in the ovine hypothalamus. *Endocrinology* 146, 4776–4785.
- Iqbal J, Henry BA, Pompolo S, Rao A and Clarke IJ 2003. Long-term alterations in bodyweight and food restriction does not affect gene expression of either preproorexin or prodynorphin in the sheep. *Neuroscience* 118, 217–226.
- Iqbal J, Pompolo S, Dumont LM, Wu CS, Mountjoy KG, Henry BA and Clarke IJ 2001a. Long-term alterations in body weight do not affect the expression of melanocortin receptor-3 and -4 mRNA in the ovine hypothalamus. *Neuroscience* 105, 931–940.
- Iqbal J, Pompolo S, Murakami T, Grouzmann E, Sakurai T, Meister B and Clarke IJ 2001b. Immunohistochemical characterization of localization of long-form leptin receptor (OB-Rb) in neurochemically defined cells in the ovine hypothalamus. *Brain Research* 920, 55–64.
- Iqbal J, Pompolo S, Sakurai T and Clarke IJ 2001c. Evidence that orexin-containing neurons provide direct input to gonadotropin-releasing hormone neurons in the ovine hypothalamus. *Journal of Neuroendocrinology* 13, 1033–1041.
- Kumar B, Francis SM, Suttie JM and Thompson MP 1998. Expression of obese mRNA in genetically lean and fat selection lines of sheep. *Comparative Biochemistry and Physiology – Part B, Biochemistry and Molecular Biology* 120, 543–548.
- Kurose Y, Iqbal J, Rao A, Murata Y, Hasegawa Y, Terashima Y, Kojima M, Kangawa K and Clarke IJ 2005. Changes in expression of the genes for the leptin receptor and the growth hormone-releasing peptide/ghrelin receptor in the hypothalamic arcuate nucleus with long-term manipulation of adiposity by dietary means. *Journal of Neuroendocrinology* 17, 331–340.
- Lincoln GA, Rhind SM, Pompolo S and Clarke IJ 2001. Hypothalamic control of photoperiod-induced cycles in food intake, body weight, and metabolic hormones in rams. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 281, R76–R90.
- Marie M, Findlay PA, Thomas L and Adam CL 2001. Daily patterns of plasma leptin in sheep: effects of photoperiod and food intake. *Journal of Endocrinology* 170, 277–286.
- McMahon CD, Buxton DF, Elsasser TH, Gunter DR, Sanders LG, Steele BP and Sartin JL 1999. Neuropeptide Y restores appetite and alters concentrations of GH after central administration to endotoxic sheep. *Journal of Endocrinology* 161, 333–339.
- McMillen IC, Edwards LJ, Duffiels J and Mülhäusler BS 2006. Regulation of leptin synthesis and secretion before birth: implications for early programming of adult obesity. *Reproduction* 131, 415–427.
- McShane TM, May T, Miner JL and Keisler DH 1992. Central actions of neuropeptide-Y may provide a neuromodulatory link between nutrition and reproduction. *Biology of Reproduction* 46, 1151–1157.
- McShane TM, Petersen SL, McCrone S and Keisler DH 1993. Influence of food restriction on neuropeptide-Y, proopiomelanocortin, and luteinizing hormone-releasing hormone gene expression in sheep hypothalamus. *Biology of Reproduction* 49, 831–839.
- Miller DW, Findlay PA, Morrison MA, Raver N and Adam CL 2002. Seasonal and dose-dependent effects of intracerebroventricular leptin on 1 h secretion and appetite in sheep. *Journal of Endocrinology* 175, 395–404.
- Miller DW, Harrison JL, Bennett EJ, Findlay PA and Adam CL 2007. Nutritional influences on reproductive neuroendocrine output: insulin, leptin, and orexigenic neuropeptide signaling in the ovine hypothalamus. *Endocrinology* 148, 5313–5322.
- Miner JL, Della-Fera MA and Paterson JA 1990. Blockade of satiety factors by central injection of neuropeptide Y in sheep. *Journal of Animal Science* 68, 3805–3811.
- Miner JL, Della-Fera MA, Paterson JA and Baile CA 1989. Lateral cerebroventricular injection of neuropeptide Y stimulates feeding in sheep. *American Journal of Physiology* 257, R383–R387.
- Morrison CD, Wood R, McFadin EL, Whiley NC and Keisler DH 2002. Effect of intravenous infusion of recombinant ovine leptin on feed intake and serum concentrations of GH, LH, insulin, IGF-1, cortisol, and thyroxine in growing prepubertal ewe lambs. *Domestic Animal Endocrinology* 22, 103–112.
- Morrison CD, Daniel JA, Holmberg BJ, Djiane J, Raver N, Gertler A and Keisler DH 2001. Central infusion of leptin into well-fed and undernourished ewe lambs: effects on feed intake and serum concentrations of growth hormone and luteinizing hormone. *Journal of Endocrinology* 168, 317–324.
- Morrison CD, Daniel JA, Hampton JH, Buff PR, McShane TM, Thomas MG and Keisler DH 2003. Luteinizing hormone and growth hormone secretion in ewes infused intracerebroventricularly with neuropeptide Y. *Domestic Animal Endocrinology* 24, 69–80.
- Mülhäusler BS, Adam CL, Findlay PA, Duffield JA and McMillen IC 2006. Increased maternal nutrition alters development of the appetite regulating network in the brain. *FASEB Journal* 20, 1257–1259.
- Mülhäusler BS, McMillen IC, Rouzard G, Findlay PA, Marrocco EM, Rhind SM and Adam CL 2004. Appetite regulatory neuropeptides are expressed in the sheep hypothalamus before birth. *Journal of Neuroendocrinology* 16, 502–507.
- Mülhäusler BS, Adam CL, Marrocco EM, Findlay PA, Roberts CT, McFarlane JR, Kauter KG and McMillen IC 2005. Impact of glucose infusion on the structural and functional characteristics of adipose tissue and on hypothalamic gene expression for appetite regulatory neuropeptides in the sheep fetus during late gestation. *Journal of Physiology* 565, 185–195.
- Parkes DG 1996. Diuretic and natriuretic actions of melanin concentrating hormone in conscious sheep. *Journal of Neuroendocrinology* 8, 57–63.
- Pillon D, Caraty A, Fabian-Nys C and Bruneau G 2003. Early decrease of proopiomelanocortin but not neuropeptide Y mRNA expression in the mediobasal hypothalamus of the ewe, during the estradiol-induced preovulatory LH surge. *General and Comparative Endocrinology* 134, 264–272.
- Polkowska J and Gladysz A 2001. Effect of food manipulation on the neuropeptide Y neuronal system in the diencephalon of ewes. *Journal of Chemical Neuroanatomy* 21, 149–159.
- Polkowska J, Wańkowska M and Wójcik-Gładysz A 2006a. Expression of NPY-immunoreactive neurons in the hypothalamus of the cycling ewe. *Folia Histochemica et Cytobiologica* 44, 13–16.
- Polkowska J, Wójcik-Gładysz A and Wańkowska M 2006b. The effect of intracerebroventricular infusions of leptin on the immunoreactivity of neuropeptide Y and gonadotropin releasing hormone neurons in the hypothalamus of prepubertal sheep in conditions of short fasting. *Journal of Chemical Neuroanatomy* 31, 130–138.
- Qi Y, Iqbal J, Oldfield BJ and Clarke IJ 2008. Neural connectivity in the medialbasal hypothalamus of the sheep brain. *Neuroendocrinology* 87, 91–112.
- Qi Y, Henry BA, Oldfield BJ and Clarke IJ 2010. The action of leptin on appetite-regulating cells in the ovine hypothalamus: demonstration of direct action in the absence of the arcuate nucleus. *Endocrinology* 151, 2106–2116.

- Relling AE, Pate JL, Reynolds CK and Loerch SC 2010. Effect of feed restriction and supplemental dietary fat on gut peptide and hypothalamic neuropeptide mRNA concentrations in growing wethers. *Journal of Animal Science* 88, 737–748.
- Sartin JL, Wagner CG, Marks DL, Daniels J, McMahon CD, Obese FY and Partridge C 2005. Appetite regulation in sheep: a potential site for therapeutic intervention in disease models. *Domestic Animal Endocrinology* 29, 446–455.
- Sartin JL, Marks DL, McMahon CD, Daniel JA, Levasseur P, Wagner CG and Whitlock BK 2008. Central role of the melanocortin-4 receptors in appetite regulation after endotoxin. *Journal of Animal Science* 86, 2557–2567.
- Sartin JL, Dyer C, Matteri R, Buxton D, Buonomo F, Shores M, Baker J, Osbourne JA, Braden T and Steele B 2001. Effect of intracerebroventricular orexin-B on food intake in sheep. *Journal of Animal Science* 79, 1573–1577.
- Schwartz MW, Woods SC, Porte D, Seely RJ and Baskin DG 2000. Central nervous system control of food intake. *Nature* 404, 661–671.
- Soliman M, Abdelhady S, Fattou I, Ishioka K, Kitamura H, Kimura K and Saito M 2001. No alteration in serum leptin levels during acute endotoxemia in sheep. *Journal of Veterinary Medical Science* 63, 1143–1145.
- Sorenson A, Adam CL, Findlay PA, Marie M, Thomas L, Travers MT and Vernon RG 2002. Leptin secretion and hypothalamic neuropeptide and receptor gene expression in sheep. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 282, R1227–R1235.
- Thomas MG, Gazal OS, Williams GL, Stanko RL and Keisler DH 1999. Injection of neuropeptide Y into the third cerebroventricle differentially influences pituitary secretion of luteinizing hormone and growth hormone in ovariectomized cows. *Domestic Animal Endocrinology* 16, 159–169.
- Thomas SA, Preston JE, Wilson MR, Farrell CL and Segal MB 2001. Leptin transport at the blood – cerebrospinal fluid barrier using the perfused sheep choroid plexus model. *Brain Research* 895, 283–290.
- Tillet Y, Batailler M and Fellman D 1996. Distribution of melanin-concentrating hormone (MCH)-like immunoreactivity in neurons of the diencephalon of sheep. *Journal of Chemical Neuroanatomy* 12, 135–145.
- Valassi E, Scacchi M and Cavagnini F 2008. Neuroendocrine control of food intake. *Nutrition, Metabolism and Cardiovascular Diseases* 18, 158–168.
- Wagner CG, McMahon CD, Marks DL, Daniel JA, Steele B and Sartin JL 2004. A role for agouti-related protein in appetite regulation in a species with continuous nutrient delivery. *Neuroendocrinology* 80, 210–218.
- Whitlock BK, Daniel JA, McMahon CD, Buonomo VC, Wagner CG, Steele B and Sartin JL 2005. Intracerebroventricular melanin-concentrating hormone stimulates food intake in sheep. *Domestic Animal Endocrinology* 28, 224–232.
- Williams LM, Adam CL, Mercer JG, Moar KM, Slater D, Hunter L, Findlay PA and Hoggard N 1999. Leptin receptor and neuropeptide Y gene expression in the sheep. *Journal of Neuroendocrinology* 11, 165–169.
- Xu R, Roh SG, Gong C, Hernandez M, Ueta Y and Chen C 2003. Orexin-B augments voltage-gated L-type Ca<sup>2+</sup> current via protein kinase C-mediated signaling pathway in ovine somatotropes. *Neuroendocrinology* 77, 141–152.
- Xu R, Wang Q, Yan M, Hernandez M, Gong C, Boon WC, Murata Y, Ueta Y and Chen C 2002. Orexin-A augments voltage-gated Ca<sup>2+</sup> currents and synergistically increases growth hormone (GH) secretion with GH-releasing hormone in primary cultured ovine somatotropes. *Endocrinology* 143, 4609–4619.
- Yildiz S, Blache D, Celebi F, Kaya I, Saatci M, Cenesiz M and Guven B 2003. Effects of short-term high carbohydrate or fat intakes on leptin, growth hormone and luteinizing hormone secretions in prepubertal fat-tailed Tuj lambs. *Reproduction in Domestic Animals* 38, 182–186.
- Zaralis K, Tolkamp BJ, Houdijk JG, Wylie AR and Kyriazakis I 2008. Changes in food intake and circulating leptin due to gastrointestinal parasitism in lambs of two breeds. *Journal of Animal Science* 86, 1891–1903.
- Zaralis K, Tolkamp BJ, Houdijk JG, Wylie AR and Kyriazakis I 2009. Consequences of protein supplementation for anorexia, expression of immunity and plasma leptin concentrations in parasitized ewes of two breeds. *British Journal of Nutrition* 101, 499–509.
- Zhang S, Blache D, Veroe PE, Adam CL, Blackberry MA, Findlay PA, Eidne KA and Martin GB 2005. Expression of orexin receptors in the brain and peripheral tissues of the male sheep. *Regulatory Peptides* 124, 81–87.