

#### University of Tennessee, Knoxville Trace: Tennessee Research and Creative **Exchange**

Field & Commercial Crops

**UT Extension Publications** 

1-1-2010

### SP721 Sunflower: An Alternative Crop for Tennessee Producers - Production Guidelines and Tennessee Hybrid Trials

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: http://trace.tennessee.edu/utk agexcrop



Part of the Agronomy and Crop Sciences Commons

#### Recommended Citation

"SP721 Sunflower: An Alternative Crop for Tennessee Producers - Production Guidelines and Tennessee Hybrid Trials," The University of Tennessee Agricultural Extension Service, R12-5110-046-011-10 SP721-1M-1/10 10-0131, http://trace.tennessee.edu/ utk\_agexcrop/56

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the UT Ag Research website. This Production is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Field & Commercial Crops by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.



## Sunflower: An Alternative Crop for Tennessee Producers

Production Guidelines and Tennessee Hybrid Trials



# **Sunflower: An Alternative Crop for Tennessee Producers**Production Guidelines and Tennessee Hybrid Trials

M. Angela McClure, Associate Professor Fred L. Allen, Professor Richard D. Johnson, Research Associate Larry G. Heatherly, Adjunct Professor Department of Plant Sciences

#### Introduction

Tennessee producers are interested in crops that can be grown to diversify and/or complement their current cropping systems that include corn, cotton, soybeans and wheat. There is a growing demand for birdseed, and sunflower is an important component of that feed, either as a sole ingredient or in a blend with other seeds such as millet, corn, sesame, sorghum, wheat and oats. Sunflower oil is used for human consumption and is also a suitable feedstock for biodiesel. An increasing demand for biodiesel will increase the demand for oilseed crops such as sunflower. Thus, there may be increased opportunities for some Tennessee producers to grow sunflower for these markets.

This publication provides general information about the growth, development and production of sunflower. Results from four years of sunflower hybrid trials conducted in Tennessee are included to provide producers with information about how these hybrids perform under Tennessee conditions.

#### History - Past and Recent

Sunflower is one of a few crops that originated in the U.S., with the southwestern U.S. likely its center of origin. Records show that wild sunflower was used as a food by Native Americans and was domesticated and spread by their movements (Seiler and Rieseberg, 1997). Archaeological evidence uncovered from a site in Middle Tennessee indicates that sunflower was being grown in Tennessee by Native Americans more than 4000 years ago (Crites, 1993). Following the discovery and settlement of the U.S., sunflower was spread to other parts of the world, with European countries and Russia being the major producers (Putt, 1997).

Modern sunflower varieties in North American trace much of their lineage back to reintroduced varieties that were developed in Europe and Russia. Sunflower was not an important agronomic crop in the U.S. until the 1950s, and oilseed sunflower has been an economically important crop in the U.S. only since the mid-1960s. Expanded world production of sunflower resulted from development of high-oil varieties and more recently from the development of hybrids.

Table 1. Acres of sunflower	harvested in the U.S. in 20	008.	
State	Seed for oil	Seed for non-oil use	All types
Colorado	155,000	23,000	178,000
Kansas	205,000	20,000	225,000
Minnesota	77,000	38,000	115,000
Nebraska	42,000	18,000	60,000
North Dakota	915,000	145,000	1,060,000
South Dakota	522,000	48,000	570,000
Texas	59,000	33,000	92,000
All other states	74,000	11,000	85,000
United States total	2,049,000	336,000	2,385,000
Source: USDA-NASS, 200	8		

Presently, most U.S. commercial sunflower production is in the Great Plains states of Colorado, Kansas, Minnesota, Nebraska, North Dakota, South Dakota and Texas (USDA-NASS, 2007). Most acreage is grown to produce seeds for vegetable oil (**Table 1**) with a small portion used for birdseed production. In 2009, average U.S. yield of sunflower seed was 1538 pounds per acre (USDA-NASS, 2009). Even though sunflower is adapted to Tennessee conditions, its primary production is in the western and upper Great Plains.

#### Types and Uses

There are two types of sunflower hybrids: 1) the oilseed type that is grown for vegetable oil, and 2) the confection or non-oilseed type (**Fig. 1**). The oilseed type has a higher oil composition in the seeds than the non-oilseed type. Oilseed types produce smaller black seeds and the oil is primarily used for human consumption. The oilseed types are also marketed as a sole ingredient for birdseed or in birdseed blends. The non-oilseed type produces the large, striped seeds that are used for human food snacks in the shell or as kernels, in baking ingredients, and in birdseed mixes. Because each type has a separate and distinct market, they cannot be mixed in storage (Johnson et al., 2009).

Commercial oilseed sunflower hybrids are divided into three categories based on the fatty acid profiles (types of saturated and unsaturated fats) of the oil in the seeds. The categories are 1) standard or linoleic, 2) NuSun or midoleic, and 3) high oleic. A comparison of the fatty acid profiles of seeds of these sunflower types and other oilseed crops is shown in **Table 2**.

Vegetable oil from sunflower seeds is lower in saturated fats than most vegetable oils. Linoleic oil processed from sunflower oil is used as a low-saturated fat cooking oil. Linoleic types were the predominant oil-sunflower hybrid produced, but their acreage has decreased. NuSun is currently the predominant oil-type sunflower grown, because seeds produce a healthier oil that contains less saturated fat than oil from linoleic types. Oil from seeds of NuSun types does not have to be hydrogenated, which makes it an excellent frying oil with a long shelf life. High-oleic hybrids produce seeds that contain specialty oil that is very low in saturated fats. This specialty oil is used in lubricants (both food grade and industrial) and food coatings. It is grown by contract only (Johnson et al., 2009).

Oil from sunflower seeds comprises 7.8 percent of the world's vegetable oil consumption, which is the fourth leading oil consumed behind palm (31.8 percent), soybean (30 percent) and rapeseed (14 percent) (ASA, 2008). Its seed typically contain about 38 to 44 percent oil and 18 to 25 percent protein. In contrast, soybean seeds typically contain about 20 percent oil and about 40 percent protein, or a mirror image of sunflower oil and protein content. Non-dehulled or partly dehulled sunflower meal has been substituted for soybean meal in diets for ruminant animals, swine, poultry and catfish.



Fig. 1. The two classes of sunflower based on seed characteristics: (1) oilseed type grown for oil and meal, and (2) non-oilseed or confection type grown for human and bird food. [From Berglund (2007a, Fig. 1); original photo credited to Gerhardt Fick; used with permission]

Oil source	Monounsaturated	Polyunsaturated	Saturated
		%	
High-oleic sunflower	82	9	9
Olive	72	11	17
NuSun sunflower	65	26	9
Rapeseed (Canola)	62	32	6
Peanut	49	33	18
Lard	47	12	41
Beef fat	44	4	52
Palm	39	10	51
Butter fat	34	2	64
Corn	25	62	13
Soybean	24	61	15
Linoleic sunflower	20	69	11
Cottonseed	18	55	27
Safflower	13	77	10

#### Growth and Development

Sunflower is an annual, erect, broadleaf plant with a strong taproot and prolific lateral root system. It emerges from the soil with two large cotyledons. Emergence will take four to five days when planted an inch deep in warm

soil, but will take a few days longer in cooler soils or when planted deeper. Soil crusting can make it difficult for the large-cotyledon seedlings to push out of the soil. A rotary hoeing may be necessary to mitigate the soil crusting problem for easier emergence.

Sunflower grows rapidly, producing large, rough leaves. Current sunflower varieties in Tennessee reach an average of 6 feet in height, varying between 5 and 7 feet depending on planting date, variety and soil conditions. After reaching full height at blooming, heads on commercial cultivars turn downwards, a trait that inhibits bird feeding on the seeds.

Each sunflower head, or inflorescence, is not a single flower, but rather 1,000 to 2,000 individual flowers joined at a common receptacle (**Fig. 2**). The head is actu-




Fig. 2. Details of a sunflower head with selected parts labeled. [From Berglund (2007a, Fig. 3); original photo credited to J. Miller and Christian Y. Oseto; used with permission]


ally composed of two types of flowers. What appear to be yellow petals around the edge of the head are actually individual ray flowers. The face of the head is comprised of hundreds of disk flowers, which each form into a seed (achene). Commercial sunflower has flowers that are self-compatible for pollination, meaning they do not require a pollinating insect. However, some studies have shown that bee pollination provides a slight yield boost. Sunflower heads turn with, or track, the sun early in their development, but later stay east-facing before facing downwards. A common practice is to plant rows north and south so that the heads can lean into the between-row space rather than bumping against an adjacent in-row plant and causing some seeds to fall. Heads on commercial varieties turn downward after blooming, which makes it more difficult for birds to eat the seeds.

Table 3 contains a description of sunflower growth stages. Pictorial views of the growth stages defined in **Table 3** are shown in **Fig. 3**. Determining stage of development is based on using the main branch or head and not branch heads. Generally, sunflower reaches R1 or bloom stage about 65 to 70 days after planting and maturity about 105 to 115 days after planting (Aiken, 2005). Hybrid differences in maturity are usually associated with differing lengths of the vegetative period before the head is visible (Putnam et al., 1990).


Table 3. Description	of sunflower growth stages. See Fig. 3 for color photos of various growth
stages.	
Stage†	Description
VE	Emergence
V1 to n— Vegetative stages	Determined by counting the number of true leaves at least 1.5 inches in length beginning as V-1, etc. If lower leaves have dropped, count leaf scars.
V20	20 true leaves
R1—beginning of Reproductive stages	The terminal bud forms a miniature floral head rather than a cluster of leaves. When viewed from above, the immature bracts have a many-pointed, star-like appearance.
R2	Immature terminal bud < 1 inch above nearest leaf attached to the stem. Disregard leaves attached to the back of the bud.
R3	Immature bud > 1 inch above the nearest leaf.
R4	The inflorescence or bud begins to open. When viewed from above, immature ray flowers (on outer edge of head) are visible.
R5	Beginning of flowering. Can be divided into sub-stages dependent on the percentage of the head area (interior disk flowers) that has completed or is in flowering; e.g., R 5.3 = 30 percent of head area completed flowering, R 5.8 = 80 percent, etc.
R5.5	50 percent flowered
R6	Flowering is complete and ray flowers (on outer edge of head) are wilting.
R7	Back of head has started turning pale yellow.
R8	Back of head is yellow but bracts (behind ray flowers) remain green.
R9	Bracts become yellow and brown. Physiological maturity.
Berglund (2007a) (A	lso color photos of various growth stages.)

#### **Vegetative Stages**

Fig. 3. Stages of sunflower development. See Table 3 for description of stages. [From Berglund (2007a, Fig. 4); original credited to A. A. Schneiter and J. F. Miller; used with permission]



#### **Reproductive Stages**



#### Cultural Practices

Guidelines for growing sunflower in the U.S. are available. The following information is a composite of the material from sources shown in **Table 4** plus sources cited at specific locations throughout the report.

Table 4. Information sources for U.S. sunflower produc	tion.
Publication	Institution and Web site
ProCrop Sunflower Menu (ProCrop, 2008)	North Dakota State Univ. Ext. Serv.
Sunflower – Alternative Field Crops Manual (Putnam et al., 1990)	Univ. of Wisconsin and Univ. of Minnesota Ext. Serv.
Sunflower Production (Berglund, 2007b)	North Dakota State Univ. Ext. Serv.
High Plains Sunflower Production Handbook (Meyer et al., 2009)	Kansas State Univ. Coop. Ext. Serv.
Sunflower (Myers, 2008)	Thomas Jefferson Agricultural Institute

#### **Hybrid Selection**

Almost all commercial varieties of sunflower are hybrids, so new seeds should be purchased each year. Hybrids should be selected on the basis of high yield with high seed oil content (at least 40 percent), a test weight of at least 25 pounds per bushel, and disease and insect resistance if available. Hybrids with resistance to rust, Verticillium wilt and certain races of downy mildew are available. Given a choice, select a high-oil hybrid instead of a low-oil hybrid with the same yield potential. The oilseed market pays a premium for seeds with more than 40 percent oil (at 10 percent moisture), and discounts seeds with less than 40 percent oil.

The University of Tennessee conducts periodic sunflower variety trials at the Research and Education Centers to identify varieties with desired traits that are less prone to lodging and more productive under Tennessee growing conditions. Hybrids with traditional (linoleic), mid-oleic (NuSun) and high-oleic oil composition are tested. Newer Clearfield® hybrids are tolerant to Beyond® herbicide. Lodging and yield data from the Tennessee tests are available at http://varietytrials.tennessee.edu.

#### Planting

Sunflower should be planted 1 to 2 inches deep. The shallower planting depth is preferred in cool, wet soils or when planting small seeds. Sunflower can be planted anytime after soil has warmed to about 44 degrees F; however, it is preferable to delay planting until soil has warmed to at least 50 degrees F. This occurs in early April at Jackson (USDC-NCDC, 2008). Planting early generally will result in higher yield, test weight and oil content of the seeds in the northern U.S. sunflower-growing regions (Meyer et al., 2009; ProCrop, 2008). It remains to be seen how early planting will affect sunflower yield and seed characteristics in the southern U.S. Planting date can be used in the management of some insects, but not all affected species respond similarly to early or late planting. Sunflower can be doublecropped with wheat, but yields will be considerably lower.

#### **Row Spacing and Plant Population**

A 30-inch row spacing is the most popular and considered standard, although narrower rows and solid seeding can be used. Wide rows offer more options for weed management and allow harvesting with a row-crop header. Sunflower will compensate to some extent for differences in plant population through adjustments in head size, number of seeds per plant and seed size. Plant populations for oilseed hybrids should be between 14,000 and 22,000 final plants per acre. The higher populations may be helpful for weed management. Sunflower is not particularly sensitive to seeding rate, since head size (and seed number) per plant will increase in a thinner stand. A lower population of 14,000 final plants/acre is suitable for non-oilseed types in order to ensure large seeds.

Sunflower seeds for planting are sold either by weight or seeds per bag; however, sunflower seeding should be based on number of seeds per acre and not weight. Oil-type hybrid seed sizes are #2 (largest), #3 and #5 (smallest). Size #3 is most commonly planted. Size of the seeds can affect maximum depth of planting and what type of

planter modifications are necessary, such as seed plates or finger pickups. The size of planted seeds has no apparent effect on agronomic performance.

Dr. Emerson Nafziger (Nafziger, 2008) of the University of Illinois Department of Crop Sciences has published an online calculator for determining soybean seeding rate, seeds per foot of row and associated costs for an intended plant population in a chosen row spacing. This calculator also can be used for sunflower seeding rate calculations. An example calculation using this calculator is as follows:

A row spacing of 30 inches with a desired final population of 20,000 plants per acre is entered. This example assumes that 80 percent of planted seeds will become viable plants. Thus, 20,000/0.80 = 24,700 seeds per acre to plant. In 30-inch-wide rows, this equates to 14.3 seeds per 10 feet of row, or

Table 5. Seed spacing (distance between seeds in a row rounded to nearest 0.5 inch) required for desired final populations in indicated row spacings assuming 80 percent final stand (90 percent germination and 10 percent stand loss).

Final stand		Row s <sub>1</sub>	pacing (i	nches)	
(plants per acre)	(7.5)	(15)	(20)	(30)	(40)
14,000	48.0	24.0	18.0	12.0	9.0
16,000	42.0	21.0	15.5	10.5	8.0
18,000	37.0	18.5	14.0	9.5	7.0
20,000	33.5	16.5	12.5	8.5	6.5
22,000	30.5	15.0	11.5	7.5	6.0
24,000	28.0	14.0	10.5	7.0	5.5
26,000	26.0	13.0	9.5	6.5	5.0
Source: Nafziger (200	08).				

8.5 inches between seeds within a row. **Table 5** contains other values (obtained from the online calculator) that can be used to make the proper planter setting to achieve a desired final stand.

#### Soil Fertility and Fertilizer Recommendations

A summary of typical nutrient content of sunflower stover and seeds is shown in **Table 6**. Using these values as a guide, a sunflower seed yield of 1,500 pounds per acre removes about 75, 23 and 55 pounds of N, P and K from an acre of soil. Thus, these amounts must be present to ensure this yield level according to these values. A soil test should be used to determine nutrient levels in the soil before growing sunflower. Only those nutrients determined to be below levels for optimum production should be re-supplied as fertilizer.

Research results indicate that sunflower requires 6 to 7 pounds of N for every 100 pounds of seed production. Using these values, a yield of 1,500 pounds of seed per acre

Table 6. Nutrient content (pounds per acre) in a
sunflower crop producing 1,000 pounds of seed
per acre.

Per derei			
Element	Seed	Stover	Total
Nitrogen (N)	30	18	48
Phosphorus (P)	12	3	15
Potassium (K)	8	28	36
Sulfur (S)	2	4	6
Magnesium (Mg)	2	5	7
Calcium (Ca)	1.2	18.5	19.7
Source: Vigil, Hergert, a	nd Meno	rel (2009	)

requires 90 to 105 pounds of N per acre. Fertilizer N rates should be lowered if sunflower is planted after wheat or if legumes are grown in rotation before sunflower. In Tennessee, this legume would likely be soybean, which provides an N credit of up to 20 pounds per acre (Savoy and Joines, 2009). It is important to apply only the amount of N needed to reach the desired yield goal, because excessive N can result in decreased oil content and increased lodging, as well as N loss to the environment. When properly applied, N source materials are not agronomically different; thus, N source should be based on cost and N content of the various N fertilizers (USDA-ERS, 2008) and availability. Nitrogen application can be made preplant, sidedress or a combination of the two. Applications should be timed to ensure N is available for rapid growth and development.

Phosphorus (P) and potassium (K) fertilization should be done according to soil test recommendations. If soils test medium or higher, response to P fertilization likely will be small to none. Phosphorus fertilization is recommended for soils that test below medium. Periodic soil tests will determine if adequate P remains in the soil to produce a crop in subsequent years. Phosphorus should be applied preplant-broadcast, preplant-knifed or banded at planting. Potassium deficiencies are not likely unless soil test levels are low. Potassium should be applied preplant-

broadcast and incorporated. Selection of P and K fertilizer materials should be based on cost (USDA-ERS, 2008) and availability. Liming is recommended for sunflower on soils with a pH of 6.0 or less.

#### Weed Management

Using best management practices for sunflower production will reduce the negative effects of weeds. However, sunflower does not provide a quick ground cover, so early-season weed control is essential. Thus, good production practices must be supplemented with chemical and cultural weed control measures, particularly in the southern states. Conventional tillage sunflower can be row-cultivated as late as the 4- to 6-leaf stage, as long as the soil is not disturbed any closer to the row than the plant leaf spread. This will minimize root pruning during cultivation.

Only a limited number of herbicides are labeled for use in sunflowers, but good weed control can be obtained with proper application of what is available. Included in these herbicides are those that can be applied preplant, preemergent or post-emergent. Annual grass weeds and small-seeded broadleaf weeds can be controlled with soil-applied pendimethalin, trifluralin, ethafluralin and S-metolachlor. The tillage-incorporation requirement for some of these herbicides will interfere with no-till production. Sulfentrazone can be applied preplant if sunflower is planted in rows where seed depth can be assured to prevent injury. After sunflower emergence, emerged grasses can be controlled with clethodim or sethoxydim. Where Clearfield® sunflower hybrids are used, emerged broadleaf weeds can be controlled with imazamox. Where non-Clearfield® hybrids are planted, row-cultivation is the only option for controlling broadleaf weeds once the crop has emerged.

Sunflower is sensitive the carryover of sulfonylurea (Steadfast\*, Resolve\*, Classic\*), imidazolinone (Pursuit\*) and sulfonamide (FirstRate\*) classes of herbicides. These products require a rotation interval of up to 18 months between herbicide use and planting back to sunflower. Check all herbicide labels for crops grown in the year preceding sunflower for precautions and rotation restrictions.

Two southern information sources provide up-to-date sunflower weed control recommendations. They are University of Tennessee Extension Publication PB 1580 (Steckel, 2008) entitled "Weed Control Manual for Tennessee" and Mississippi State University Publication 2434 (Rankin, 2007) entitled "Sunflower Weed Control Recommendations for Mississippi."

#### Insect Management

Insects are occasionally a problem in sunflower production. Sunflower seeds can be treated commercially with an insecticide such as imidacloprid (Gaucho\*) or thiamethoxam (Cruiser\*) for protection from soil insect pests. Some insects such as the sunflower midge, sunflower beetle, sunflower stem weevil, red sunflower seed weevil, headclipping weevil and the banded sunflower moth, need to be monitored for infestation levels that justify application of control measures. Growers should minimize insect damage by applying integrated pest management practices that combine biological, cultural and chemical control measures to minimize economic, health and environmental risks. Application of control measures should be based on economic threshold levels, when available, to maintain pest populations below levels that cause unacceptable crop quality and yield losses.

Treatment thresholds should be used as a guide based on yield potential and crop value. For example, lower thresholds might be warranted where higher crop values or yields are expected. This is the case for non-oilseed types because of the requirement for seeds that are free of insect damage. In fact, insect-damaged non-oilseed types will be discounted or rejected by processors, whereas premiums for large, insect-free seeds are common. Conversely, when the crop is expected to be lower in yield or value, higher thresholds might be used.

Many insects feed on sunflower foliage but the defoliation level rarely is high enough to cause significant yield loss during vegetative development (**Table 7**). Defoliating insects cause the most damage from about R1 to R3. Insects that infest flowers and seeds are the most damaging to yield and quality potential; therefore, thorough scouting should be conducted during the reproductive period to monitor levels of these insects.

Insect management in sunflower is covered in detail by Knodel and Charlet (2007) and Sloderbeck et al. (2009). Both of these sources have textual and pictorial descriptions to aid in the identification and symptoms of damage of the various insect species, and provide information on scouting methods, economic thresholds if available, and management and/or control options for common insect species. Insecticides labeled for sunflower are available,

and their use is described by Sloderbeck, Michaud, and Whitworth (2008). Not all insecticide products that are labeled for use in the Great Plains area can be used in Tennessee. Check all insecticide labels prior to their use.

#### Disease Management

The most serious diseases of sunflower are caused by fungi. Major diseases include sclerotinia stalk and head rot, verticillium wilt, rust, phoma black stem and downy mildew (Putnam at al., 1990). Sclerotinia (white mold) is also found in soybean, canola and certain other broadleaf plants. In cool wet soils, seeds or seedlings may be attacked by fungi, so seeds are typically treated with fungicide [e.g., Maxim XL (McMullen and Bradley, 2007)]. Hybrids with tolerance to races of rust, some races of downy mildew, verticillium wilt, and other disease pathogens are

Table 7. Approximate percentage yield reduction when sunflower is defoliated the indicated amount at each indicated growth stage (See Table 3 for growth stages).

		Percen	tage defo	liation	
Plant stage	10	30	50	70	100
	(	Approxi	mate % y	vield loss	;)
V-4 to V-5	0	2	4	5	21
V-9 to V-11	0	3	5	7	24
R1	2	6	7	16	47
R3	2	15	24	44	99
R5	1	7	16	37	90
R7	0	3	10	16	22
R8	0	2	5	8	11
Source: Sloderh	secla et al	(2009)			

Source: Sloderbeck et al. (2009)

available. The most economical and effective management of sunflower diseases is the planting of resistant or tolerant hybrids, and using a rotation scheme that allows a minimum of three to four years between successive sunflower crops. Excellent and detailed textual and pictorial descriptions of disease symptoms can be found in Bradley et al. (2007) and Jardine (2009). Specific management recommendations for the various sunflower disease pathogens can be found in ProCrop (2008).

#### **Bird Control**

Birds can be major pests in sunflower. Ripening seeds are exposed and the large head serves as a ready perch during feeding. Problem birds feed on insects and weed seeds in sunflower fields before the crop is vulnerable to damage, and thus become accustomed to feeding in that location. Cultural practices in combination with mechanical and chemical practices can and should be used to control birds in sunflower fields.

Bird damage to sunflower is imminent. Thus, prevention of bird infestations that will damage sunflower is the first line of defense. Sunflower should not be planted near cattail marshes or woodlots. All planting in a region should be done at or near the same time because earlier and later-maturing fields will sustain more damage. Weeds and insects in the crop should be controlled early because insects and weed seeds are often a source of food for the birds before the crop becomes susceptible to bird feeding. Planting a small plot of oilseed sunflower near high-value non-oilseed and oilseed crops can provide a trap crop to protect the higher-value crop. Providing alternative food plots will likely enhance the efficacy of any repellent or scare device and should be included in bird damage management plans. Delayed plowing down of harvest stubble until after harvest is completed should provide an alternate feeding area. Harvesting sunflower as early as possible will avoid prolonged exposure of the seeds to bird damage.

Physical disruptions to bird feeding on sunflower seeds include guns, automatic exploders and electronic frightening and pyrotechnic devices. However, none are 100 percent effective. Repellent baits are available, but their efficacy is inconsistent. The effectiveness of bird-repellent materials applied to the head is limited because the orientation of the head prevents contact of the applied material with the developing seeds. This is an important factor to consider before using any chemical bird repellent since sunflower heads are downward-oriented when bird damage potential is greatest. A detailed outline of measures to control bird damage to sunflower is provided by Linz et al. (2006) and Latzke (2007).

The U.S. Department of Agriculture's Animal and Plant Health Inspection Service (USDA-APHIS) has a Wildlife Damage Management Division. Within this agency, there is a Wildlife Services Division that conducts research to evaluate methods for managing and/or controlling bird damage to crops. Online information for the

Wildlife Services Division in Tennessee can be accessed by going to the above Web site, searching "Tennessee," and then accessing the "Wildlife Services – Tennessee" listing.

#### Crop Rotation and Residue Management

Sunflowers can be rotated with corn, soybean and/or sorghum. Yields of sunflower doublecropped with wheat may be too low to be profitable. Regardless of cropping system, sunflower should not be planted in the same field more than once every three to four years. Sunflower, like soybean, does not leave very much residue, so a fall cover crop should be considered on erosive fields. Sunflower grows best on well-drained soils, which means they are probably not suited for the alluvial clay soils along the Mississippi River.

#### Harvesting and Storage

Sunflower seeds are generally physiologically mature when the back of the flower head is yellow. When the head turns brown on the back, seeds are usually ready for harvest. Platform, row-crop and corn heads can be used to successfully harvest sunflower. Row-crop heads can be used without modification. Corn heads need to be modified with a stationary cutting knife. Platform heads can be used without modification, but often have a higher amount of seed and head loss than a row head. Adding pans to the front of the platform and/or modifying the reel can improve efficiency (Myers 2008). Because getting sunflower heads into the combine is probably the biggest problem when harvesting, header performance can often dictate combine efficiency

Combine settings must be adjusted for sunflower versus other crops. A good description of these settings is provided by Myers (2008). The overall goal of the threshing process should be passing the head nearly intact through the combine, or in a few large pieces, with all developed seeds removed from the head. If the head is being ground up into small pieces, the grain will have an excessive trash content (Myers 2008).

During colder periods, sunflower can be safely stored at 10 percent moisture or less, but during warmer months the storage moisture should be at 8 percent or less. Proper moisture sampling procedures during storage, storage bin requirements, and air temperatures for drying are provided by (Myers 2008). Be aware that sunflower dries more rapidly than corn or soybeans, and should be monitored to avoid overdrying. Also, be aware that sunflower drying has a higher risk of fire hazard than some crops, because small fibers that rub off the sunflower hulls and float in the air can readily burn. Precautions to avert this hazard are provided by Myers (2008).

#### Marketing and Economics

The easiest way to market sunflower is often to the birdseed market. However, many current or potential sunflower producers are interested in non-oilseed confection sunflower because of its higher price. As stated previously, this market has a more demanding standard for a high-quality, undamaged seeds suitable for human food. Most sunflower producers grow the oilseed type.

With the emergence of NuSun and high oleic types, opportunities for a price premium of 10 to 20 percent exist. Until sufficient markets develop for these specialty oil types, growers may be required to ship them long distances and this should be considered before growing for this market. No processing plants are located in Tennessee at this time. The nearest processors accepting sunflower for oil are in Georgia (AG Strong, Dalton, GA), Alabama (AG Strong, Athens, AL), and Missouri. It is critical that growers make local arrangements for storage and shipping and have a contract with a processor prior to planting a crop of sunflower where the intended market is crushing for oil and meal.

Representative budgets using October 2008 input costs and commodity prices for oil type and non-oilseed type sunflower are provided by Dumler et al. (2008) of Kansas State University. According to their budgets (with modifications for Tennessee conditions and lower fertilizer prices), breakeven yields needed to cover all costs except management and overhead for oil type and non-oilseed type sunflower are about 1800 and 1200 pounds per acre, respectively. These yields are based on receiving \$17.06 and \$26.50 per hundredweight for the two types, respectively. These yield levels will be used to assess yields obtained from the following Tennessee yield trials. Of course, alternative prices paid (especially for fertilizers) and received, as well as inputs and their amounts used (fertilizers, herbicides, and pesticides) by individual Tennessee producers will result in deviations from these estimates.

#### 2004 - 2007 Performance of Sunflower Hybrids in Tennessee

#### Acknowledgements

The authors are grateful to Rob L. Myers, Director of Programs, and other personnel at the Jefferson Institute (www.jeffersoninstitute.org) for providing most of the seed each year for the yield tests, and for providing publications and references that were helpful in developing this publication.

#### **Experimental Procedures**

Sunflower hybrid trials were conducted at the East Tennessee (Knoxville) and Milan Research and Education Centers (REC). The trials contained 15 (2004), 17 (2005), 25 (2006) and 31 (2007) hybrids at each location. Nitrogen at 90 pounds per acre was applied in split applications to each test site each year. A recent Mississippi study confirms that this amount of N is sufficient for maximum sunflower yield in the southern U.S. (Zheljazkov et al., 2008). Plots were two rows wide (30-inch row width) and 30 feet long. Plots at each location were replicated three times in a randomized complete block design.

Planting dates, seeding rates, harvest dates and soil series for each site are shown in **Table 8**. All monocropped tests were planted in May and doublecropped tests at Milan in 2006 and 2007 were planted in June.

#### Interpretation of Data

Tables list entries in descending order of performance at each location. All yields were adjusted to 10 percent moisture. LSD (least significant difference) values for each test are shown at the bottom of each respective table. Average yields of any two varieties being compared must differ by at least the LSD to be considered different in yielding ability at the 5 percent level of probability. For example, if the LSD for a test is 450 lbs/a and the average yield of Hybrid A is 1700 pounds per acre and the average yield of Hybrid B is 1300 pounds per acre, then the two hybrids are not statistically different in yield because the difference of 400 pounds per acre is less than the minimum of 450 pounds per acre required for them to be significantly different. Similarly, if the average yield of Hybrid C is 2200 pounds per acre then its yield is significantly greater than yields of both Hybrids A and B because the difference between B and C (900 pounds) and between A and C (500 pounds) exceeds the LSD value of 450 pounds.

The coefficient of variation (C.V.) value shown in each table is a measure of the variability associated with each experiment. A C.V. of 10 percent indicates that the error variation of the experiment is about 10 percent of the size of the experiment's average. Similarly, a C.V. of 30 percent indicates that error variation is nearly one-third as large as the experiment's average. The goal in conducting each yield test is to keep the C.V. as low as possible, preferably below 20 percent.

#### **Growing Season Weather**

**2007.** The growing season was characterized by extremes. A late frost and very low temperatures in the first portion of April caused wheat and corn crop damage. The remainder of the season was characterized by record-setting heat and drought that lowered yields. Daytime temperatures were high (several days above 100 deg. F) during flowering and seed fill at many locations throughout the state.

**2006.** The growing season was characterized by hot, dry conditions through most of the growing period. Daytime temperatures across Tennessee were high (several days above 90 deg. F) during flowering and seed fill.

**2005.** The growing season was characterized by several timely rainfall events during critical periods. Rainfall events were prompted by hurricane aftermaths (especially Dennis, Katrina and Rita) passing through the state. Daytime temperatures across Tennessee were high (several days above 90 deg. F) during flowering and seed fill.

**2004.** The growing season was characterized by very favorable temperatures and rainfall for seed production. Adequate amounts and very timely distribution of rain, as well as lower than normal day and night temperatures, resulted in an exceptionally good growing season.

#### Results

**2007.** The vast majority of the hybrids had test weights that exceeded the industry standard of 25 pounds per bushel (**Table 9**). Yields from the Knoxville site and from the doublecropped trial at the Milan site were extremely low. About half of the hybrids in the May planting at Milan produced yields that exceeded 1800 pounds per acre. Yield of the non-oilseed type 'Triumph 777 C' hybrid exceeded 1200 pounds per acre only in the May-planted Milan trial.

**2006.** Average yields of all hybrids grown at the Knoxville and Milan sites were below 1800 pounds per acre (**Table 10**). The vast majority of the hybrids had test weights that exceeded the industry standard of 25 pounds per bushel. A majority of the hybrids suffered minimal bird damage. Average yields of hybrids grown at the Knoxville site were greater than those at the Milan site. Yield from several of the hybrids grown at Knoxville exceeded 1800 pounds per acre, whereas none did at Milan. Yields from hybrids planted in June following wheat at Milan were generally greater than yields from May-planted hybrids, and two of the later-planted hybrids produced yields that exceeded 1800 pounds per acre. Yield from the non-oilseed type hybrid 'Triumph 777 C' was near or exceeded the breakeven yield in all tests.

**2005.** Only the 1750 pounds per acre average yield of 'Triumph 636' approached 1800 pounds per acre (**Table 11**). A majority of the hybrids suffered significant bird damage, and several hybrids had a test weight below the 25 pounds-per-bushel industry standard. Yields at Knoxville were generally greater than those at Milan. Yield of several of the hybrids grown at the Knoxville site exceeded 1800 pounds per acre, whereas none did at Milan. Yield of the non-oilseed type hybrid 'Triumph 777 C' exceeded the arbitrary breakeven yield of 1200 pounds per acre at Knoxville but not at Milan.

**2004.** Average yields from the Knoxville and Milan sites were generally lower than the arbitrary breakeven yield of 1800 pounds per acre (**Table 12**). Only the 'Triumph 636' average yield of 1953 pounds per acre exceeded 1800 pounds per acre. Low yields were generally associated with significant bird damage. Test weight of all hybrids exceeded the industry standard of 25 pounds per bushel. Several hybrids at the Knoxville site exceeded 1800 pounds per acre, whereas none of the hybrids grown at Milan exceeded this yield.

Combined. Average yields of all hybrids that were grown at both locations during the 2005-2007 period were below 1800 pounds per acre (**Tables 13-15**). This resulted from the erratic yield performance of all hybrids across years at both locations. Average test weights of most hybrids across the three-year period exceeded the industry standard. Over the four-year period (2004-2007), the same trend occurred in both average yields and test weights. However, the average yield of 'Triumph 636' did approach 1800 pounds per acre at Knoxville, and this hybrid had the highest average yield at Milan. Across the 2006-2007 period, average yields from the doublecropped trials at Milan were low and well below the average yields from the May-planted trials at both locations.

#### **Summary and Conclusions**

Several factors should be considered when assessing the results from these yield trials. First, the significant bird damage in some years indicates that the bird problem will have to be mitigated if sunflower is to provide consistently profitable yields in Tennessee. Second, as discussed previously, sunflower can be planted in early April in Tennessee. It is unknown how this might affect yield potential. Third, using the above-cited economic parameters and these four years of variety trial results, it is obvious that yields from sunflower grown in Tennessee will have to consistently equal or exceed those from the best hybrids in these trials. Fourth, these results indicate that yields of sunflower grown in Tennessee will be erratic across years. The generally low average yields in these trials (for whatever reason) compared to the U.S. average yield (1454 and 1334 pounds per acre for oil type and non-oilseed type, respectively, in 2007; USDA-NASS, 2008) indicate that Tennessee producers should consider producing for a premium market to realize a profit. Fifth, these limited data indicate that sunflower should not be considered as the summer crop in a doublecropping system.

There are items that should be addressed if sunflower production is to gain a foothold among Tennessee producers. First, the best planting date for maximum yields should be identified. Second, there is no knowledge about

the pressure from insect and disease infestations that may occur with increased sunflower plantings across a wide area. Third, access to markets must be ascertained before sunflower is promoted for wide-scale planting in the southern U.S. Fourth, limited herbicides for use in sunflower that has no transgenic weed control component may be problematic. Finally, rotational crops should be identified for use with sunflower since its monoculture is not recommended.

		essee Research and Ed	ucation Centers (REC)	) where sunflower hybrid
tests were conduc	ted in 2004-2007.			
REC	Planting Date	Harvest Date	Seeding Rate	Soil Type
			seed/acre	
2007		•	•	
Knoxville	May 8	September 6	25,000	Stasser Silt Loam
Milan	May 8	August 29	25,000	Grenada, Henry Silt Loam
Milan (DC†)	June 15	October 10	25,000	Grenada, Henry Silt Loam
2006		· · · · · · · · · · · · · · · · · · ·	·	
Knoxville	May 9	August 21	19,000	Sequatchie Silt Loam
Milan	May 19	August 28	25,000	Grenada, Henry Silt Loam
Milan (DC)	June 7	October 3	25,000	Grenada, Henry Silt Loam
2005	•		•	
Knoxville	May 12	August 26	25,000	Sequatchie Silt Loam
Milan	May 20	September 6	25,000	Falaya Silt Loam
2004		<del></del>	•	· ·
Knoxville	May 21	August 31	25,000	Sequatchie Silt Loam
Milan	May 20	September 13	25,000	Loring, Henry Silt Loam
†DC = doublecro	pped – planted followir	ng wheat.		

Table 9. Mean yields and agronomic characteristics of 31 sunflower hybrids evaluated in three environments in Tennessee during 2007.

		Ava. Yield+			Double	Moisture	Test		Head	Plant	)	Bird
		± Std. Err.			Crop	at Harvest	Weight	Ö	Diameter	Height	Lodging	Damage
Brand	Hybrid	(n=3)	Knoxville §	Milan	Milan	(n=1)	(n=3)	(n=3)	(n=1)	(n=3)	(n=3)	(n=1)
			lbs/a			% -	nq/sqI	%	in.	'n.	score	%
<b>Traditional Hybrids</b>	Hybrids											
Dekalb	DKF 39-01	$712 \pm 79$	285	1642	210	7.5	30.0	42.3	4.2	45	<u>+</u>	2.8
Pioneer	63A70	$666 \pm 100$	326	1363	275	7.3	27.1	44.2	5.5	45	<u>+</u>	2.7
Mycogen	8N270	$574 \pm 73$	512	696	242	7.6	30.4	4 1.1	4.5	38	6.1	2.5
Mid Oleic	Mid Oleic Hybrids (NuSun)											
Croplan	356 NS	$962 \pm 73$	729	1722	433	7.8	30.3	42.3	5.0	43	1.2	<del>6</del> .
Triumph	636	$944 \pm 95$	345	1998	489	7.4	28.1	43.2	4.8	20	2.1	2.5
Pioneer	63M80	$934 \pm 72$	298	2037	468	7.7	30.0	42.4	5.3	47	1.3	2.8
Mycogen	8N453	$930 \pm 72$	699	1871	251	7.7	27.3	44.6	4.7	48	<del>1</del> .3	2.8
Triumph	645	$911 \pm 72$	401	1798	533	7.6	30.6	44.6	4.2	49	1.9	2.0
Dekalb	DKF 37-31	$900 \pm 78$	519	1774	406	7.9	30.1	42.2	5.0	46	1.5	2.0
Triumph	12 099	$885 \pm 79$	588	1711	356	7.8	25.5	41.7	4.3	47	1.3	3.2
Dekalb	DKF 35-10	$866 \pm 73$	642	1632	323	7.8	29.1	40.1	5.2	49	4.	2.0
Mycogen	8N520DM	849 ± 78	374	1739	435	7.4	27.3	42.5	4.3	45	4.	2.7
Mycogen	8N386CL	$827 \pm 73$	334	1787	328	7.5	27.5	42.8	3.7	48	4.	2.5
Pioneer	63M91	$826 \pm 78$	184	2099	196	8.1	28.1	43.8	4.7	49	1.3 6.1	2.5
Dekalb	DKF 38-30	$826 \pm 73$	312	1937	229	7.9	31.4	4.14	8.4	46	<del>1</del> .3	2.8
Mycogen	8N462DM	$819 \pm 73$	634	1680	142	7.7	30.1	44.3	5.0	46	<u>4</u> .	3.3
Triumph	s678	$814 \pm 74$	331	1744	368	7.4	30.3	44.2	4.0	43	2.1	2.7
Mycogen	8D310	$668 \pm 73$	334	1203	468	7.5	24.7	38.9	4.2	20	<del>1</del> .3	1.7
Triumph	999	$650 \pm 80$	232	1535	181	7.4	29.7	43.6	4.2	47	2.0	2.3
Triumph	620 CL	$631 \pm 95$	236	1623	32	7.4	27.1	42.2	3.3	47	1.7	2.8
Dekalb	DKF 38-80 CL	$564 \pm 78$	204	1334	154	7.5	26.9	40.1	4.3	42	1.2	3.0
Croplan	3080 DMR NS	$513 \pm 78$	290	1020	230	9.7	24.3	42.6	4.3	42	1.2	3.0
Mycogen	8N337DM	$479 \pm 72$	391	964	85	7.8	24.7	41.2	4.7	44	<b>4</b> .	2.2
High Oleic	High Oleic Hybrids ¶											
Triumph	847HO CL	$925 \pm 79$	227	2130	419	7.4	28.1	43.4	4.3	25	2.2	2.7
Triumph	855HO	$925 \pm 79$	354	2250	171	7.7	24.7	43.8	4.7	54	2.4	2.2
Croplan	378 DMR HO	$829 \pm 73$	306	1762	419	8.0	30.9	43.3	5.5	20	<u>+</u>	3.0
Mycogen	8H419CL	$816 \pm 79$	355	1740	351	7.4	24.9	42.7	4.2	46	<del>1</del> .3	2.7
Pioneer	64H41	$807 \pm 72$	204	1908	311	7.6	30.5	42.6	4.0	47	<del>د</del> .	3.0
Triumph	845HO	$743 \pm 72$	324	1518	388	7.3	26.7	4.4	4.0	45	2.4	2.0
Triumph	859HO CL	$728 \pm 100$	394	1634	156	8.2	22.6	41.0	8.4	40	1.2	3.5
Confection	Confectionary (non-oilseed)	Hybrids										
Triumph	777 C ‡	36	636	1956	387	7.4	20.2	29.2	5.2	54	2.4	1.3
Avg.		829	387	1682	329	7.7	28.2	42.4	4.5	46	1.5	2.5
L.S.D05		199	269	449	285							
C.V. (%)		24.6	42.2	15.8	44.1							
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1:toiom /00/ 20 toting		- Langeritand			0 +004 020100	to thought of	, ollinorayi	the contract of	20000		

§ Severe heat and drought at Knoxville during the growing season. Lodging = 1 to 5 scale; where 1 = 95% of plants errect; 2.5 = ~50% of plants leaning at an angle ≥ 45°; 5 = 95+% of plants leaning at an angle ≥ 45°. ¶CL = tolerant to imazamox (Beyond) herbicide # Confectionary Type † All yields adjusted to 10% moisture lbs / ac ÷ 25 = bushels per acre

Bird Damage = 1 to 5 scale; where 1 = 95%+ of plant seed remaining; 2.5 = ~50% of plant seed eaten; 5 = 95+% of plant seed eaten.

Table 10. Mean yields and agronomic characteristics of 25 sunflower hybrids evaluated in three environments in Tennessee during 2006.

		Avg. Yield†			Double	Moisture	Test	Head	Plant	ı	Bird
		± Std. Err.	:		Crop	at Harvest	Weight	Diameter	Height	Lodging	Damage
Brand	Hybrid	(n=3)	Knoxville	Milan	Milan	(n=1)	(n=3)	(n=1)	(n=3)	(n=3)	(n=1)
			lbs/a			%	nq/sql	.⊑	. <u>:</u>	score	%
<b>Traditional Hybrids</b>	<u> </u>										
Dekalb	DKF 39-01	$1688 \pm 160$	2803	1154	1108	15.4	25.5	6.3	22	<b>1</b> .	1.3
Mycogen	8N270	$1478 \pm 159$	1727	1293	1415	17.2	26.7	5.7	55	1.3	1.7
Mid Oleic Hy	Mid Oleic Hybrids (NuSun) ¶										
Mycogen	8N453	$1628 \pm 160$	2307	840	1736	14.7	28.6	6.3	61	1.2	1.7
Triumph	s678	$1573 \pm 173$	1756	1454	1509	17.9	25.7	5.5	51	1.2	1.5
Triumph	636	$1502 \pm 159$	2357	992	1158	18.4	25.2	5.7	29	1.6	1.5
Triumph	645	$1405 \pm 159$	1987	1346	883	16.3	24.1	5.0	62	1.7	1.5
Mycogen	8N386CL	$1397 \pm 172$	1302	1685	1204	13.3	25.8	5.8	62	1.1	1.3
Mycogen	8D310	$1389 \pm 160$	1467	1257	1442	12.2	22.3	4.6	26	1.1	1.5
Dekalb	DKF 35-10	$1318 \pm 160$	1222	928	1804	17.9	26.4	4.4	29	1.3	1.5
Dekalb	<b>DKF 38-80 CL</b>	$1316 \pm 160$	1498	1226	1223	17.2	25.2	5.0	51	1.3	1.5
Mycogen	8N462DM	$1298 \pm 159$	1060	1345	1489	17.1	27.2	5.3	28	1.1	2.2
Triumph	658	$1290 \pm 160$	1230	1189	1453	16.2	24.5	5.3	61	1.8	1.3
Dekalb	DKF 38-30	$1235 \pm 173$	1237	1141	1327	14.2	26.2	5.1	09	1.5	1.7
Triumph	620 CL	$1189 \pm 173$	1061	1225	1282	14.1	26.3	4.6	26	4.1	2.0
Mycogen	8N337DM	$1082 \pm 160$	992	1203	1050	11.7	26.3	4.3	22	1.6	2.0
Triumph	960 CL	$1039 \pm 159$	840	922	1355	21.3	24.5	8.4	28	1.2	1.8
Triumph	<b>TRX S5322CL</b>	$1037 \pm 173$	1078	1207	826	16.2	24.2	4.6	46	1.6	1.5
Dekalb	DKF 37-31	$1021 \pm 173$	1177	756	1130	14.7	24.4	4.5	54	1.3	1.5
Mycogen	8N520DM	$1018 \pm 160$	1001	763	1288	15.0	25.4	5.2	28	1.2	1.8
Triumph	s672	$976 \pm 159$	971	787	1170	12.9	26.2	4.6	45	1.2	1.5
Triumph	999	$910 \pm 159$	877	629	1174	12.2	25.5	5.6	22	1.7	1.5
High Oleic Hybrids	lybrids ¶										
Triumph	845HO	$1432 \pm 173$	1150	1850	1297	19.5	23.2	4.9	09	1.3	1.5
Mycogen	8N419CL	$1334 \pm 159$	1370	1365	1269	12.6	25.9	5.2	62	1.0	2.0
Mycogen	8H350DM	$1265 \pm 185$	1125	1290	1379	10.9	25.0	5.0	28	1.2	1.5
Confectiona	Confectionary (non-oilseed) Hybrids	lybrids									
Triumph	777 C ‡	$1392 \pm 159$	1210	1122	1845	18.2	16.2	5.2	64	1.1	1.7
Avg.		1289	1409	1164	1290	15.5	25.1	5.1	22	1.3	1.6
L.S.D <sub>05</sub>		442	1027	755	504						
C.V. (%)		35.9	43.1	37.3	23.8						
† All yields adju.	† All yields adjusted to 10% moisture		# Confectionary Type	уре	¶ CL = tolerar	¶ CL = tolerant to imazamox (Beyond) herbicide	leyond) herbi		lbs / ac ÷ 25 =	lbs / ac ÷ 25 = bushels per acre	cre

Lodging = 1 to 5 scale; where 1 = 95% of plants errect; 2.5 = -50% of plants leaning at an angle  $\geq 45$ °; 5 = 95+% of plants leaning at an angle  $\geq 45$ °. ¶CL = tolerant to imazamox (Beyond) herbicide Bird Damage = 1 to 5 scale; where 1 = 95% + of plant seed remaining; 2.5 = ~50% of plant seed eaten; 5 = 95 + % of plant seed eaten. # Confectionary Type † All yields adjusted to 10% moisture

Table 11. Mean yields and agronomic characteristics of 17 sunflower hybrids evaluated in two environments in Tennessee during 2005.

		Ava Viold+			Moieture	Toct	Нозд	Dlant		Ring
		- 10 - 10 - 10 - 10 - 10 - 10 - 10 - 10				Woight	Diamotor.	- tale	201200	Since Operation
Brand	Hybrid	± 3td. Eff. (n=2)	Knoxville	Milan	at narvest (n=2)	(n=1)	(n=1)	neigiit (n=2)		Dailiage (n=1)
			lbs/a		%	nq/sql	.⊑	ï.	score	%
Traditional Hybrids	brids									
Mycogen	SF270	$1557 \pm 162$	1601	1512	7.4	26.2	5.0	69	3.0	3.0
Mid Oleic Hybrids (NuSun)	rids (NuSnn) ¶									
Triumph	989	$1750 \pm 163$	2250	1250	7.8	23.6	6.3	20	3.2	2.0
Dekalb	DKF 35-10	$1575 \pm 162$	1761	1389	7.7	25.3	6.4	89	2.4	2.8
Monsanto	MH 4331 B	$1571 \pm 163$	1980	1162	7.6	25.4	5.3	29	2.6	2.5
Triumph	658	$1539 \pm 164$	1559	1519	7.8	23.2	5.0	69	3.4	3.5
Mycogen	8N429 CL	$1482 \pm 162$	1964	666	7.5	24.7	4.9	72	3.0	2.3
Mycogen	8N352	$1475 \pm 163$	1568	1381	7.4	26.9	4.7	99	3.3	4.0
Dekalb	DKF 38-30	$1454 \pm 162$	1492	1417	7.6	25.3	4.2	89	2.3	3.3
Advanta Pacific		$1428 \pm 162$	1458	1399	7.4	24.5	8.4	69	2.5	1.5
Triumph	645	$1387 \pm 163$	1580	1195	7.3	22.8	5.3	72	3.8	3.5
Triumph	620 CL	$1153 \pm 162$	206	1399	7.4	26.6	4.5	65	3.2	4.0
Dekalb	<b>DKF 38-80 CL</b>	$1126 \pm 164$	1396	822	7.5	24.1	5.1	63	3.2	2.0
Triumph	s672	$1117 \pm 164$	966	1239	7.4	26.8	4.5	20	2.1	3.2
Mycogen	8N251	$1074 \pm 161$	1392	755	7.4	23.0	8.4	09	2.8	<del>1</del> .8
Advanta Pacific	: 461 NS	$1057 \pm 164$	806	1206	7.6	23.0	4.1	64	2.2	2.3
Triumph	CT 099	$934 \pm 163$	649	1219	8.1	23.5	4.4	89	3.5	4.0
Confectionary	Confectionary (non-oilseed) Hybrids	<u>brids</u>								
Triumph	777 C ‡	$1249 \pm 162$	1717	781	8.9	16.0	5.5	79	3.6	1.5
Avg.		1344	1473	1215	9.2	24.2	4.9	29	2.9	2.8
L.S.D <sub>05</sub>		440	830	341						
C.V. (%)		28.3	33.8	16.8						

# Confectionary Type

TCL = tolerant to imazamox (Beyond) herbicide

lbs / ac ÷ 25 = bushels per acre

Lodging = 1 to 5 scale; where 1 = 95% of plants errect; 2.5 = ~50% of plants leaning at an angle ≥ 45°; 5 = 95+% of plants leaning at an angle ≥ 45°.

Bird Damage = 1 to 5 scale; where 1 = 95% + of plant seed remaining; 2.5 = ~50% of plant seed eaten; 5 = 95 + % of plant seed eaten.

Table 12. Mean yields and agronomic characteristics of 15 sunflower hybrids evaluated in two environments in Tennessee during 2004.

		Avg. Yield†			Moisture	Test	Head	Plant	•	Bird
		± Std. Err.			at Harvest	Weight	Diameter	Height	Lodging	Damage
Brand	Hybrid	(n=2)	Knoxville	Milan	(n=1)	(n=2)	(n=2)	(n=2)	(n=2)	(n=1)
		***************************************	lbs/a		%	nq/sql	ï.	ï.	score	%
Traditional Hybrids	Hybrids									
Mycogen	SF270	$801 \pm 155$	206	897	9.4	33.2	5.8	20	3.1	3.0
Mid Oleic Hy	//Id Oleic Hybrids (NuSun) ¶									
Triumph	636	na	1	1776	9.3	29.8	7.3	72	3.1	1
Mycogen	8N429 CL	$1624 \pm 137$	1520	1729	9.2	30.3	6.4	89	2.3	1.5
Deklab	DKF 38-80 CL	na	1	1772	9.5	30.5	6.5	29	2.8	1
Monsanto	MH 4231	$1430 \pm 154$	2222	639	11.2	31.3	5.9	65	2.3	1.5
Interstate	HySun 450	$1376 \pm 137$	1440	1313	9.6	31.5	6.3	99	2.0	2.0
Triumph	620 CL	$1329 \pm 155$	1520	1138	9.6	31.8	6.1	69	3.1	2.0
Mycogen	8N251	$1245 \pm 137$	1116	1373	9.5	33.4	6.1	99	2.5	3.0
Triumph	645	na		1015	9.3	31.2	9.9	20	3.8	
Interstate	4880 NS/CL	$1184 \pm 137$	1175	1192	9.5	32.3	6.7	69	2.4	1.5
Deklab	DKF 38-30	$1180 \pm 137$	1296	1063	10.5	32.3	6.3	89	2.5	3.5
Deklab	DKF 35-10	$1119 \pm 198$	1801	438	9.6	33.9	5.5	69	4.1	1.5
Triumph	299	$1091 \pm 137$	1743	438	9.5	32.8	5.9	51	2.7	2.5
Triumph	s675	$1014 \pm 154$	1057	971	10.0	32.9	5.7	48	4.8	4.0
Mycogen	8N352	$933 \pm 155$	536	1331	9.7	33.5	6.2	99	2.4	2.0
Avg.		1267	1410	1154	9.7	31.9	6.2	92	2.8	2.4
L.S.D05		428	944	444						
C.V. (%)		26.5	30.4	22.2						

# Confectionary Type

TCL = tolerant to imazamox (Beyond) herbicide

lbs / ac ÷ 25 = bushels per acre

Lodging = 1 to 5 scale; where 1 = 95% of plants errect; 2.5 = -50% of plants leaning at an angle  $\ge 45$ °; 5 = 95 + % of plants leaning at an angle  $\ge 45$ °.

Bird Damage = 1 to 5 scale; where 1 = 95% + of plant seed remaining; 2.5 = ~50% of plant seed eaten; 5 = 95 + % of plant seed eaten.

Table 13. Mean yields and agronomic characteristics of 21 sunflower hybrids evaluated in three environments for two years (2006-2007) in Tennessee.

III Tennessee.	ee.										
		Avg. Yield†			Double	Moisture	Test	Head	Plant		Bird
		± Std. Err.			Crop	at Harvest	Weight	Diameter	Height	Lodging	Damage
Brand	Hybrid	(n=6)	Knoxville	Milan	Milan	(n=4)	(n=6)	(n=2)	(u=e)	(n=6)	(n=2)
			lbs/a			%	nq/sql	'n.	Ü.	score	%
<b>Traditional Hybrids</b>	Hybrids										
Dekalb	DKF 39-01	$1200 \pm 92$	1544	1398	629	9.5	27.8	5.2	51	4.1	2.1
Mycogen	8N270	$1026 \pm 88$	1120	1131	829	10.0	28.6	5.1	47	1.6	2.1
Mid Oleic H	Mid Oleic Hybrids (NuSun) ¶										
Mycogen	8N453	$1279 \pm 87$	1488	1355	994	9.5	28.0	5.5	54	1.2	2.3
Triumph	929	$1223 \pm 103$	1351	1495	824	10.7	26.7	5.3	22	1.8	2.0
Triumph	s678	$1194 \pm 92$	1043	1599	938	9.3	28.1	4.8	47	1.6	2.1
Triumph	645	$1158 \pm 87$	1194	1572	208	8.6	27.6	4.6	22	1.8	1.8
Mycogen	8N386CL	$1112 \pm 91$	818	1736	782	0.6	26.6	4.7	22	1.3	6.
Dekalb	DKF 35-10	$1092 \pm 88$	932	1280	1064	10.4	27.8	4.8	54	<b>1</b> .	1.8
Mycogen	8N462DM	$1058 \pm 88$	847	1512	815	10.1	28.7	5.1	25	1.3	2.8
Dekalb	DKF 38-30	$1030 \pm 91$	774	1539	778	9.5	28.9	5.0	53	<del>1</del> .	2.3
Mycogen	8D310	$1028 \pm 88$	006	1230	922	8.7	23.7	4 4.	53	1.2	1.6
Triumph	960 CL	$962 \pm 91$	714	1317	826	11.2	25.0	4.6	53	1.3	2.5
Dekalb	DKF 37-31	$960 \pm 94$	848	1265	208	9.3	27.5	4.8	20	<del>1</del> .	1.8
Dekalb	DKF 38-80 CL	$940 \pm 91$	851	1280	688	10.0	26.0	4.7	46	1.3	2.3
Mycogen	8N520DM	$934 \pm 91$	688	1251	861	9.3	26.5	4.8	25	1.3	2.3
Triumph	620 CL	$910 \pm 106$	649	1424	629	8.6	26.8	4.0	53	1.6	2.4
Triumph	999	$780 \pm 92$	222	1107	829	8.6	27.8	4.9	21	1.9	9.1
	8N337DM	$780 \pm 88$	691	1084	266	8.8	25.5	4.5	51	1.5	2.1
High Oleic F	Hybrids ¶										
Triumph	845HO	$1088 \pm 91$	737	1684	842	9.5	25.2	4.5	53	1.8	1.8
Mycogen	8H419CL	$1075 \pm 92$	863	1552	810	8.7	25.4	4.7	24	1.2	2.3
Confections	Confectionary (non-oilseed) Hybrids	brids									
Triumph	777 C ‡	$1193 \pm 92$	923	1539	1116	10.1	18.1	5.2	29	1.8	1.5
Avg.		1049	930	1398	818	9.5	26.5	4.8	52	1.5	2.1
L.S.D05		290	262	208	380						
C.V. (%)		33.9	50.4	24.1	30.5						
	200 to 200 /00/ at 12 cts										

<sup>†</sup> All yields adjusted to 10% moisture

<sup>‡</sup> Confectionary Type

<sup>¶</sup>CL = tolerant to imazamox (Beyond) herbicide

lbs / ac ÷ 25 = bushels per acre

Lodging = 1 to 5 scale; where 1 = 95% of plants errect; 2.5 = ~50% of plants leaning at an angle ≥ 45°; 5 = 95+% of plants leaning at an angle ≥ 45°. Bird Damage = 1 to 5 scale; where 1 = 95% + of plant seed remaining; 2.5 = ~50% of plant seed eaten; 5 = 95 + % of plant seed eaten.

Table 14. Mean yields and agronomic characteristics of nine sunflower hybrids evaluated in two environments for three years (2005-2007) in Tennessee.

		Avg. Yield†			Moisture	Test	Head	Plant		Bird
		± Std. Err.			at Harvest	Weight	Diameter	Height	Lodging	Damage
Brand	Hybrid	(u=e)	Knoxville	Milan	(n=5)	(n=5)	(n=3)	(u=6)	(n=6)	(n=3)
			lbs/a		%	nq/sql	in.	.⊑	score	%
Mid Oleic H	Mid Oleic Hybrids (NuSun) ¶									
Triumph	636	$1532 \pm 95$	1651	1413	9.8	26.0	5.6	28	2.1	2.0
Triumph	645	$1384 \pm 96$	1322	1446	9.1	25.9	4.8	61	2.1	2.3
Dekalb	DKF 35-10	$1262 \pm 95$	1208	1316	9.7	27.2	4.8	28	1.6	2.1
Dekalb	DKF 38-30	$1256 \pm 99$	1013	1498	8.9	27.7	4.7	28	1.6	2.6
Dekalb	DKF 37-31	$1228 \pm 99$	1225	1231	8.7	26.9	4.9	22	1.6	2.0
Dekalb	DKF 38-80 CL	$1086 \pm 96$	1033	1138	9.4	25.1	4.8	25	6.1	2.2
Triumph	620 CL	$1075 \pm 99$	735	1416	8.3	27.3	4.1	28	2.1	2.9
Triumph	12 099	988 <del>+</del> 96	692	1284	10.5	25.1	4.5	20	6.1	3.0
Confection	Confectionary (non-oilseed) Hybrids	ybrids								
Triumph	777 C ‡	$1237 \pm 95$	1188	1286	10.1	16.8	5.3	29	2.2	1.5
Avg.		1228	1119	1337	9.4	25.3	4.9	28	1.9	2.3
L.S.D <sub>05</sub>		379	612	455						
C.V. (%)		32.5	44.5	22.9						
11 - 1-1-1-1										

# Confectionary Type

¶CL = tolerant to imazamox (Beyond) herbicide

lbs / ac ÷ 25 = bushels per acre

Lodging = 1 to 5 scale; where 1 = 95% of plants errect; 2.5 = -50% of plants leaning at an angle  $\ge 45\%$ ; 5 = 95% of plants leaning at an angle  $\ge 45\%$ .

Bird Damage = 1 to 5 scale; where 1 = 95% + of plant seed remaining; 2.5 = -50% of plant seed eaten; 5 = 95 + % of plant seed eaten.

Table 15. Mean yields and agronomic characteristics of six sunflower hybrids evaluated in two environments for four years (2004-2007) in Tennessee.

		Avg. Yield†			Moisture	Test	Head	Plant		Bird
		± Std. Err.			at Harvest	Weight	Diameter	Height	Lodging	Damage
Brand	Hybrid	(n=8)	Knoxville	Milan	(n=6)	(n=7)	(n=5)	(n=8)	(n=8)	(n=4)
			lbs/a		%	nq/sql	in.	in.	score	%
Mid Oleic	Mid Oleic Hybrids (NuSun) ¶									
Triumph	636	$1637 \pm 90$	1770	1504	9.7	26.6	6.3	62	2.4	2.0
Triumph	645	$1346 \pm 90$	1353	1338	9.2	26.8	5.5	63	2.6	2.3
Dekalb	DKF 38-30	$1237 \pm 83$	1084	1390	9.2	29.1	5.3	61	1.8	2.9
Dekalb	DKF 35-10	$1227 \pm 90$	1356	1097	9.7	28.7	5.1	61	2.2	2.0
Dekalb	DKF 38-80 CL	$1193 \pm 90$	1089	1297	9.4	26.0	5.5	54	2.1	2.1
Triumph	620 CL	$1139 \pm 85$	931	1346	8.5	28.6	4.9	61	2.3	2.8
Avg.		1296	1264	1329	9.3	27.6	5.4	09	2.2	2.3
L.S.D <sub>05</sub>		368	602	437						
C.V. (%)		31.6	42.0	22.9						

# Confectionary Type

¶ CL = tolerant to imazamox (Beyond) herbicide

lbs / ac ÷ 25 = bushels per acre

Lodging = 1 to 5 scale; where 1 = 95% of plants errect; 2.5 = -50% of plants leaning at an angle  $\geq 45$ °; 5 = 95+% of plants leaning at an angle  $\geq 45$ °.

Bird Damage = 1 to 5 scale; where 1 = 95% + 0f plant seed remaining; 2.5 = ~50% of plant seed eaten; 5 = 95 + % of plant seed eaten.

#### References

Aiken, R. M. 2005. Applying thermal time scales to sunflower development. Agron. J. 97:746-754. http://agron.scijournals.org/cgi/content/full/97/3/746

American Soybean Association (ASA). 2008. Soy Stats: World vegetable oil consumption 2007. American Soybean Assoc., St. Louis, MO. http://www.soystats.com/2008/page\_35.htm

Berglund, D. R. 2007a. Introduction. Pp. 1-5. In D. R. Berglund (ed.). Sunflower Production. Publication A-1331. North Dakota State Univ., Fargo, ND. http://www.sunflowernsa.com/uploads/Sunflower\_Production\_Handbook\_2007.pdf

Berglund, D. R. (ed.). 2007b. Sunflower Production. Publication A-1331. North Dakota State Univ., Fargo, ND. http://www.sunflowernsa.com/uploads/Sunflower\_Production\_Handbook\_2007.pdf

Bradley, C., S. Markell, and T. Gulya. 2007. Pest management: Diseases of sunflower. Pp. 54-77. In D. R. Berglund (ed.). Sunflower Production. Publication A-1331. North Dakota State Univ., Fargo, ND. http://www.sunflowernsa.com/uploads/Sunflower\_Production\_Handbook\_2007.pdf

Crites, G. D. 1993. Domesticated sunflower in fifth millennium b.p. temporal context: New evidence from Middle Tennessee. Am. Antiq. 58:146-148.

Dumler, T. J., D. M. O'Brien, and B. L. S. Olson. 2008. Sunflower cost-return budget in western Kansas. Farm Mgmt. Guide MF-887. Kansas State Univ., Manhattan, KS. www.agmanager.info

Jardine, D.J. 2009. Diseases. Pp. 31-35. In High Plains Sunflower Production Handbook. MF-2384. Kansas State Univ., Manhattan, KS. http://www.ksre.ksu.edu/library/crpsl2/mf2384.pdf

Johnson, J.J., R.F. Meyer, J.M. Krall, J.P. Shroyer, A.J. Schlegel, J.S. Falk, and C.D. Lee. 2005. Agronomic Practices. Pp. 1-4. In High Plains Sunflower Production Handbook. MF-2384. Kansas State Univ., Manhattan, KS. <a href="http://www.ksre.ksu.edu/library/crpsl2/mf2384.pdf">http://www.ksre.ksu.edu/library/crpsl2/mf2384.pdf</a>

Knodel, J., and L. Charlet. 2007. Pest management: insects. Pp. 26-53. In D. R. Berglund (ed.). Sunflower Production. Publication A-1331. North Dakota State Univ., Fargo, ND. http://www.sunflowernsa.com/uploads/Sunflower\_Production\_Handbook\_2007.pdf

Latzke, J. M. 2007. Research continues into effective blackbird control. Mar. 8, 2007. High Plains/Midwest Ag Journal, Dodge City, KS. http://www.hpj.com/archives/2007/mar07/mar12/Researchcontinuesintoeffect.cfm

Linz, G. M., and J. Hanzel. 2007. Birds. Pp. 85-88. In D. R. Berglund (ed.). Sunflower Production. Publication A-1331. North Dakota State Univ., Fargo, ND. http://www.sunflowernsa.com/uploads/Sunflower\_Production\_Handbook\_2007.pdf

Linz, G. M., R. A. Dolbeer, J. J. Hanzel, and L. E. Huffman. 2006. Controlling blackbird damage to sunflower and grain crops in the northern Great Plains. Agric. Information Bull. 679. U.S. Dept. of Agriculture, Animal and Plant Inspection Service (USDA-APHIS), Jamestown, ND. http://www.npwrc.usgs.gov/resource/birds/blkbird/index.htm

Linz, G. M., A. A. Slowik, L. B. Penry, and H. J. Homan. 2004. Evaluation of registered sunflower insecticides as candidate blackbird repellents. U.S. Department of Agriculture-Animal and Plant Inspection Services (USDA-APHIS), Bismarck, ND. http://www.aphis.usda.gov/wildlife\_damage/nwrc/publications/04pubs/linz041.pdf

McMullen, M. P., and C. A. Bradley. 2007. 2007 North Dakota Field Crop Fungicide Guide. North Dakota State Univ., Fargo, ND. http://www.ag.ndsu.edu/pubs/plantsci/pests/pp622/pp622.pdf

Meyer, R., D. Belshe, J. Falk, S. Patten, and D. O'Brien (Ed.). 2009. High Plains Sunflower Production Handbook. MF-2384. Kansas State Univ., Mahattan, KS. http://www.ksre.ksu.edu/library/crps12/mf2384.pdf

Myers, R. L. 2008. Sunflower: A Native Oilseed with Growing Markets. Thomas Jefferson Agricultural Institute, Columbia, MO. http://www.jeffersoninstitute.org/pubs/sunflower\_guide.pdf

Nafziger, E. D. 2008. Chapter 3: Soybeans. In R. Hoeft and E. Nafziger (eds.). Illinois Agronomy Handbook, Dept. of Crop Sciences, Univ. of Illinois, Champagne, IL. http://iah.aces.uiuc.edu/index.php?ch=drop.html&s=soy

North Dakota Agricultural Weather Network (NDAWN). 2008. Sunflower development and growing degree days (GDD). North Dakota State Univ., Fargo, ND. http://ndawn.ndsu.nodak.edu/help.html?topic=sunflowergdd

ProCrop. 2008. Crop production information: Sunflower menu. North Dakota State Univ., Fargo, ND. http://www.ag.ndsu.edu/procrop/sun/index.htm

Putnam, D. H., E. S. Oplinger, D. R. Hicks, B. R. Durgan, D. M. Noetzel, R. A. Meronuck, J. D. Doll, and E. E. Schulte. 1990. Sunflower. Alternative Field Crops Manual. Univ. of Minnesota, St. Paul, MN, and Univ. of Wisconsin, Madison, WI. http://www.hort.purdue.edu/newcrop/afcm/sunflower.html

Putt, E.D. 1997. Early History of sunflower. p. 2-20. In A.A. Schneiter (ed.) Sunflower Technology and Production. Agron. Monogr. 35, ASA, CSSA, SSSA, Madison, WI.

Rankin, Al, Jr. 2007. Sunflower weed control recommendations for Mississippi. Publ. 2434. Miss. State Univ., Mississippi State, MS. http://msucares.com/pubs/publications/p2434.pdf

Savoy, H.J. and D. Joines. 2009. BEES Info #100 Lime and Fertilizer Recommendations for the Various Crops of Tennessee. Chapter II Agronomic Crops. http://soilplantandpest.utk.edu/pdffiles/soiltestandfertrecom/chap2-agronomic\_mar2009.pdf

Seiler, G.J. and L.H. Rieseberg. 1997. Systematics, origin, and germplasm resources of the wild and domestic sunflower. p. 21-65. In A.A. Schneiter (ed.) Sunflower Technology and Production. Agron. Monogr. 35, ASA, CSSA, SSSA, Madison, WI.

Sloderbeck, P.E., A. Gebre-Amlak, J.P. Michaud, F.B. Peairs, and G.L. Hein. 2009. Insect Pest Identification and Control. Pp. 17-26. In High Plains Sunflower Production Handbook. MF-2384. Kansas State Univ., Manhattan, KS. http://www.ksre.ksu.edu/library/crpsl2/mf2384.pdf

Sloderbeck, P. E., J. P. Michaud, and R. J. Whitworth. 2008. Sunflower insect management 2008. MF-814. Kansas State Univ., Manhattan, KS.

Steckel, L. 2008. Sunflower weed control. In T. C. Mueller (ed.). 2008 Weed Control Manual for Tennessee. Univ. of Tennessee, Knoxville, TN. www.weeds.utk.edu

U.S. Department of Agriculture-Economics Research Service (USDA-ERS). 2008. U.S. fertilizer use and price. USDA-ERS, Washington, D.C. http://www.ers.usda.gov/Data/FertilizerUse/

U.S. Department of Agriculture-National Agricultural Statistics Service (USDA-NASS). 2007. Sunflower: Production per harvested acre by county. USDA-NASS, Washington, D.C. <a href="http://www.nass.usda.gov/Charts\_and\_Maps/Crops\_County/sf-pr.asp">http://www.nass.usda.gov/Charts\_and\_Maps/Crops\_County/sf-pr.asp</a>

U.S. Department of Agriculture-National Agricultural Statistics Service (USDA-NASS). 2008. U.S. and all states data-crops. USDA-NASS, Washington, D.C. http://www.nass.usda.gov/QuickStats/

U.S. Department of Agriculture-National Agricultural Statistics Service (USDA-NASS). 2009. U.S. and all states data-crops. USDA-NASS, Washington, D.C. http://www.nass.usda.gov/QuickStats/

U.S. Department of Commerce-National Climatic Data Center (USDC-NCDC). 2008. Record of Climatological Observations. USDC-NCDC, Washington, D.C. http://www7.ncdc.noaa.gov/IPS/coop/coop.html

Vigil, M. F., G.W. Hergert, and D.B. Mengel. 2009. Nutrient Management. Pp. 4-8. In High Plains Sunflower Production Handbook. MF-2384. Kansas State Univ., Manhattan, KS. http://www.ksre.ksu.edu/library/crpsl2/mf2384.pdf

Zheljazkov, V. D., B. A. Vick, M. W. Ebelhar, N. Buehring, B. S. Baldwin, T. Astatkie, and J. F. Miller. 2008. Yield, oil content, and composition of sunflower grown at multiple locations in Mississippi. Agron. J. 100:635-642. http://agron.scijournals.org/cgi/reprint/100/3/635



This publication is posted online at: http://varietytrials.tennessee.edu and www.UTCrops.com