



2-11-2011

SP290-D Insects: Twospotted Spider Mites

Frank A. Hale

Follow this and additional works at: http://trace.tennessee.edu/utk_agexdise

 Part of the [Agricultural Science Commons](#), and the [Agriculture Commons](#)

Recommended Citation

"SP290-D Insects: Twospotted Spider Mites," Frank A. Hale,
SP290-D
, http://trace.tennessee.edu/utk_agexdise/61

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#). This Home, Lawn & Garden Insects & Pests is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Insects, Pests, Plant Diseases and Weeds by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Insects



Twospotted Spider Mites

Frank A. Hale, Professor

*Originally developed by Harry Williams, Professor Emeritus
Entomology and Plant Pathology*

Twospotted spider mites occur as injurious pests on more than 180 host plants, including over 100 cultivated species. Violet, chickweed, pokeweed, wild mustard and blackberry are common foci from which infestations develop on nearby ornamental plants.

Spider mites are not actually insects. More closely related to spiders than insects, they derive their name from the thin web that some mite species spin.

Mites pierce the outer leaf surface with their sharp, slender mouthparts as they feed. When they extract the sap, the tissue of the leaf collapses in the area of the puncture. Soon a chlorotic (yellow) spot forms at each feeding site. After a heavy attack, an entire plant may become yellowed, bronzed, partially defoliated or completely killed. The mites may spin so much webbing over the plant that it becomes entirely covered.


Spider mites pass through four separate life stages – egg, larva, nymph and adult. The eight-legged adult can be rusty green, greenish amber, yellowish, brick red or rust to almost black. There are usually two (sometimes four) black spots on the top of the 0.5-0.7 mm long mites. A good hand lens is needed to see this. The spherical eggs, laid on the leaf, range from transparent and colorless to a milky white or opaque straw yellow.

The six-legged larva is colorless, pale green or yellow. Similar to the adult except in size, the nymph has eight legs and is pale green to brownish-green. Large spots of black may develop on each side.

Twospotted spider mites overwinter as adult females in protected sites in the soil and duff, or on ground hugging plants, primarily weeds. In mild weather they continue to feed and lay eggs, although development in the winter is much slower than in the summer.

Adults mate soon after emerging in warm weather, and females soon begin to lay eggs. Each female may lay

more than 100 eggs in her lifetime and up to 19 eggs per day. Development is rapid in hot, dry weather. Each generation may take as many as 20 or as few as five days to mature.



Control Measures

Inspect plants for mites at frequent regular intervals (at least once each week). Observe the plants for webbing between the leaf petiole, stems and leaf edges. Observe the plant for chlorotic discoloration, stippling, and eggs or cast skins on the underside of the leaf. A hand lens or other magnification is a useful aid for seeing mites on the foliage.

Hold a white card under the plant foliage. Shake the foliage to dislodge mites onto the card. Circle the small dots that appear with a pencil mark. Hold the card in the sunlight and observe. Mites will crawl out of the circles. Dust and dirt will not.

Spray the under surfaces of leaves with miticide at five-day intervals until mite infestations are suppressed. Usually two to three applications are required. Do not use

horticultural oil (various brand names) more than once per week or apply captan two weeks before or after an oil spray. Some miticides can only be used once per year or need to be applied early in the season at specific growth

stages of the host plant. Follow label directions. Do not apply any miticide to ornamental, vegetable or fruit plants that are not listed on the label.

| Crop | Miticide Recommendations |
|--|---|
| Ornamental Plants Outdoor trees Shrubs and Flowers Interiorscapes Commercial ornamental greenhouses | http://eppserver.ag.utk.edu/redbook/pdf/ornamentalinsects.pdf http://eppserver.ag.utk.edu/redbook/pdf/interiorscapeinsects.pdf http://eppserver.ag.utk.edu/redbook/pdf/ghinsects.pdf |
| Apple | http://www.ces.ncsu.edu/fletcher/programs/apple/2010orchard-management.pdf |
| Peach | http://www.ent.uga.edu/peach/peachguide.pdf |
| Small Fruit (blueberries, bramble, bunch grapes, muscadines, strawberries) | http://www.smallfruits.org/SmallFruitsRegGuide/index.htm |
| Vegetables: Commercial Residential | http://www.citrusandvegetable.com/TheSoutheasternUSVegetableCropHandbook/tabid/791Default.aspx http://eppserver.ag.utk.edu/redbook/pdf/homegardeninsects.pdf |

Precautionary Statement

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store, or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label. Persons who do not obey the law will be subject to penalties.

Disclaimer Statement

Pesticides recommended in this publication were registered for the prescribed uses when printed. Pesticides registrations are continuously reviewed. Should registration of a recommended pesticide be canceled, it would no longer be recommended by the University of Tennessee. Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others which may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product.

THE UNIVERSITY of TENNESSEE 
 INSTITUTE of AGRICULTURE

Visit the UT Extension Web site at
<http://utextension.tennessee.edu/>

SP290-D (Rev) 2/11 11-0090