



April 2010

## W027-Cotton Insects: Stink Bugs

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: [http://trace.tennessee.edu/utk\\_agexcrop](http://trace.tennessee.edu/utk_agexcrop)



Part of the [Entomology Commons](#), and the [Plant Sciences Commons](#)

---

### Recommended Citation

"W027-Cotton Insects: Stink Bugs," The University of Tennessee Agricultural Extension Service, 04-0077 E12-4615,  
[http://trace.tennessee.edu/utk\\_agexcrop/77](http://trace.tennessee.edu/utk_agexcrop/77)

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Insect, Pest and Disease Control - Cotton is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Field & Commercial Crops by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

# Cotton Insects

## Stink Bugs

Scott D. Stewart, Associate Professor  
Entomology and Plant Pathology

### Classification and Description

Stink bugs (Hemiptera: Pentatomidae) are true bugs with piercing-sucking mouthparts and incomplete metamorphosis. The immature stages (nymphs) and adults of several species may damage cotton. Predatory stink bugs may also be found in cotton.

The green stink bug (*Acrosternum hilare*) is the most common species that feeds on cotton in Tennessee. The brown stink bug (*Euschistus servus*) is another common component of the stink bug complex.



Green stink bug

### Hosts, Life History and Distribution

Both green and brown stink bugs are found throughout the state. A third species, the southern green stink bug (*Nezara viridula*), is less common in Tennessee and often confined to the southernmost counties.

Stink bugs have a wide host range that includes many cultivated crops, including corn and soybeans as well as

numerous uncultivated plants. Overwintering adults become active in the spring. Stink bug eggs are characteristically laid in a mass of 20-100. Eggs hatch in 6-7 days during the summer. Nymphs tend to remain aggregated until the third or fourth instar. All species pass through five immature (nymphal) life stages, which require 23-25 days at optimal tempera-



Brown stink bug



Stink bug nymphs

ture. Adults may live several weeks or months. Several generations per year occur in Tennessee.

### Pest Status and Injury

Stink bugs have become a common pest problem in cotton. Bt cotton and boll weevil eradication efforts have reduced the overall number of insecticide applications made to cotton. Because stink bugs are generally susceptible to many insecticides, this reduction in insecticide use has provided an opportunity for stink bugs to establish in cotton fields. Plant-feeding species of





*Boll injury from stink bug feeding*

7-21 days in age. External signs of feeding injury include the appearance of circular black lesions on the surface of bolls (i.e., “catfacing”). These sunken lesions are typically about 1/16th of an inch in diameter. The lint of bolls may be stained, seed may be destroyed and feeding warts may be observed on the internal surface of the boll wall. Damaged bolls may rot due to secondary infection by plant pathogens, or lint production may be reduced in one or more locks. Other true bugs, such as the clouded plant bug and tarnished plant bug, may cause similar injury. External feeding signs on bolls are not always associated with internal damage.

stink bugs are typically seed feeders, and thus, injury to cotton bolls is the primary concern. Economic damage in pre-flowering cotton is very rare. Stink bugs target the seed inside developing bolls, pre-f

### **Management Considerations and Thresholds**

Adult stink bugs are strong fliers and typically migrate into cotton from alternate hosts outside the field, including adjacent fields of maturing corn and soybeans. Stink bugs are difficult to scout. Adults tend to aggregate, and the distribution of stink bugs within a field may be highly concentrated, particularly along borders. Despite being relatively large insects, they can be difficult to see. Stink bugs can be scouted for visually, but most thresholds are based on drop cloth samples or the occurrence of internal symptoms of feeding damage to bolls.

Current UT thresholds call for a treatment when one or more stink bugs are found per drop cloth sample (6 row feet) or when 20 percent or more of thumb-sized bolls show internal evidence of stink bug feeding, including feeding warts or stained lint. Insecticide options are listed in the Tennessee Cotton Insect Control Guide (Extension PB 387). Brown stink bugs are relatively tolerant of pyrethroid insecticides. Thus, organophosphate insecticides such as Bidrin, Orthene or methyl parathion are typically preferred for stink bug control. However, pyrethroid insecticides are effective on green or southern green stink bugs.

For information about the management of the major field crops grown in Tennessee, visit [www.utcrops.com](http://www.utcrops.com)

#### **Precautionary Statement**

To protect people and the environment, pesticides should be used safely. This is everyone’s responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store, or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label. Persons who do not obey the law will be subject to penalties.

#### **Disclaimer Statement**

Pesticides recommended in this publication were registered for the prescribed uses when printed. Pesticides registrations are continuously reviewed. Should registration of a recommended pesticide be canceled, it would no longer be recommended by the University of Tennessee. Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others which may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product.

04-0077 E12-4615

The Agricultural Extension Service offers its programs to all eligible persons regardless of race, religion, color, national origin, sex, age, disability or veteran status and is an Equal Opportunity Employer.

COOPERATIVE EXTENSION WORK IN AGRICULTURE AND HOME ECONOMICS

The University of Tennessee Institute of Agriculture, U.S. Department of Agriculture, and county governments cooperating in furtherance of Acts of May 8 and June 30, 1914.

Agricultural Extension Service Charles L. Norman, Dean