



April 2010

SP503-I-Identifying the Flathead Appletree Borer (*Chrysobothris femorata*) and Other Buprestid Beetle Species in Tennessee

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: https://trace.tennessee.edu/utk_agexgard



Part of the [Entomology Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

"SP503-I-Identifying the Flathead Appletree Borer (*Chrysobothris femorata*) and Other Buprestid Beetle Species in Tennessee," The University of Tennessee Agricultural Extension Service, SP503-I 09-0198, https://trace.tennessee.edu/utk_agexgard/85

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Landscaping - Insects & Diseases is brought to you for free and open access by the UT Extension Publications at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Home Garden, Lawn, and Landscape by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Insects

Identifying the Flatheaded Appletree Borer (*Chrysobothris femorata*) and Other Buprestid Beetle Species in Tennessee

Jason A. Hansen, Frank A. Hale¹, Professor, Entomology and Plant Pathology, and William E. Klingeman, Associate Professor, Plant Sciences

Introduction

The flatheaded appletree borer, *Chrysobothris femorata* (Olivier) (FHATB) is a well-documented, native pest of deciduous trees. It has a wide host range and is particularly problematic in commercial nurseries and urban landscapes, where it can cause rapid decline of economically important hosts. When infestations are high, FHATB has been known to attack healthy trees as well as those stressed by drought, plant disease, mechanical injury and other environmental factors. Its distribution is ubiquitous, covering the entire continental United States and extending into Canada.

Life History

Throughout its range, FHATB completes one generation per year. Adult females lay eggs on the sunny side of trees in late spring and throughout the summer months, ovipositing after a meal of tender bark is taken from the host plant. Eggs hatch about seven days later, at which time larvae chew through the bottom of the egg directly into the host tree, avoiding desiccation and

predation. Inside the host, larvae feed on the actively dividing cambium tissues, in addition to sapwood. Galleries, which are feeding paths made by larvae, maintain optimal humidity for larval growth. Feeding activity continues even in cold winter months on sun-warmed portions of the trunk. When fully developed, larvae bore into the heartwood and form pupal chambers with entrances that are tightly plugged with frass. Frass appears as sawdust particles after passage of the plant tissues through the beetle digestive tract. Pupation occurs in late spring to early summer and lasts 1-2 weeks, after which the adult emerges by cutting a distinctive D-shaped exit hole in the bark (Fig. 1).



Figure 1. FHATB exit holes. Photo by Frank Hale, Entomology and Plant Pathology, University of Tennessee

¹ **Corresponding Author:** Frank Hale, University of Tennessee Soil, Plant and Pest Center, 5201 Marchant Drive, Nashville, TN 37211-5112. Email: fahale@utk.edu



Damage Symptoms

The most serious injury to host plants is caused by larval feeding activity beneath the bark that damages the cambium layer and disrupts the flow of vital nutrients throughout the tree. A single larva is capable of girdling a young tree within one season. Evidence of larval activity can be found under bark of infested trees as sinuous feeding tunnels packed with frass (Fig. 2). Portions of the trunk may show signs of infestation by noticeable oozing of sap. Trees that survive borer attacks are often left scarred and unmarketable (Fig. 3).



Figure 2. Frass-filled galleries. Photo by Nadeer Youssef, TSU Otis Floyd Nursery Research Station, McMinnville, TN

By contrast, adult feeding activity typically causes little damage, though there have been reports of trees being completely defoliated by FHATB when beetle population densities are unusually high. Adult beetles can also gnaw on woody tissues in branch crotches, around bud scars and at the base of leaf petioles.

Identification

Legless larvae are readily recognized by their flattened, sclerotized (hardened) thoracic area, with the remainder of body segments having a slightly off-white color (Fig. 4)

Adult beetles are often described as “bullet-shaped” and are about 0.3–0.6 inches long, with large compound eyes. From above, the back of FHATB is ornamented with several irregular brassy spots (Fig. 5). Beneath the wings,



Figure 3. Tree scar from FHATB. Photo by Frank Hale, Entomology and Plant Pathology, University of Tennessee



Figure 4. Larva of FHATB. Photo by Joshua Basham, TSU Otis Floyd Nursery Research Station, McMinnville, TN


the abdomen is a metallic purple, while the ventral surface is metallic bronze. Forelegs of male beetles are armed with several small teeth (Fig. 6). Antennae are a dark reddish color, 11-segmented, with the last segment tapering apically (Fig. 7).

The FHATB is part of a larger species complex in which all the members have a similar appearance (Fig. 5). In most cases, it is difficult to determine which species is



Figure 5. *Chrysobothris femorata* complex and others buprestids similar in appearance:
**A) *C. femorata*, B) *C. adelpha*, C) *C. rugosiceps*, D) *C. viridiceps*, E) *C. quadriimpressa*, F) *C. shawnee*, G) *C. cribaria*, H) *C. sexsignata*, and I) *C. wintu* (western U.S. species, not found in Tennessee).
 Photos by Jason Hansen, University of Tennessee**

present in the field without closer microscopic examination. Some can be separated using characters on the face of the beetle (Fig. 8). There is some genetic evidence that interbreeding may be occurring between different species in this complex. Eight members of this group occur in the southeastern U.S. and it is still unclear which of them, in



**Figure 6. Male foreleg characteristics of select *Chrysobothris* species: A) *C. sexsignata*, with smooth foreleg and B) *C. femorata*, with spined teeth (arrow).
 Photos by Jason Hansen, University of Tennessee**




Figure 7. Antennal comparison of two species in the *Chrysobothris femorata* “species complex”:
A) *Chrysobothris rugosiceps*, with last antennal segment quadrate and B) *C. femorata*, with last antennal segment tapered. Photos by Jason Hansen, University of Tennessee

addition to FHATB, may be economic and aesthetic threats. Plant host range is known to overlap significantly within this group and correct identification is important before control strategies are implemented.




Figure 8. Face of four common species in the *Chrysobothris femorata* “species complex” with distinguishing characters indicated (arrows): A) *Chrysobothris viridiceps* male, B) *C. rugosiceps* male, with irregular purple spots, C) *C. femorata* male, with semicircular clypeus (compare with D) and deeply cut central notch, and D) *C. adelpha* male, with clypeus having a more straight angle, rather than semicircular (compare with C). Photos by Jason Hansen, University of Tennessee

Management Options

- **Monitoring** – populations of FHATB can be surveyed using purple panel traps (Fig. 9) covered with adhesive pastes (eg. Pestick® or Tanglefoot®) starting mid-April. This new trapping method attracts a wide range of buprestids, including many in the *Chrysobothris femorata* complex, and it is important to know which pest species is a problem when checking them (Figs. 5 & 10). If males are trapped, genitalia can easily be used to separate the six most common species in the southeastern U.S. (Fig. 11). Alternatively, trees can be inspected on an individual basis to determine presence of injury in early spring before emergence begins.
- **Cultural control** – Proper plant management and maintenance are extremely important, as borers prefer to attack stressed or dying trees, while tending to avoid vigorously growing specimens.
- **Biological control** – Natural predators can play a significant role in reducing infestations, by causing reductions of borer populations ranging from 7 to 58 percent, depending on environmental conditions. Though not commercially available, several species of wasps that have been reported to attack FHATB, including *Labena grillator* Say, *Cryptohelcostizus*

chrysobothridis Cushman (Hymenoptera: Ichneumonidae); *Phasgonophora sulcata* Westwood (Hymenoptera: Chalcididae); *Eusandalum* spp. (Hymenoptera: Eupelmidae); *Heterospilus astigmus* Ash. and *Atanycolus rugosiventris* Ash. (Hymenoptera: Braconidae), may occur in Tennessee. Beetles from the clerid family (*Chariessa pilosa* Forst., *C. pilosa onusta* Say) as well as an asilid fly (*Andrenosoma fulvicauda* Say) have also been reared from FHATB.

- **Chemical control** – Several pesticides are available to treat for FHATB by drench treatments applied from February to mid-April that include imidicloprid (e.g., Merit 2F®, Merit 75WSP®, Bayer Advanced Tree & Shrub Insect Control® for ornamental uses in landscapes) used either alone or in addition to cyfluthrin (e.g., Discus® for use in commercial nurseries) or bifenthrin (e.g., Allectus SC® for ornamental uses in landscapes) or Allectus GC SC®, which is labeled for use on golf courses and sod farms. Adult beetles can be controlled by spraying tree trunks in mid-May and late June when borers are at peak emergence with permethrin (e.g., Astro® for ornamental uses in landscapes, Perm-Up 3.2® for use in commercial nurseries), bifenthrin (Onyx® for ornamental uses in landscapes, Onyx Pro® for use in commercial nurseries), or chlorpyrifos (Dursban 4E®, Dursban 50W® for use in commercial nurseries).

Related References & Additional Reading:

- Fenton FA. 1942. The flatheaded apple tree borer (*Chrysobothris femorata* (Olivier)). Bull. B-259. Oklahoma Agricultural Experiment Station, Stillwater. 31p.
- Fisher, WS 1942. A revision of the North American species of Buprestid beetles belonging to the tribe Chrysobothrini. United States Department of Agriculture Miscellaneous Publication 470: 1-274.
- Hansen, J., W.E. Klingeman, J.K. Moulton, J.B. Oliver and M.T. Windham. 2008. Species variation within the *Chrysobothris femorata* “complex” (Flatheaded Appletree Borer): evidence and implications of DNA sequencing. Proc. Ann. Res. Conf. Southern Nursery Assoc. 53: 320-323.
- Oliver J, DC Fare, N Youssef and W Klingeman. 2004. Collection of adult flatheaded borers using multicolored traps. Southern Nursery Association Proceedings. 48: 193-199.



Figure 9. A purple panel trap design, which was first developed by Jason Oliver and others at the Tennessee State University's Otis Floyd Nursery Research Station in McMinnville, TN. Photo by Nadeer Youssef, TSU Otis Floyd Nursery Research Station, McMinnville, TN




Figure 10. Common buprestids attracted to purple panel traps in Tennessee: A) *Acmaeodera pulchella*, B) *A. tubulus*, C) *Ptosima gibbicollis*, D) *Chrysobothris azurea*, E) *C. femorata*, F) *Agrilus bilineatus*, G) *C. dentipes*, H) *Buprestis lineata*, and I) *Dicerca obscura*. Photos by Jason Hansen, University of Tennessee

Potter D.A., GM Timmons and FC Gordon. 1988. Flatheaded apple tree borer (Coleoptera: Buprestidae) in nursery-grown red maples: phenology of emergence, treatment timing, and response to stressed trees. J. Environ. Hort. 6(1): 18-22.

Wellso, SG and GV Manley. 2007. A revision of the *Chrysobothris femorata* (Olivier, 1790) species group from North America, north of Mexico (Coleoptera: Buprestidae). Zootaxa 1652: 1-26.

Wygant ND. 1938. The relation of insects to shelterbelt plantations in the Great Plains. Journal of Forestry 36(10): 1011-1018.




Figure 11. Genitalia characteristics of adult male *Chrysobothris* beetles are critical identification guides for differentiating species within the *Chrysobothris femorata* “species complex.” Examples of species collected on purple panel traps in Tennessee include: A) *Chrysobothris viridiceps*, with constriction (arrow) near the apex of one lobe of the genitalia, B) *C. rugosiceps*, tip of aedeagus (arrow) slightly narrower than the tips on adjacent lobes, C) *C. femorata*, with the tip of aedeagus broad (arrow), D) *C. adelpha*, with spines (arrow) protruding perpendicularly to lateral lobes, E) *C. shawnee*, with narrow lateral lobes that gradually taper to the tip, and F) *C. quadriimpressa*, with lateral lobes constricted (arrow) behind strong lateral bulges. Photos by Jason Hansen, University of Tennessee

Precautionary Statement

To protect people and the environment, pesticides should be used safely. This is everyone’s responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label. Persons who do not obey the law will be subject to penalties.

Disclaimer Statement

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator’s responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.

SP503-I 09-0198

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.
UT Extension provides equal opportunities in programs and employment.