



April 2010

SP290-M Maple Petiole Borer

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: http://trace.tennessee.edu/utk_agexfores



Part of the [Entomology Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

"SP290-M Maple Petiole Borer," The University of Tennessee Agricultural Extension Service, SP290M 06-0263,
http://trace.tennessee.edu/utk_agexfores/92

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Insects, Pests, Diseases & Weeds is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Forestry, Trees, and Timber by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Insects

Maple Petiole Borer

*Frank A. Hale, Professor
originally developed by Jaime Yanes, Jr., former Assistant Professor
and Harry E. Williams, Professor Emeritus
Entomology and Plant Pathology*

The maple petiole borer, *Caulocampus acericaulis* (MacGillivray), was first introduced into the United States from Europe. This sawfly is wasp-like in appearance and is rarely seen because of its small size. Although several species of maple are subject to attack by this insect, sugar maples are preferred.

Life Cycle


Adult sawflies emerge in early to mid-April. After mating, the female uses her ovipositor (egg-laying structure) to puncture the petioles and lay a single

egg near the leaf blades. The larvae are yellowish and smooth with brownish heads. Larvae are weevil-like in appearance because the abdominal legs are very small. Full-grown larvae are one-third inch long. After hatching, larvae tunnel and consume the contents of the petioles about one-half inch from the leaf blade.

Early larval activity results in discoloration of the upper portion of the petiole. This area later becomes black and shrunk. The attached leaf shrivels and changes color. After a short period of time, the petioles break and damaged leaves fall to the ground (during late April to May). Larvae continue to feed in the portion of the petiole remaining attached to the twig for seven to 10 days. Larvae drop to the ground, burrow down 2 to 3 inches into the soil and overwinter. In the spring, mature larvae pupate and emerge as adults in early to mid-April. There is one generation a year.



*Petiole hollowed out in sugar maple larva
Charles D. Pless, University of Tennessee, www.forestryimages.org*



Control

Defoliation seldom exceeds a third of the total foliage. Damage of this type causes considerable concern to the homeowner. However, the leaf drop seldom causes injury to the tree and control recommendations are usually not warranted.

Some of the maple petiole borer damage can be prevented by applying Scimitar, Talstar, Tempo, Decathlon, Bayer Advanced Rose & Flower Insect Killer, or Tempo SC Ultra sprays to the foliage in mid-April.



Leaves severed from petioles in sugar maple
Charles D. Pless, University of Tennessee, www.forestryimages.org

Precautionary Statement

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label. Persons who do not obey the law will be subject to penalties.

Disclaimer Statement

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.

SP290M 06-0263

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.
UT Extension provides equal opportunities in programs and employment.