

University of Tennessee, Knoxville Trace: Tennessee Research and Creative **Exchange**

Field & Commercial Crops

UT Extension Publications

4-21-2010

W125 Sicklepod

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: http://trace.tennessee.edu/utk_agexcrop



Part of the Plant Sciences Commons

Recommended Citation

"W125 Sicklepod," The University of Tennessee Agricultural Extension Service, 06-00311, http://trace.tennessee.edu/utk agexcrop/ 113

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the UT Ag Research website. This Weeds and Herbicide Application is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Field & Commercial Crops by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Sicklepod

Larry Steckel, Assistant Professor, Plant Sciences

Sicklepod Senna obtusifolia (L.)

Also known as coffeebean, Java bean, Cassia obtusifolia (L.), Cassia tora (L.)

Classification and Description

Sicklepod is a member of the *Fabaceae* (Leguminoseae) or bean (pea) family. Sicklepod is an annual, herbaceous to semi-woody plant that is thought to be native to the American tropics. It can be found throughout the southeastern United States. The cotyledons of sicklepod are rounded and have three to five distinct veins. The stems are erect, branched, lack hairs (glabrous) and can reach heights of 1 - 6 feet. Sicklepod has alternate leaves comprised of 4 - 6 leaflets that are egg-shaped and arranged pinnately compound, which means that the leaflets are opposite one another. The pair of leaflets furthest from the main stem is the largest and the pair closest to the stem is the smallest. The egg-shaped leaflets can be 1 - 3.5 inches long. This weed reproduces by seed. The brownish seeds are housed in long (4 - 8 inches), sickle-shaped. hairless pods. Pods are a typical fruiting structure of a legume. Flowers have distinctive yellow petals and one or two arise from leaf axils (where stem meets leaf). Sicklepod has a stout taproot. This weed can be easily confused with coffee senna (Cassia occidentalis). However, the leaves of sicklepod are blunt while coffee senna's are pointed.

Weed Status and Injury

Sicklepod can be found in Tennessee in agronomic crops, waste places, moist forests and barnyards. Sicklepod can be a weed pest in cotton, corn, and soybeans. Sicklepod has been found to reduce cotton yield by 2.79 percent per weed per 30 row-feet. The critical time of control is the first four weeks after planting in most crops. Sicklepod can survive in a shaded environment but loses some competitive advantage once a crop has shaded the soil. It is one of the world's worst weeds.

Interesting Facts

Though sicklepod is a legume, it does not fix nitrogen. *Senna* is from ancient Greek and means an aromatic plant. This is in reference to sicklepod being rank-smelling when crushed. The Latin *obtusifolia* refers to the shape of the leaflets (*obtuse* blunt and *folia* leaves or blunt



Newly emerged sicklepod



First true leaf

leaves). Sicklepod seeds have had medicinal properties since 4000 B.C. when they were used as laxatives. Sicklepod hosts several insects and diseases of agricultural importance, including Asian soybean rust.

Management Considerations

Sicklepod is a very common weed in Tennessee row crop agriculture. It can be very competitive to cotton, soybeans and corn. Fortunately, a number of herbicides that can be sprayed on various crops provide good control of this weed. Please refer to the *Weed Control Manual for Tennessee Row Crops* (UT Extension PB 1580) for specific herbicides and management recommendations within various crops.

References

Gleason, H. A. and A. Cronquist. 1963. Manual of Vascular Plants. Willard Grant Press. Boston, MA, USA. 810 p.

Holm, L., J. Doll, E. Holm, J. Pancho and J. Herberger. 1997. World Weeds: Natural Histories and Distribution. John Wiley and Sons, Inc. New York, NY, USA. 1129 p.

Coker, T., K. Hurst, T. Kirk, J. Rupe, C. Tingle and M. Trent. FSA-7531 Asian soybean rust. Accessed May 21, 2006 http:// www.uaex.edu/Other_Areas/publications/ PDF/FSA-7531.pdf

Photo credits: L Steckel



Sicklepod competing with cotton



Sicklepod in cotton