



3-8-2018

Complete Genome Sequences of Three Staphylococcus pseudintermedius Strains Isolated from Botswana

Mohamed Adel Salaheldin Abouelkhair
University of Tennessee, Knoxville, mabouelk@vols.utk.edu


Stephen A. Kania
University of Tennessee, Knoxville

David A. Bemis
University of Tennessee, Knoxville

Matthew C. Riley
University of Tennessee, Knoxville

Riley Thompson
University of Tennessee, Knoxville

Follow this and additional works at: http://trace.tennessee.edu/utk_compmedpubs

 Part of the [Medical Genetics Commons](#), [Medical Immunology Commons](#), [Medical Microbiology Commons](#), [Medical Molecular Biology Commons](#), [Veterinary Infectious Diseases Commons](#), and the [Veterinary Microbiology and Immunobiology Commons](#)

Recommended Citation

Abouelkhair MA, Thompson R, Riley MC, Bemis DA, Kania SA. 2018. Complete genome sequences of three Staphylococcus pseudintermedius strains isolated from Botswana. *Genome Announcements*, March 2018 vol. 6 no. 10:e01599-17. <https://doi.org/10.1128/genomeA.01599-17>.

This Article is brought to you for free and open access by the Veterinary Medicine -- Faculty Publications and Other Works at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Faculty Publications and Other Works -- Biomedical and Diagnostic Sciences by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.



Complete Genome Sequences of Three *Staphylococcus pseudintermedius* Strains Isolated from Botswana

Mohamed A. Abouelkhair,^a Riley Thompson,^a Matthew C. Riley,^b David A. Bemis,^a  Stephen A. Kania^a

^aDepartment of Biomedical and Diagnostic Sciences, University of Tennessee College of Veterinary Medicine, Knoxville, Tennessee, USA

^bCenter for Genome Sciences, United States Army Medical Research Institute of Infectious Diseases, Fort Detrick, Maryland, USA

ABSTRACT We report here the first whole-genome sequences for 3 strains of *Staphylococcus pseudintermedius* (112N, 113N, and 114N) isolated in Africa. Samples of this opportunistic pathogen were collected from nasal swabs obtained from healthy carrier dogs in Botswana. The sequence information will facilitate spatial phylogenetic comparisons of staphylococcal species and other bacteria at the genome level.

Staphylococcus pseudintermedius is a Gram-positive, coagulase-positive bacterium commonly found as part of the normal skin and nasal flora of healthy dogs (1, 2). It is frequently associated with skin, soft tissue, and wound infections and less often with pneumonia and urinary tract infections similar to those caused by *Staphylococcus aureus* (2, 3). *S. pseudintermedius* infrequently colonizes the nasal mucosa of humans, where it has been associated with infections primarily from bite wounds or in immunocompromised individuals (3).

Most phenotypic and genotypic studies have been performed with methicillin-resistant *S. pseudintermedius* strains (1, 4–6). However, variability of phenotypic and genotypic features, including virulence, molecular epidemiology, and biological characteristics, have not been fully explored in methicillin-susceptible *S. pseudintermedius* (MSSP) isolates (7).

Whole-genome shotgun projects for *S. pseudintermedius* strains have been deposited at DDBJ/ENA/GenBank, including European *S. pseudintermedius* isolates (4, 6) and genomes from dominant clonal lineages in North America (6), but no complete genomes for *S. pseudintermedius* strains from the African continent are available and distinct geographical strain differences have been observed in this species (1, 4, 6, 8–13).

Whole-genome sequencing of MSSP isolates from Botswana will facilitate identification of genetic relatedness of these isolates with previously published MRSP and MSSP strains (1, 4–6, 12, 13) and facilitate evolutionary studies of staphylococcal species and other bacteria at the genome level.

All Botswana *S. pseudintermedius* strains were sequenced using Illumina MiSeq (Illumina, Inc., USA), with two runs (75 bp forward and reverse) generating a total of 1,091,288, 1,219,855, and 1,171,515 reads for *S. pseudintermedius* 112N, 113N, and 114N strains, respectively, with overall coverage of >250-fold. The resulting reads for each sequence were assembled using Geneious version 11.0.4 (14). The NCBI Prokaryote Genome Annotation Pipeline software version 4.3 was used for annotating the genomes.

The 112N genome is 2,516,677 bp (291 contigs of 500 bp or greater) with a 37.7% GC content, 2,375 predicted coding sequences, and 67 predicted RNAs. The 113N genome is 2,490,161 bp (96 contigs) with a 37.6% GC content, 2,257 predicted coding

Received 3 January 2018 Accepted 8 February 2018 Published 8 March 2018

Citation Abouelkhair MA, Thompson R, Riley MC, Bemis DA, Kania SA. 2018. Complete genome sequences of three *Staphylococcus pseudintermedius* strains isolated from Botswana. *Genome Announc* 6:e01599-17. <https://doi.org/10.1128/genomeA.01599-17>.

Copyright © 2018 Abouelkhair et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stephen A. Kania, skania@utk.edu.

sequences, and 72 predicted RNAs. The 114N genome is 2,546,835 bp (84 contigs) with a 37.5% GC content, 2,301 predicted coding sequences, and 72 predicted RNAs. The strains have unique multilocus sequence types (ST) (12) including ST887, ST888, and ST889 for *S. pseudintermedius* 112N, 113N, and 114N, respectively.

Accession number(s). These whole-genome sequences of *Staphylococcus pseudintermedius* 112N, 113N, and 114N strains have been deposited at DDBJ/ENA/GenBank under the accession numbers [PJUS00000000](https://doi.org/10.1128/genomeA.01651-16), [PJUR00000000](https://doi.org/10.1128/genomeA.01651-16), and [PJUQ00000000](https://doi.org/10.1128/genomeA.01651-16), respectively. The *Staphylococcus pseudintermedius* 112N, 113N, and 114N versions described in this paper are versions PJUS02000000, PJUR01000000, and PJUQ01000000, respectively.

ACKNOWLEDGMENTS

We thank the University of Tennessee Institute of Agriculture Center of Excellence in Livestock Diseases and Human Health for funding and support.

We acknowledge the University of Tennessee College of Veterinary Medicine Bacteriology Lab for isolating strains and for technical assistance.

REFERENCES

1. Abouelkhair MA, Riley MC, Bemis DA, Kania SA. 2017. Complete genome sequence of *Staphylococcus pseudintermedius* type strain LMG 22219. *Genome Announc* 5:e01651-16. <https://doi.org/10.1128/genomeA.01651-16>.
2. Devriese LA, Vancanneyt M, Baele M, Vanechoutte M, De Graef E, Snauwaert C, Cleenwerck I, Dawyndt P, Swings J, Decostere A, Haesebrouck F. 2005. *Staphylococcus pseudintermedius* sp. nov., a coagulase-positive species from animals. *Int J Syst Evol Microbiol* 55:1569–1573. <https://doi.org/10.1099/ijs.0.63413-0>.
3. Börjesson S, Gómez-Sanz E, Ekström K, Torres C, Grönlund U. 2015. *Staphylococcus pseudintermedius* can be misdiagnosed as *Staphylococcus aureus* in humans with dog bite wounds. *Eur J Clin Microbiol Infect Dis* 34: 839–844. <https://doi.org/10.1007/s10096-014-2300-y>.
4. Moodley A, Riley MC, Kania SA, Guardabassi L. 2013. Genome sequence of *Staphylococcus pseudintermedius* strain E140, an ST71 European-associated methicillin-resistant isolate. *Genome Announc* 1:e00207-12. <https://doi.org/10.1128/genomeA.00207-12>.
5. Videla R, Solyman SM, Brahmabhatt A, Sadeghi L, Bemis DA, Kania SA. 2018. Clonal complexes and antimicrobial susceptibility profiles of *Staphylococcus pseudintermedius* isolates from dogs in the United States. *Microb Drug Resist* 24:83–88. <https://doi.org/10.1089/mdr.2016.0250>.
6. Riley MC, Perreten V, Bemis DA, Kania SA. 2016. Complete genome sequences of three important methicillin-resistant clinical isolates of *Staphylococcus pseudintermedius*. *Genome Announc* 4:e01194-16. <https://doi.org/10.1128/genomeA.01194-16>.
7. Melter O, Svec P, Tkadlec J, Doskar J, Kinska H, Pantucek R. 2017. Characterisation of methicillin-susceptible *Staphylococcus pseudintermedius* isolates from canine infections and determination of virulence factors using multiplex PCR. *Veterinárni Medicína* 62:81–89. <https://doi.org/10.17221/105/2016-VETMED>.
8. Han JI, Rhim H, Yang CH, Park HM. 2018. Molecular characteristics of new clonal complexes of *Staphylococcus pseudintermedius* from clinically normal dogs. *Vet Q* 38:14–20. <https://doi.org/10.1080/01652176.2017.1400710>.
9. Pires dos Santos T, Damborg P, Moodley A, Guardabassi L. 2016. Systematic review on global epidemiology of methicillin-resistant *Staphylococcus pseudintermedius*: inference of population structure from multilocus sequence typing data. *Front Microbiol* 7:1599. <https://doi.org/10.3389/fmicb.2016.01599>.
10. Worthing KA, Abraham S, Coombs GW, Pang S, Saputra S, Jordan D, Trott DJ, Norris JM. 2018. Clonal diversity and geographic distribution of methicillin-resistant *Staphylococcus pseudintermedius* from Australian animals: discovery of novel sequence types. *Vet Microbiol* 213:58–65. <https://doi.org/10.1016/j.vetmic.2017.11.018>.
11. Ruscher C, Lübke-Becker A, Semmler T, Wleklinski C-G, Paasch A, Šoba A, Stamm I, Kopp P, Wieler LH, Walther B. 2010. Widespread rapid emergence of a distinct methicillin- and multidrug-resistant *Staphylococcus pseudintermedius* (MRSP) genetic lineage in Europe. *Vet Microbiol* 144: 340–346. <https://doi.org/10.1016/j.vetmic.2010.01.008>.
12. Kjellman EE, Slettemeås JS, Small H, Sunde M. 2015. Methicillin-resistant *Staphylococcus pseudintermedius* (MRSP) from healthy dogs in Norway—occurrence, genotypes and comparison to clinical MRSP. *MicrobiologyOpen* 4:857–866. <https://doi.org/10.1002/mbo3.258>.
13. Solyman SM, Black CC, Duim B, Perreten V, van Duijkeren E, Wagenaar JA, Eberlein LC, Sadeghi LN, Videla R, Bemis DA, Kania SA. 2013. Multilocus sequence typing for characterization of *Staphylococcus pseudintermedius*. *J Clin Microbiol* 51:306–310. <https://doi.org/10.1128/JCM.02421-12>.
14. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.