

Keywords:

Drought tolerance, Cold acclimation, Membrane remodeling, Leaf senescence, Reactive oxygen species (ROS), UDP-glycosyltransferase activity, Specialized metabolism, Flavonoid glycosides, Absciscic acid (ABA) metabolism, Fatty acid elongation, Cell wall biosynthesis, Transporter activity, Signal transduction, Local adaptation

Shared Mechanistic Insights: Core Climate Adaptation Mechanisms

1. Stress Response Integration: Drought and Cold

- **ERD7 (EARLY RESPONSE TO DEHYDRATION 7)** is a critical hub gene with roles in both **dehydration (drought)** and **cold stress responses** [Barajas-Lopez et al., 2021 (PMID: 33165601); Wu et al., 2024 (PMID: 38279327)].
 - **Cold Acclimation:** ERD7 mutants show increased susceptibility to cold, mediated via altered membrane lipid composition and reduced membrane fluidity. This results in impaired adaptation to low temperature.
 - **Dehydration and ABA Response:** ERD7 is responsive to water deprivation and abscisic acid (ABA), a key drought hormone. ABA-modulated genes are directly relevant to periodic drought in the Pacific Northwest.
 - **Molecular Mechanism:** ERD7 binds selectively to phospholipids and impacts phosphoinositide (PIP2) levels during cold, suggesting it is crucial for membrane remodeling under stress.

2. Regulation of Leaf Senescence and ROS Homeostasis

- **Senescence as an Adaptive Response:** ERD7 acts as a positive regulator of both age-dependent and stress-induced leaf senescence, with **phosphorylation states (Thr-225 and Ser-262)** modulating its activity [Wu et al., 2024 (PMID: 38279327)].
- **ROS Management:** ERD7 regulates accumulation of hydrogen peroxide via upregulation of **AtrbohD and AtrbohF** (NADPH oxidases).
 - **Functional Relevance:** Controlled leaf senescence and ROS detoxification provide a mechanism for plastic resource allocation and cellular protection during climate stresses.

3. Secondary Metabolism and Glycosylation

- **UDP-glycosyltransferase Activity (UGTs)**

- **GO Enrichment:** Significant enrichment for UDP-glycosyltransferase activity (GO:MF; p=0.0075) and UDP-glucosyltransferase activity (GO:MF; p=0.0306), indicating a core role in glycosylation pathways [GO/InterPro: IPR004224; PMID:11846783, 19858195].
- **Functional Outcomes:** UGTs catalyze glycosylation of flavonoids (e.g., kaempferol, quercetin, rutin) and hormonal metabolites (e.g., ABA glucosylation), enhancing metabolite stability, transport, and stress signaling.
- **Pathway Evidence:** Multiple genes participate in **kaempferol/quercetin/rutin glycoside biosynthesis** (KEGG), providing antioxidant capacity, UV protection, and membrane stabilization under stress.

4. Specialized Metabolism and Natural Variation

- **Functional Diversification:** Specific UGTs (e.g., UGT89A2) show natural variation in sugar donor specificity, affecting accumulation patterns of glycosylated metabolites across Arabidopsis accessions [Chen & Li, 2017 (PMID: 27411741)].
 - **Implication:** Polymorphisms influencing glycosylation patterns may underlie fine-scale local adaptation to diverse climates, paralleling Populus adaptation across the Pacific Northwest gradient.

5. Cell Wall & Lipid Remodeling

- **Cellulose and VLCFA Biosynthesis:** Genes in this set also map to **cellulose biosynthesis** and **very long chain fatty acid (VLCFA) biosynthesis** pathways (KEGG), essential for fortifying cell walls and modulating lipid composition under environmental stress.
 - **Relevance:** Structural cell wall and membrane adaptations buffer cells against desiccation, freezing, and mechanical stress, vital for perennial trees like Populus.

6. Transporter and Signal Functions

- **Monosaccharide and Nitrate Transporters** (AT5G61520, WR3): Support metabolic flexibility and nutrient uptake during stress adaptation.
- **Kinases and Signal Transduction** (e.g., AT1G66920, RLP9): Enable rapid signal relay during abiotic challenges.

7. Link to Populus trichocarpa Local Adaptation

- **Conserved Toolkits:** The involvement of UGTs, stress-response regulators (like ERD7), and membrane/cell wall remodeling genes in Arabidopsis underscores the likelihood that Populus uses similar mechanisms—membrane lipid modification, metabolic reconfiguration, regulated senescence, and specialized metabolites—to adapt to gradients of drought, cold, and other climate variables in the Pacific Northwest. This is supported by the original GEA study targeting

genetic associations with climate variables derived from 50-year regional TerraClimate records.

Unique or Distinctive Adaptive Mechanisms

Below are mechanisms identified in only one analysis—additional insights or points for future *Populus* research:

A. Natural UGT Variation and Sugar Donor Specificity

- **Summary:** Chen & Li (2017) identified key amino acids controlling donor specificity in UGT89A2, affecting dihydroxybenzoic acid (DHBA) glycoside accumulation patterns among *Arabidopsis* accessions (PMID: 27411741).
- **Relevance:** Polymorphic variation in specialized metabolism could allow tree populations to fine-tune stress metabolite profiles to heterogeneous microclimates of the Pacific Northwest.
- **Experimental Suggestion:** In *Populus*, GWAS or GEA approaches coupled with metabolic profiling could test for similar correlations between UGT variation and adaptive metabolite profiles.

B. Specific Role of Fatty Acid Elongation (KCS11)

- **Pathway:** Very long chain fatty acid (VLCFA) biosynthesis is only linked to a single gene (AT2G26640/KCS11) in this set, with known functions in cuticular wax and membrane lipid synthesis.
- **Relevance:** Enhanced cuticular waxes can improve water retention and freezing tolerance—key for trees exposed to both summer drought and winter cold.
- **Testing in *Populus*:** Manipulating VLCFA pathway genes (CRISPR or overexpression) and measuring cuticle properties, drought/cold tolerance phenotypes.

C. Nutrient/Resource Allocation (WR3)

- **Summary:** WR3 is annotated with nitrate transport and wound response—potentially unique aspects of nutrient and stress connectivity not widely enriched in other analyses.
 - **Relevance:** Variation in nitrate uptake could influence tree vigor and stress recovery in nutrient-variable sites.
 - ***Populus* Applicability:** Introduce labeled nitrate uptake assays in conjunction with climate variables to test genotype × environment interaction effects.
-

Citations and Data Sources

- **GO/KEGG/Gene Enrichment:** GO:MF enrichments, KEGG pathway mapping (as above). [IPR004224, PMID:11846783, 19858195]
 - **Experimental Literature:**
 - Barajas-Lopez JD et al., 2021. "EARLY RESPONSE TO DEHYDRATION 7 Remodels Cell Membrane Lipid Composition during Cold Stress in Arabidopsis." *Plant Cell Physiol.* 62:242-256. (PMID: 33165601, DOI:10.1093/pcp/pcaa139)
 - Wu R et al., 2024. "Phosphorylation of Thr-225 and Ser-262 on ERD7 Promotes Age-Dependent and Stress-Induced Leaf Senescence..." *IJMS* 25:21328. (PMID: 38279327, DOI:10.3390/ijms25021328)
 - Chen HY & Li X., 2017. "Identification of a residue responsible for UDP-sugar donor selectivity of a dihydroxybenzoic acid glycosyltransferase..." *Plant J.* 87: 634-646. (PMID: 27411741, DOI:10.1111/tpj.13271)
 - **Original Populus GEA Study Context:** 14 climate variables, 50 years of TerraClimate data, Pacific Northwest population.
-

Conclusion

The integrated analysis reveals a tightly coordinated set of stress response, metabolic, and structural genes and pathways—conserved between *Arabidopsis* and *Populus*—that underpin adaptation to climate variability. These include membrane and cell wall modification, ROS and senescence regulation, and broad modulation of specialized metabolites via glycosylation—all likely critical for local adaptation in forest trees across environmental gradients. Unique mechanisms such as natural UGT variation or nitrate transport add additional layers for experimental validation in *Populus*, advancing our understanding of forest tree climate adaptation.