Efficacy of Extending the Voluntary Waiting Period in Lactating Dairy Cows to Improve Fertility of Sex-Sorted Semen

A Thesis Presented for the

Master of Science

Degree

The University of Tennessee, Knoxville

Sarah E. Orr

December 2015

DEDICATION

I would like to dedicate this thesis to my family and friends. I could not have done this without your love, support, and encouragement!

ACKNOWLEDGMENTS

I have been blessed to have received assistance and guidance from many people throughout my graduate education, and though I am going to attempt to acknowledge everyone, I am positive I will miss someone along the way. To those that I forget to mention, thank you!

First and foremost, thank you to Dr. Edwards and Dr. Schrick for agreeing to be co-mentors and take me on as a graduate student. Both of you have been great about pushing me out of my comfort zone, challenging me to do and be more than I ever thought possible, and helping me to grow and develop into a better person. Thanks to Dr. Rhinehart for agreeing to be on my Master's committee. I am grateful to each of you for the opportunities and experiences I have been provided as a result of working with each of you. Also, thanks to Dr. Saxton for teaching me how to analyze my data and guiding me through the process of statistical analyses.

A huge thank you must be extended to Charlie Young at the Little River Dairy for the many ways he helped me out. Whether it was teaching the beef cattle girl all about real-life dairy production, guiding me through the intimidating time of data and statistical analyses, teaching me how to tail bleed a dairy cow, or believing in me enough to let me perform AI and embryo transfer on your cows, thank you for answering all my questions and always being patient with me. I cannot begin to express how grateful I am to have had your day-to-day guidance out at the barn or how thankful I am that you were always willing to help me out (usually by breeding a cow at 6 a.m.) whenever my schedule was particularly crazy. Somehow all the cookies in the world do not seem like thanks enough. I must also mention my appreciation for the rest of the ETREC-Little River staff—Mark, Tate, Taylor, Shawn, Jacob, and Kody. Thank you all for all your assistance with my project, always saying hello, and bringing a smile to my face to make my time spent out there very pleasant.

A simple thank you is not nearly enough to show my gratitude for Leah Hooper and Megan Goodwin. These girls were there to show me the ropes about so many things when it came to being a graduate student with Dr. Edwards, whether it was how to be a teaching assistant, knowing what Schrick and Edwards actually wanted versus what they said they wanted, or how to develop a love for Starbucks. You all were both so helpful when I needed assistance with my project and for that I am very thankful! Megan, from the first day of graduate school, you took me under your wing and taught me all there was to know about how to handle this somewhat intimidating time in my life. To this day, you still periodically check in on me to provide encouraging words and motivation to finish, and somehow they always come at just the right time. Leah, bless your heart for sharing an office with me; I am pretty sure you still have not developed an appreciation for country music or boots and probably never will—but at least I can say I tried! All jokes aside, you were a fabulous office mate, and I really appreciate you taking time to teach me so many important things like how to write announcements for class or what all the names of the ninja turtles were. Furthermore, I must express my thanks to Laci Gunthorpe for her assistance in her limited time here. Thank you for being a walking-talking reminder of my to-do list, helping me remember everything I was supposed to do, listening to me vent from time-to-time, and helping me to let loose at the end of a long week.

The undergraduates that volunteered to help with my study cannot go unmentioned because without them, heat checking would have been much more difficult some days and not nearly as fun. Good company while watching cows “stand around chewing cud” at 6 a.m. made for a better start to the day. The work ethic and dedication of Tyler Trowbridge, Emma Bishop, Bryce BeVier, Hannah McDonald and Clare Welsch is evident because they willingly came early, stayed late, and did anything I asked them to help with. I cannot count the number of

times I sent Tyler to heat check or run blood while I handled class and other teaching assistant duties, or how many times I had to rely on Emma to process blood on late evenings because a cow came in and changed all our plans; I never doubted you all and always trusted you would handle things the right way. Bryce—thank you for always making heat checking an interesting experience and for challenging me to be patient and appreciate different points of view. Hannah, I honestly do not have a clue what I would have done without your help with data this past summer. I am certain you were completely tired of looking at excel spreadsheets, as was I, but please know that every little thing you did was a huge help to me. Thanks also for bringing new and interesting experiences into my life—like shark's teeth and shark eggs. Clare, even though your time out at the barn was limited, I thoroughly appreciate your dedication, willingness to help with anything—my research, class responsibilities, and everything else, and especially your moral support when I just need to talk. I could not have done this without each of you, and I wish you all nothing but the best in your future endeavors!

Kaitlyn Hess and Morgan Bragg were fantastic undergraduate lab workers and helped out with many class and lab related things. Thank you for getting up early to meet me at JARTU at 6 a.m. for lab preparation, staying late for clean up, and always working hard at anything I asked you to help with. Your hard work and dedication did not go unnoticed!

Drs. Rebecca Payton and Louisa Rispoli were instrumental in my graduate education in so many ways. Much thanks to each of you for all your help along the way! Becca, thank you for being my go-to person for any kind of question I had—SAS related for when I could not figure out the many errors I got or the right code to fix it, class and lab related for when I came to you with "Edwards said she wanted (insert request here), can you help me with that?", or the plethora of other things I asked about—my favorite of which is definitely "what in the devil is

poutine?”. Thank you for always lending a hand when I needed help, encouraging me through the tough times and the tears, and wiping the cow manure off my face at the barn! Louisa, I do not think I, my thesis, or any other computer documents would have survived my graduate school experience without you. Heaven knows you tried to save my computer, but it turns out I am pretty good at taking them out since I am on my third one now! Thanks for all your technical support as well as the mental and emotional support. I am grateful you included me in some of your projects, especially the ones that got me to the farm! Thanks to both of you for helping, supporting, and guiding the non-lab graduate student on her journey.

Last, but certainly not least, I owe the biggest debt of gratitude to my family and friends for getting me through this experience. Momma, Daddy, Heather, Emily and the rest of my family and friends too numerous to list here, have been phenomenal over the past two and a half years. Your understanding and support as I furthered my education have been unbelievable and definitely way more than I deserved. I know that at times I have not been the most pleasant person to be around, but you all never stopped loving me, listening to me, letting me cry on your shoulder or vent over the phone, encouraging me, and believing in me even when I did not believe in myself. Thanks for being strong for me when I was not strong enough on my own. I am so lucky and blessed to have each of you in my life.

ABSTRACT

The objective of this study was to compare fertility of sex-sorted semen in lactating cows using a voluntary waiting period (**VWP**) of approximately 55 days (**VWP55**) or 85 days (**VWP85**). At 21 days in milk (**DIM**), cows were randomly assigned to begin estrus synchronization at 55 or 85 DIM. Only cows confirmed cyclic by 55 DIM were synchronized. At 55 or 85 DIM, cows [VWP55 (n=44); VWP85 (n=45)] having a corpus luteum (**CL**) were administered $PGF_{2\alpha}$ [two alpha]. Estrus was monitored continuously by the HeatWatch® [registered sign] system (**HW**) and by twice daily visual observation seven days following each $PGF_{2\alpha}$. Cows exhibiting estrus were artificially inseminated with sex-sorted Holstein semen using one of two different bulls having similar sire conception rates. After 14 days, $PGF_{2\alpha}$ was administered to any cow that did not exhibit estrus. Cows not responding to either $PGF_{2\alpha}$ administration were administered GnRH and a controlled intravaginal drug releasing device (**CIDR**). Seven days later, CIDR was removed and $PGF_{2\alpha}$ was administered. Pregnancy determination occurred 28 to 35 days after AI and was monitored at 60, 80, and 150 days. No difference in overall percentage of cows expressing estrus (79.55% and 77.78%, P = 0.8393), hours to estrus following either the first or all $PGF_{2\alpha}$ administrations (65.79 h and 72.65 h, P = 0.4908; 62.28 h and 73.00 h, P = 0.2218, respectively), number of mounts (9.05 and 10.54, P = 0.5246), average mount duration (4.59 s and 4.99 s, P = 0.2602), and total estrus length (8.16 h and 9.17 h, P = 0.6389) was observed between VWP55 and VWP85 cows. Conception rates 30 days after AI were significantly higher for VWP85 cows compared to VWP55 (51.43% vs. 22.86%, respectively; P = 0.0181). Conception rates were monitored at 60 (48.57% vs. 20.00%, respectively; P = 0.0166), 80 (42.86% vs. 20.00%, respectively; P = 0.0471), and 150 days after AI (42.86% vs. 20.00%, respectively; P = 0.0471) to monitor embryonic loss. Fertility of sex-

sorted semen in lactating cows can be improved after a VWP of 85 days under specific management conditions.

TABLE OF CONTENTS

CHAPTER 1 INTRODUCTION 1
CHAPTER 2 REVIEW OF LITERATURE 4
Introduction 4
Increasing World Population and the Need for Improved Dairy Cow Fertility 4
Declining Fertility of Lactating Dairy Cows 5
Milk Production and Calving Related Challenges in Dairy Cows 5
Negative Energy Balance 7
Ovarian Cyclicity 8
Managing Fertility of the Lactating Dairy Cow 9
Voluntary Waiting Period 9
Increased Fertility Coincident with Increased Days in Milk 10
Increased Fertility after Estrus Detection at Later Days in Milk 10
Increased Fertility after Timed Artificial Insemination at Later Days in Milk 11
Fertility of Sex-Sorted Semen 12
Dairy Heifers 13
Lactating Cows 14
Economic Aspects 16
CHAPTER 3 MATERIALS AND METHODS 19
Treatment Assignments 19
Estrus Synchronization and Detection 21
Artificial Insemination & Pregnancy Determination 22
Statistical Analyses 22
CHAPTER 4 RESULTS AND DISCUSSION 24
Estrus Activity 25
Artificial Insemination and Conception Rates 29
Conclusions 35
Implications 37
LITERATURE CITED 38
VITA 53

LIST OF TABLES

Table 4.1. Estrus characteristics for VWP55 and VWP85 cows…….26

Table 4.2. Influence of parity on milk production……...33

Table 4.3. Influence of parity on reproductive tract scores……..34

Table 4.4. Influence of parity and milk production on resumption of ovarian cyclicity…….36

LIST OF FIGURES

Figure 3.1. Estrus synchronization protocol for VWP55 and VWP85 cows.....20

Figure 4.2. Percentage of estrus expression over time following $PGF_2\alpha$ administration……...28

Figure 4.3. Conception rates for VWP55 and VWP85 cows…….30

CHAPTER 1

INTRODUCTION

Sex-sorted semen is a valuable tool for the dairy industry to increase the number of heifer calves available. Fertility of sex-sorted semen is lower than conventional semen. For instance, in dairy heifers, fertility of sex-sorted semen is on average 30% lower (range: +14% to -77%) than conventional semen (DeJarnette et al., 2008; Seidel and Schenk, 2008; DeJarnette et al., 2009; Schenk et al., 2009; DeJarnette et al., 2010; Norman et al., 2010; DeJarnette et al., 2011; Healy et al., 2013). In lactating dairy cows, fertility of sex-sorted semen is reduced by 26% (range: 0% to -54%) compared to conventional semen (Andersson et al., 2006; DeJarnette et al., 2008; Schenk et al., 2009; DeJarnette et al., 2010; Norman et al., 2010; Karakaya et al., 2014). This is compounded by conventional semen conception rates that are only approximately 32% in lactating dairy cows (Norman et al., 2009; Schefers et al., 2010; Wiltbank and Pursley, 2014), resulting in many recommendations that sex-sorted semen usage be limited to dairy heifers (DeJarnette et al., 2009; Norman et al., 2010; Healy et al., 2013). Multiple studies have evaluated methods to address the fertility deficit associated with sex-sorted semen including altered uterine deposition sites (Seidel and Schenk, 2008), sperm cell concentrations (DeJarnette et al., 2008), sorting and storage processes (Schenk et al., 2009), timing of insemination (Sales et al., 2011), and integration with timed AI (Sá Filho et al., 2013), yielding only marginal, if any, increases in conception rates for lactating cows.

A critical component of dairy breeding strategies that impacts fertility is the length of the voluntary waiting period (**VWP**), which is a defined time period after calving that producers elect to wait before efforts to inseminate those cows begin. Waiting a defined time period after calving is critical to allow a cow adequate time to go through uterine involution, recover from

any postpartum health issues, and resume ovarian cyclicity. Many studies have reported benefits of waiting to inseminate cows until later in lactation. Summarizing seven studies, Britt (1975) reported that fertility was 25% for cows inseminated in the first 20 days after calving compared to greater than 60% for cows inseminated after 70 days postpartum. One study demonstrated improved conception rates in cows bred for the first time after 50 days in milk (**DIM**) compared to those bred before 50 DIM (Hillers et al., 1984). Others found that cows inseminated ≥74 DIM had higher fertility than cows inseminated earlier in lactation (Whitmore et al., 1974). Tenhagen et al. (2003) indicated that Holstein cows submitted for timed insemination using conventional semen after an OvSynch protocol had increased first service conception rates as the stage of lactation progressed from 53 to 59 DIM, to 73 to 81 DIM, and 94 to 102 DIM. Stevenson and Phatak (2005) noted improvements in pregnancy rate as DIM increased in cows that were originally synchronized for timed insemination but were inseminated following standing estrus before timed insemination.

Despite obvious improvements of inseminating cows with conventional semen later in lactation (Whitmore et al., 1974; Britt, 1975; Hillers et al., 1984; Weller and Folman, 1990; Tenhagen et al., 2003; Stevenson and Phatak, 2005; Caraviello et al., 2006; Schefers et al., 2010), two producer surveys of 103 and 673 US herds report current practices that employ an average VWP of 52 ± 1.3 and 56 ± 0.6 days, respectively (Caraviello et al., 2006; DeJarnette et al., 2007). Because fertility of conventional semen increases when cows are inseminated later in lactation, we hypothesized that extending the VWP beyond 55 DIM would be beneficial for improving fertility for lactating cows inseminated with sex-sorted semen. In support of this, Schenk et al. (2009) retrospectively found that lactating dairy cows inseminated between 84 to 98 DIM had increased pregnancy rates compared to inseminations earlier in lactation. Thus, the

objective of this study was to directly compare the fertility of sex-sorted semen in lactating cows using a VWP of approximately 55 days or a VWP of 85 days.

CHAPTER 2

REVIEW OF LITERATURE

Introduction

The following is a review of the pertinent scientific literature describing the factors contributing to the current fertility status of lactating dairy cows. Use of sex-sorted semen in both dairy heifers and lactating cows will be discussed as well as the impact of the voluntary waiting period on fertility. Finally, economic considerations for both sex-sorted semen and extended VWPs will be reviewed.

Increasing World Population and the Need for Improved Dairy Cow Fertility

In 2011, it was estimated that world population would reach 9.15 billion people by the year 2050 (FAO, 2011). With that estimate, production of dairy products needs to increase by more than 121% to accommodate the projected consumption (FAO, 2011). More current estimates by the United Nations suggest that world population will be even greater—9.7 billion people—by 2050, further attesting to the need for more dairy products to feed a growing world (United Nations, 2015).

To this end, US milk production has increased by 1.3% yearly in recent years and is likely to continue at this rate into the future (Santos et al., 2010). More milk produced per cow translates into fewer cows needed to produce an equivalent amount of milk. It has been suggested that 11.1% fewer cows will be needed to meet milk production requirements in 40 years (Santos et al., 2010). Increasing milk production per cow however does not come without a cost. Genetic selection for increased milk production over the past five decades is one of many factors that have been coincident with reduced fertility of lactating dairy cows (reviewed by Lucy, 2001). Because pregnancy and subsequent parturition are required for initiating lactation,

efforts to improve fertility of the dairy cow are important to efficiently meet the production needs of the future.

Declining Fertility of Lactating Dairy Cows

Dairy cow fertility worldwide has declined dramatically since the 1950s (reviewed by Lucy, 2001; Washburn et al., 2002). In 1998, after reviewing the literature, Butler determined that conception rates of lactating dairy cows had decreased from approximately 65% in 1951 to only 40% in 1996 for New York dairy herds (Butler, 1998). From 1975 to 1997, the decline was estimated to be at a rate of 0.5% per year (Beam and Butler, 1999). Increased services per conception and average conception rates of less than 40% have led to a decline in calving rate of approximately one percent per year (Royal et al., 2000b). Reduced fertility results in decreased genetic gain within the herd (Hohenboken, 1999; reviewed by Lucy, 2001; Seidel, 2007; McCullock et al., 2013) and fewer replacement females. These negative consequences result in decreased profitability for the producer (Royal et al., 2000b; McCullock et al., 2013).

Recent evidence from analysis of nearly 20 million inseminations from Dairy Herd Improvement breeding records obtained since 1995 demonstrated that first service conception rates for Holstein cows have begun to improve (Norman et al., 2009). According to Norman et al. (2009), the average first service conception rate for lactating Holsteins rose from 27% to 32% between 2001 and 2007 and has been attributed to the adoption of estrus synchronization or timed artificial insemination protocols. Although fertility in lactating cows is improving, it is still at suboptimal levels, which emphasizes the need for additional work toward improvement.

Milk Production and Calving Related Challenges in Dairy Cows

Numerous reports in the literature have associated decreased fertility with increased milk production. Rolling herd average milk production rose from approximately 6,500 kg per

lactation in the 1970s to nearly 9,000 kg per lactation in the late 1990s (reviewed by Lucy, 2001). Corresponding reductions in conception rates, increased calving intervals, and increased services per conception suggest an antagonistic relationship between milk production and fertility (reviewed by Lucy, 2001). In support of this notion, Dhaliwal et al. (1996) determined that high yielding cows required 0.42 more services and needed an additional 25 days to conceive when inseminations began before 80 DIM. Nevertheless, others have reported weak or no impact of milk yield on fertility (Darwash et al., 1997; Loeffler et al., 1999; Gröhn and Rajala-Schultz, 2000).

Many other factors may also negatively influence fertility. Dystocia increased the number of services required for conception, interval to first estrus, interval to first service, days open, and calving interval compared to cows having normal parturitions (Dematawena and Berger, 1997; Gaafar et al., 2011). Furthermore, cows experiencing severe cases of dystocia have reduced conception rates by 200 DIM compared to cows experiencing mild cases of dystocia or having a normal parturition (Tenhagen et al., 2007). Other calving-related issues such as twinning also affect fertility (Roche, 2006). After parturition, uterine involution must occur, whereby the uterus regresses in size and expels fetal membranes and associated tissue attachments. Uterine involution typically occurs by 30 to 40 DIM (Okano and Tomizuka, 1987). However, the uterus may not be capable of supporting a new pregnancy until 60 or more DIM (Roche, 2006). Bacterial contamination of the uterus after calving is a common occurrence. If the immune response is delayed, other postpartum related disorders such as retained placenta, metritis and endometritis may reduce fertility (Thatcher et al., 2006). Furthermore, subclinical and clinical mastitis early in lactation have been reported to negatively impact fertility by increasing services per conception and days open (Barker et al., 1998; Schrick et al., 2001).

Negative Energy Balance

The transition period from gestation to early lactation requires a high level of nutrition to meet increased energy demands for milk production in early lactation. Feed intake must increase to partially accommodate the partitioning of nutrients for lactation as well as maintenance requirements with the remaining energy needs being met by mobilization of fat tissue (Bauman and Currie, 1980; Wathes et al., 2007; Roche et al., 2009). Decreased feed intake and fat mobilization prepartum and early postpartum often cause high producing cows to experience an energy deficit, known as negative energy balance [**NEB;** (Bauman and Currie, 1980)]. Negative energy balance typically extends from approximately two weeks before calving to 8 to 12 weeks postpartum and can lead to negative effects lasting into the breeding period (Butler, 2003, 2005).

Negative energy balance has adverse effects on reproductive functions and fertility (Butler, 2003; Wathes et al., 2007). For instance, cows in NEB may have altered luteinizing hormone pulse frequency and reduced glucose, insulin, and insulin-like growth factor I levels which contributes to increased length of anovulation (Butler, 2005; Wathes et al., 2007). These alterations may lead to reduced estradiol synthesis which slows follicular growth and extends time to first postpartum ovulation (Butler, 2003; Wathes et al., 2007). Negative energy balance also causes the liver of the cow to metabolize steroid hormones at an increased rate, resulting in lower circulating levels of estradiol and progesterone which further reduces the likelihood of ovulation (Butler, 2003; Wiltbank et al., 2006). Furthermore, mobilization of body tissues for energy results in the production of non-esterified fatty acids and beta-hydroxybutyrate acid which negatively impact the follicular environment to impair oocyte quality and developmental competence (Wathes et al., 2007). Subsequent embryo development may be compromised because lower progesterone levels shortly after fertilization may fail to adequately prepare the

uterus to support pregnancy (Butler, 2003; Wathes et al., 2007). It is also suggested that NEB impedes uterine repair by hampering the clearance of pathogens from the uterine environment following calving (Wathes et al., 2007).

Negative energy balance can be indirectly monitored through the use of body condition scoring (**BCS**), a subjective method involving assessment of fat stores and assignment of a numerical score, generally using a scale of one (emaciated) to five (obese) (Ferguson et al., 1994). The BCS of the cow at the start of lactation and the amount of body condition lost during early lactation can be important predictors of subsequent fertility (Gillund et al., 2001; Roche et al., 2009). BCS loss of one or more numerical unit(s) after calving resulted in longer calving to conception intervals and required more artificial inseminations for cows to conceive (Gillund et al., 2001). Notably, BCS loss did not affect calving to first service interval, which helps to explain why more services were required for conception (Gillund et al., 2001). Similarly, in a large, U.S. based, commercial dairy herd, when BCS decreased by 0.4 or 0.8 points during the first month of lactation, cows were 1.17 and 1.36 times less likely to conceive after the first AI, respectively (Domecq et al., 1997).

Ovarian Cyclicity

One of the most important factors affecting cow fertility is cyclicity status. Research suggests that atypical cyclicity patterns such as prolonged luteal phase, follicular cysts, and cessation of cyclicity in lactating cows have increased over time (Royal et al., 2000a). Multiple studies reported that approximately 20 to 50% of cows remained anovular by 50 to 60 DIM (Moreira et al., 2001; El-Zarkouny et al., 2004; Santos et al., 2004; Shrestha et al., 2004). Shrestha et al. (2004) classified cows that took more than 45 DIM to return to cyclicity by causative issues including prolonged or shortened luteal phase, delayed first ovulation, or

cessation of cyclicity and found that majority of delayed cows had delayed first ovulation or prolonged luteal phase. Evidence from a Japanese study of primiparous and multiparous Holsteins demonstrated that time to first ovulation and first estrus was shorter for first lactation females compared to multiparous cows (Sakaguchi et al., 2004), while others found that first parity cows took longer to return to cyclic status compared to multiparous cows (Tanaka et al., 2008; Chebel and Santos, 2010). Though submitting cows to GnRH-based timed artificial insemination (**TAI**) may induce cyclicity in some cows, pregnancy rates are still reduced compared to cows that regained cyclicity without hormonal induction (Cerri et al., 2004; Santos et al., 2004; Chebel et al., 2010; Chebel and Santos, 2010). The percentage of cows that resumed ovarian cyclicity increased as DIM increased (Shrestha et al., 2004), and correspondingly others have reported that more estrus events before insemination results in increased fertility (Thatcher and Wilcox, 1973; Sakaguchi, 2010).

Managing Fertility of the Lactating Dairy Cow

Voluntary Waiting Period

The voluntary waiting period (**VWP**) is an essential management component in dairy cattle production and is a defined length of time after parturition during which cows are purposely not submitted for insemination (Chebel and Santos, 2010). The VWP allows time for the cow to undergo healing processes including recovering from any postparturient disorders, completing uterine involution, and returning to a cyclic status before to insemination (Stevenson, 2001; Miller et al., 2007; Wathes et al., 2007; Chebel and Santos, 2010). Two surveys of 103 and 673 herds found that the average length of the VWP in American dairy herds is approximately 52 and 56 days, respectively (Caraviello et al., 2006; DeJarnette et al., 2007). In some cases, the range of VWP lengths employed can vary from 40-70 days (Stevenson, 2001).

Though some may vary the length of the VWP for multiple reasons such as post-calving health, parity, and milk yield, waiting at least ~ 55 days before first insemination is imperative. (DeJarnette et al., 2007; Miller et al., 2007). Approximately 20-50% of cows may still be anovular by 50-60 DIM (Linderoth, 2005; Miller et al., 2007; Chebel and Santos, 2010), suggesting that breeding immediately after the traditional VWP of 50-60 DIM may result in fewer successful conceptions.

Increased Fertility Coincident with Increased Days in Milk

Fertility in lactating dairy cows may increase as DIM increase (Caraviello et al., 2006; Schefers et al., 2010) as evidenced by improved conception rates when cows were inseminated between 120 and 180 DIM (Niozas et al., 2015). However, majority of these conclusions are based on retrospective analysis of data, and few instances exist where controlled studies examined the effects of altering the VWP, primarily to address summer heat stress (Gobikrushanth et al., 2014) and economic impacts of extending or altering calving intervals (Arbel et al., 2001; Niozas et al., 2015).

Increased Fertility after Estrus Detection at Later Days in Milk

In a 1975 review, sevens studies were summarized and concluded that fertility increased from 25% when artificial inseminations after detected estrus occurred after 20 DIM compared to more than 60% after 70 DIM (Britt, 1975). In a study designed to determine if artificially inseminating cows earlier postpartum would impact future reproductive performance, conception rates to first inseminations after the first postpartum estrus (~39 DIM) were lower compared to inseminations after 74 DIM [67% vs. 37%, respectively; (Whitmore et al., 1974)]. Both Britt (1975) and Whitmore et al. (1974) reported a decrease in services per conception from approximately 2.2 to 1.5 when cows were inseminated at later DIM. Whitmore et al. (1974)

followed cows over three lactations and noted increased fertility of 43, 42, and 22 percentage points higher for cows inseminated at first estrus after 74 DIM for the first, second, and third lactation, respectively. Furthermore, Hillers et al. (1984) reported significantly lower conception rates when cows were inseminated before 50 DIM compared to those that were inseminated between 50-79 DIM (32% vs. 49-54%, respectively). Weller and Folman (1990) concluded that inseminations at 83 DIM achieved desirable fertility without compromising milk production later in lactation.

Increased Fertility after Timed Artificial Insemination at Later Days in Milk

The goal of timed AI (**TAI**) is to utilize different pharmaceuticals to synchronize ovulation to occur at a predicted time, allowing inseminations to be performed at a specific time (Wiltbank and Pursley, 2014). After each $PGF_{2\alpha}$ injection(s) in a given protocol, an opportunity exists for cows to exhibit estrus, and thus insemination may occur if estrus is detected. A common management practice among dairy operations utilizing TAI synchronization practices is to inseminate cows that exhibit estrus before TAI (Stevenson and Phatak, 2005). Holstein cows inseminated with conventional semen after detected estrus during two different TAI protocols had increased pregnancy rates as DIM increased from 54 to 77 DIM [22.6% vs 48.7%, respectively; (Stevenson and Phatak, 2005)]. Inseminations based on estrus during the TAI protocols compared to those occurring only during the $PGF_{2\alpha}$ pre-synchronization demonstrated pregnancy rates that were significantly increased at greater DIM [31.3% at 59 DIM vs. 45.3% at 74 DIM (Stevenson and Phatak, 2005)]. Chebel and Santos (2010) reported numerically higher conception rates for cows inseminated at 72 DIM compared to cows inseminated at 65 DIM (39.6% vs. 33.0%, respectively; $P = 0.14$; $n = 639$). However, it is important to note that cows in the shorter VWP were inseminated after estrus during a pre-synchronization of two $PGF_{2\alpha}$

injections, but if estrus was not expressed, short VWP cows were time inseminated along with the longer VWP cows which were only time inseminated following the OvSynch protocol (Chebel and Santos, 2010). Gobikrushanth et al. (2014) found that extending the VWP from an average of 60 DIM (range: 57 to 63 DIM) to an average of 83 DIM (range: 64 to 121 DIM) resulted in slight improvements (32.1% to 37.7%; P = 0.02) in conception rates; however, part of the study was conducted during times of heat stress and could influence observed conception rates.

Modern dairy operations often utilize TAI synchronization to inseminate cows because of the benefits of eliminating estrus detection and having defined time periods for inseminations. Unfortunately, TAI following a traditional VWP of 50 to 60 days often results in decreased conception rates compared to inseminations following estrus detection (Kasimanickam et al., 2005). Tenhagen et al. (2003) reported increased first service conception rates when TAI was performed after at least 73 to 81 DIM among Holstein cows segregated by production level. Using the Ovsynch protocol, cows inseminated at 53 to 59 DIM (low production) and 73 to 81 DIM (average production) had significantly lower conception rates compared to cows inseminated at 94 to 102 DIM [high production; 14.4% vs. 28.7% vs. 41.4%, respectively; (Tenhagen et al., 2003)]. It is noted that milk production level did not have an influence on conception rates when cows were synchronized at similar DIM, thus differences in conception rate could be a result of inseminating cows at differing DIM (Tenhagen et al., 2003).

Fertility of Sex-Sorted Semen

Fertility of a dairy cow is also influenced by the service sire. Currently, two broad categories of semen types exist commercially: conventional and sex-sorted semen. Sex-sorted semen is a valuable reproductive tool for the dairy industry to aid in herd expansion, improve the

rate of genetic gain, and increase the number of replacement females available (Hohenboken, 1999; Seidel, 2007; McCullock et al., 2013). A difference in fertility exists between conventional and sex-sorted semen. Moreover, differences also exist when comparing fertility of sex-sorted semen in heifers to that of lactating cows (Bodmer et al., 2005; Andersson et al., 2006; Schenk et al., 2009; DeJarnette et al., 2010; DeJarnette et al., 2011; Healy et al., 2013; Karakaya et al., 2014). In general, fertility of sex-sorted semen is 20% to 30% lower than conventional semen for both heifers and lactating cows (Andersson et al., 2006; DeJarnette et al., 2008; Seidel and Schenk, 2008; DeJarnette et al., 2009; Schenk et al., 2009; DeJarnette et al., 2010; Norman et al., 2010; DeJarnette et al., 2011; Healy et al., 2013; Karakaya et al., 2014). Lower conception rates for sex-sorted semen combined with increased cost prevents its widespread usage (Seidel, 2003).

Dairy Heifers

In four different field trials working with Holstein heifers in multiple locations and under different management conditions, Seidel et al. (1999) found average conception rates for frozen-thawed sex-sorted semen to be 50 to 56% compared to 67% for conventional semen. These rates were not improved by altering sperm concentration, deposition site, or freezing times (Seidel and Schenk, 2008). Schenk et al. (2009) reported sex-sorted semen conception rates that were equivalent to conventional conception rates (49%, 34%, 54% and 43%, 62%, 61%, respectively) following alterations in sorting pressure, storage time, laser sources, and media. Despite these findings, DeJarnette et al. (2011) still questioned the plausibility of ever achieving equivalent conception rates for sex-sorted and conventional semen. Previous efforts were inconclusive about altering semen dosage for improving conception rates for sex-sorted semen. DeJarnette et al. (2008) reported no influence of 2.1, 3.5, or 5.0 x 10^6 sperm per insemination

dose on heifer conception rates (46.7%, 51.2%, 52.5%, $P > 0.05$), but DeJarnette et al. (2010) found that differences exist between 2.1 and 3.5 x 10^6 compared to conventional semen (44% and 46% vs. 61%, respectively; $P < 0.05$). Furthermore, Healy et al. (2013) retrospectively analyzed heifer inseminations for sex-sorted and conventional semen found predicted conception rates of 21.3% and 32.1%, respectively, which were noticeably lower than other studies. According to Bodmer et al. (2005), sex-sorted semen pregnancy rates were significantly lower than conventional semen pregnancy rates (33.3% vs. 59.3%; $P = 0.05$). While reductions are noted in dairy heifers for sex-sorted semen, even after multiple approaches to achieve improvements, usage in heifers remains a realistic option for producers due to the inherent increased fertility of nulliparous females (Butler et al., 2014) which may help to justify the added expense to obtain more heifer calves.

Lactating Cows

Conception rates may be considerably lower for sex-sorted semen in lactating cows due to fewer sperm per insemination dose and compromised fertility of the sperm (Seidel and Garner, 2002), the added stress of milk production in cows (reviewed by Lucy, 2001), the timing of the insemination (Sa Filho et al., 2010), or a combination of these or other unknown factor(s). Since the national average conception rate in lactating dairy cows is only 32% (Wiltbank and Pursley, 2014), reductions in fertility of any magnitude negatively affect usage of sex-sorted semen. When conception rates for sex-sorted semen were compared across 211 herds, slightly more than half as many lactating cows conceived compared to dairy heifers [27%, n= 2,298 vs. 45%, n= 48,200; (Schenk et al., 2009)]. In cows selected for absence of uterine and ovarian abnormalities, sex-sorted semen (2 x 10^6 sperm per straw) pregnancy rates were similar to conventional semen [10 x 10^6 sperm per straw; (40.5 ± 6.2% vs. 55.6 ± 6.5%, respectively; n =

115)] when TAI was performed at ~80 DIM (Crichton et al., 2006; Schenk et al., 2009). However, when reproductive status, insemination method, and stage of lactation were not controlled for, pregnancy rates for sex-sorted semen (2 x 10^6 sperm per straw) were significantly lower than unsorted 10 x 10^6 sperm per insemination dose controls [37.7% vs. 25.0%; (Schenk et al., 2009)]. Schenk et al. (2009) retrospectively observed an increase in pregnancy rate of 7.9% when 2 x 10^6 sex-sorted sperm packaged in 0.25 or 0.5 mL straws was used to inseminate non-selected cows after 84 to 98 DIM compared to inseminations earlier in lactation. Turkish Holstein-Friesians inseminated with sex-sorted semen after 140 DIM experienced conceptions rates similar to conventional semen (31.8% vs. 40.9%, respectively) when pregnancy was determined at 31 days post AI (Karakaya et al., 2014). However, Karakaya et al. (2014) also reported high rates of loss (19.1%) by 62 DIM, possibly due to environmental heat stress or lower quality sex-sorted semen. In unsynchronized Holstein-Friesians inseminated with low-dose (2 x 10^6) sex-sorted sperm placed in uterine horns, sex-sorted semen conception rates were significantly lower than conventional semen controls [21% vs. 46%, respectively; (Andersson et al., 2006)]. It is important to note that improper uterine horn inseminations or delaying pregnancy determination to 90 days post AI could be influential factors affecting conception rates in the Andersson et al. (2006) study. A Brazilian study in 2013 evaluated the success of sex-sorted semen when utilized with estrus detection versus timed insemination and found that, while service rates were higher for timed inseminations, pregnancy per AI was higher for cows bred following estrus expression than after two timed insemination protocols [31.7% vs. 19.4% and 23.9%, respectively (Sá Filho et al., 2013)]. Overall, there was a trend for pregnancy rate to be lower for the estrus detection group compared to the two groups of timed insemination cows (14.3% vs. 18.3% and 23.2%), likely due to increased submission rate with timed AI (Sá Filho et

al., 2013). Multiple studies evaluating fertility of sex-sorted semen when processed at varying dosages found no appreciable improvement in conception rates (23.0% to 30.3%) when concentration was increased from 2.1 million to 3.5 million or 5 million sperm per straw (DeJarnette et al., 2008; DeJarnette et al., 2010).

Economic Aspects

Sex-sorted semen provides multiple economic benefits for the dairy industry including increased production of heifer calves, allowing producers to be more selective when choosing replacement females, reduced incidence of dystocia at parturition because of smaller calves, and increased genetic progress within the herd (De Vries et al., 2008). Unfortunately, widespread usage of sex-sorted semen has been hampered because of increased cost per straw which has been estimated at $20 to $50 more per insemination dose and because of reduced conception rates compared to conventional semen (Amann, 1999; Weigel, 2004; Norman et al., 2010). For sex-sorted semen to be an economical technology to integrate into reproductive practices, general fertility of a herd must be managed well initially (Seidel, 2003). This is further shown in dairy heifers by Weigel (2004) which reports that the cost of a heifer calf increases from $97 to $106 when conception rates for sex-sorted semen decline from 45% to 35%, assuming semen costs are $50. Thus, in order for sex-sorted semen to be economically beneficial, the value of producing a heifer calf over a bull calf must be twice the cost of production [i.e., if it costs $100 dollars to use sex-sorted semen, the resultant heifer calf must be worth at least $200 more to justify the added expense (Amann, 1999; Seidel, 2003)].

Extending the VWP may help to address some of the fertility issues associated with sex-sorted semen. The increased amounts of milk produced from extending the calving interval may also help to offset the additional cost of sex-sorted semen. Extending the VWP in high

producing cows could have economic benefits in terms of milk production (Arbel et al., 2001; Linderoth, 2005; Gobikrushanth et al., 2014). Gobikrushanth et al. (2014) found that extending the VWP from an average of 60 DIM (range: 57 to 63 DIM) to an average of 83 DIM (range: 64 to 121 DIM) improved conception rates by 5% and resulted in approximately 1,600 kg of additional milk yield per lactation in primiparous cows. However, no significant improvements were found for milk production in multiparous cows, which may be attributed to varying levels of production among multiparous cows (Gobikrushanth et al., 2014). When rbST was used to boost persistency of lactation, van Amburgh et al. (1997) showed that cows subjected to extended calving intervals of 16.5 months produced more milk compared to cows having a calving interval of ~13 months. With or without the aid of bST, it seems intuitive that cows which are milked longer would produce increased amounts of milk; the caveat of extended VWPs and subsequent extensions of calving intervals would center on the input cost of maintaining lactating cows for longer periods.

Though Inchaisri et al. (2010) stated that the economically optimal VWP is generally shorter (~60 DIM) for most cows, they also hypothesized that small extensions of the VWP may not negatively impact production from an economic standpoint while allowing producers to achieve increased fertility. Extending the VWP in high producing cows resulted in economic advantages of $0.19 and $0.12 per cow per day of lactation for primiparous and multiparous cows, respectively (Arbel et al., 2001). While this only equates to approximately $11 and $7 of increased revenue, respectively, combined with increased conception rates at later DIM, it demonstrates promise that extending the VWP in lactating cows may be beneficial for fertility and could potentially help to offset the increased cost of sex-sorted semen by increased milk production. Developing a way to improve the fertility of sex-sorted semen in lactating cows that

is economically advantageous to the producer could be instrumental for increasing the appeal of this reproductive technology for producers.

CHAPTER 3

MATERIALS AND METHODS

Lactating Holsteins (n = 183), average parity of 1.98 ± 0.97 (range 1 to 5), were housed in a sand-bedded free stall barn, fed *ad libitum* a total mixed ration of corn silage, grass silage, grain and hay (NRC, 2001), provided open access to fresh water, and were milked twice daily. Animal related procedures were approved by the Institutional Animal Care and Use Committee at the University of Tennessee, Knoxville.

Treatment Assignments

Cows calving between August and April over a two year period and having no obvious adverse health or reproductive conditions by 21 DIM (n = 137) were randomly allotted to one of two voluntary waiting periods (**VWP),** either 55 days [**VWP55** (n = 71)] or 85 days [**VWP85** (n = 66); (**Figure 3.1**)]. Reproductive tract scores (**RTS**) were assigned based on the size and length of the reproductive tract with one being the smallest and three being the largest by a single technician as described by Young et al. (2010).

Beginning at approximately 21 DIM, weekly transrectal ultrasound exams were conducted to confirm ovarian cyclicity [i.e. presence of a corpus luteum (**CL**); **Figure 3.1**]. Cyclicity was confirmed by the presence of a CL on either ovary for two of three weeks or visual observation of standing estrus before 55 DIM. Independent of treatment designation, 48 cows (35%) had not resumed cyclicity by 55 DIM and were removed from the trial and returned to normal herd management practices. Additionally, cows that began the synchronization protocol and developed health concerns that could possibly bias treatment results were removed from the study (n = 1), nine cows never expressed estrus during the study, and nine began estrus synchronization but did not express estrus before the study ended.




Figure 3.1. Experimental schematic and estrus synchronization protocol for VWP55 and VWP85 cows.
Following calving, cows were assigned to VWP55 or VWP85 treatment groups at 21 DIM. From 21 DIM to 55 DIM, resumption of cyclicity was monitored by weekly ultrasound scans such that all cows utilized were cyclic before 55 DIM. Relevant for cyclic cows and initial treatment designation, estrus synchronization began at ~55 or ~85 DIM. Cows were administered $PGF_{2\alpha}$ at appropriate times if a CL was present. Following two $PGF_{2\alpha}$ administrations (14 days apart) without estrus expression, cows were administered GnRH and CIDR. Seven days later, CIDRs were removed and $PGF_{2\alpha}$ was administered a final time. Any cow exhibiting estrus was submitted for AI with sex-sorted semen from one of two different Holstein bulls.

Estrus Synchronization and Detection

On a weekly basis, cows that were confirmed cyclic by 55 DIM were body condition scored (**BCS**) by a single evaluator and reported based on the standard, dairy industry accepted scale of 1 to 5, with 1 being emaciated and 5 being obese (Ferguson et al., 1994) at either 55 to 65 DIM (**VWP55**; n = 44) or 85 to 95 DIM (**VWP85**; n = 45). HeatWatch® (**HW**; Cow Chips, LLC, Manalapan, NJ) patches were attached to the sacral portion of the spine of each cow with adhesive. Each patch contained a radiotelemetric device that recorded the number and duration of each mount for cows exhibiting estrus. Mount information was sent to a computer database where estrus intensity and duration information was compiled and stored for later analysis.

Cows detected to have a CL present by ultrasound were administered $PGF_{2\alpha}$ (25 mg/mL im; Lutalyse®, *dinoprost tromethamine*, Zoetis, USA; **Figure 3.1**). If no CL was present, cows were monitored for spontaneous estrus until the next stage of the protocol (n = 25). Estrus detection occurred continuously via the HW system and by visual observation twice daily (0600 and 1900) in a deep sand bedded concrete alley way for seven consecutive days after each $PGF_{2\alpha}$ administration (**Figure 3.1**). Estrus was defined as at least two mounts of two seconds or greater within six consecutive hours if HW was the only detection source and/or the observation of a cow standing to be mounted by another. Cows detected in estrus were artificially inseminated ten to eleven hours before predicted ovulation, which was assumed to occur approximately 28 h (Hockett et al., 2005) after first recorded HW mount. After 14 days, cows not detected in estrus were examined by ultrasound, and if a CL was present, a second injection of $PGF_{2\alpha}$ was administered (**Figure 3.1**). Cows not detected in estrus and thus not artificially inseminated in the 7 days following the second prostaglandin administration received an injection of GnRH (50 µg/mL im; Cystorelin®, *gonadorelin diacetate tetrahydrate*, Merial Limited, Duluth, GA) and a

controlled intravaginal drug releasing device (CIDR; Eazi-Breed™ CIDR® Cattle Insert, Zoetis, USA) impregnated with 1.38 g of progesterone was placed intravaginally for 7 days (**Figure 3.1**). Following removal of the CIDR, a final dose of $PGF_{2\alpha}$ was administered if a CL was present (**Figure 3.1**).

Artificial Insemination & Pregnancy Determination

Cows exhibiting estrus were submitted for AI by one technician with sex-sorted semen using one of two different Holstein bulls. Sire conception rates for the two different bulls were +1.4 and +1.5 at the beginning and +0.7 and +0.9 at the end of the study, respectively. At the time of insemination, an AI score (**AIS)** relating the general difficulty of insemination was assessed on a subset of cows [score of 1 (reproductive tract easily traversed) to 4 (semen deposited in cervix) and modeled after an embryo transfer score described by Scenna et al. (2005)]. In addition, a cervix score (**CS)** was also recorded relating the overall size of the cervix as discerned via transrectal palpation [(score of 1 to 3 as described by Young et al. (2010)]. Pregnancy status was determined between 28 to 35 days after AI using transrectal ultrasonography. Maintenance of pregnancy was assessed at 60, 80 and 150 days after AI.

Statistical Analyses

In general, data were analyzed as a completely randomized design using PROC GLIMMIX (SAS, Version 9.4, SAS, Inc., Cary, NC, USA) with a binomial or normal distribution. In instances where an effect of year was significant, some estrus characteristics (i.e., mount duration) were analyzed as a randomized block design, blocking on year. All relevant variables were included in initial analyses and manual backwards selection was used to remove non-significant ($P > 0.05$) variables one-by-one from the model until only significant variables remained. Treatment differences from all analyses were determined using F-protected

least significant differences and reported as least squares means ± standard error of the mean (**SEM**).

All milk production values were segregated into three groups using the PROC UNIVARIATE procedure such that 25% of cows were classified as low production, 50% as medium production, and 25% as high production for each analysis. Body condition score was separated into two groups such that cows were evenly divided above and below the average score of 2.5. Additionally, parity was grouped as first, second, and third or greater due to limited observations of parities greater than three. All models included treatment, parity, and year as fixed effects. The statistical model for reproductive tract scores included milk production as a fixed effect; whereas, cyclicity analyses included treatment, parity, year, milk production, and RTS. Estrus characteristics models included treatment, parity, milk production (3 d average), and year. Additionally, the PROC LIFETEST procedure was used to compare percentage of estrus expression and time to estrus for treatments. Finally, conception rate models used the following variables in the analysis: treatment, RTS, parity, sire, BCS, milk production (day of AI), and year.

CHAPTER 4

RESULTS AND DISCUSSION

Milk production was similar for VWP55 and VWP85 cows when treatments were initially assigned at 21 DIM (41.33 ± 0.85 and 39.47 ± 0.82 kg, respectively; P = 0.1165), when cyclicity was confirmed at 55 DIM (43.57 ± 0.86 and 44.48 ± 0.80 kg, respectively; P = 0.4413), when estrus synchronization efforts began (43.28 ± 0.84 and 42.51 ± 0.79 kg, respectively; P = 0.5082), and at the time of AI (40.16 ± 1.46 and 40.89 ± 1.46 kg, respectively; P = 0.7250). Reproductive tract scores were similar for cows assigned to either a 55 day VWP or 85 day VWP when evaluated at 21 DIM (1.99 ± 0.07 and 1.94 ± 0.07, respectively; P = 0.6941), at 55 DIM when cyclicity was confirmed (1.64 ± 0.08 and 1.80 ± 0.08, respectively; P = 0.1647), at the time of estrus synchronization (1.73 ± 0.09 and 1.66 ± 0.09, respectively; P = 0.5406), and at the time of AI (1.57 ± 0.10 and 1.66 ± 0.10, respectively; P = 0.5549). The percentage of VWP55 and VWP85 cows cycling by 55 DIM was similar (77.29% ± 5.90 and 79.03% ± 5.29, respectively; P = 0.8158). Additionally, parity for cows assigned to a 55 versus 85 day VWP was similar for all cows beginning estrus synchronization (1.82 ± 0.14 vs. 2.13 ± 0.14; P = 0.1200, respectively). Body condition score was not different for VWP55 and VWP85 cows at onset of estrus synchronization efforts (2.31 ± 0.06 and 2.45 ± 0.06; P = 0.0805). Average rectal temperature at AI for VWP55 cows was similar to those of VWP85 cows (38.41°C ± 0.08 vs. 38.41°C ± 0.08; P = 0.9428). In summary, random treatment allocation at 21 DIM resulted in two groups of cows there were essentially equivalent in milk production, reproductive tract scores, ovarian cyclicity, parity, and body condition score. Hence, any inferences about estrus activity or conception rates were not influenced by these factors and were the result of our imposed treatment groups of either a 55 or 85 day VWP.

Estrus Activity

The percentage of cows confirmed to be cyclic by 55 DIM that exhibited natural or pharmacologically induced estrus was similar for cows after a VWP of 55 or 85 days (**Table 4.1**). An exhaustive review of the literature determined no previous studies examining estrus activity at these specific times in lactation; however, this finding is similar to another report that examined estrus expression between 70 and 100 DIM using only a radiotelemetry system in Holstein cows (Lopez et al., 2004). Furthermore, the responsiveness of cows having a CL was also similar for VWP55 and VWP85 cows when evaluation of estrus expression within seven days from the first $PGF_{2\alpha}$ administration (**Table 4.1**). Response rates for both VWP55 and VWP85 cows are similar to those reported by Momcilovic et al. (1998), Walton et al. (1987), and Simmons et al. (1979), which evaluated $PGF_{2\alpha}$ responsiveness between 40 and 60 DIM. While some report higher response rates (81.2%) to $PGF_{2\alpha}$ when cows were inseminated at an average of 83 DIM in both visual observation and HW herds (Jobst et al., 2000), our finding is consistent with studies not specifying DIM which reported $PGF_{2\alpha}$ response rates of 58% to 72% (Lauderdale et al., 1974; Archbald et al., 1993).

The time from $PGF_{2\alpha}$ administration to estrus was measured on a subset of cows (n = 38) using HW system. The time to estrus from the initial $PGF_{2\alpha}$ administration was similar for cows assigned to a VWP of 55 days and 85 days (**Table 4.1**). When all $PGF_{2\alpha}$ administrations from the estrus synchronization protocol were considered, time to estrus was still similar for both VWP55 and VWP85 cows (**Table 4.1**). Our fining is consistent with Macmillan et al. (1980) which reported that stage of lactation [(early or peak lactation versus late lactation (> 240 DIM)] did not impact interval to estrus after $PGF_{2\alpha}$ administration. Times to estrus reported herein are similar to other works reporting approximately

Table 4.1. Estrus response and activity for VWP55 and VWP85 cows[1]

	Treatment[2]		
	VWP55	VWP85	P-Value
Overall percentage that exhibited estrus[3]	79.55[a] ± 6.08	77.78[a] ± 6.20	0.8393
Percentage exhibiting estrus within 7 d of $PGF_{2\alpha}$ [4]	64.37[a] ± 11.36	57.14[a] ± 9.57	0.6275
Time to estrus for first $PGF_{2\alpha}$ administration (h)[5]	65.79[a] ± 7.58	72.65[a] ± 6.27	0.4908
Time to estrus for all $PGF_{2\alpha}$ administrations (h)[5]	62.28[a] ± 6.68	73.00[a] ± 5.46	0.2218
Number of mounts	9.05[a] ± 1.75	10.54[a] ± 1.52	0.5246
Mount duration (s)	4.59[a] ± 0.43	4.99[a] ± 0.41	0.2602
Estrus length (h)[6]	8.16[a] ± 1.64	9.17[a] ± 1.39	0.6389

[1]The total number of cows that exhibited estrus, hours to estrus from $PGF_{2\alpha}$, and estrus characteristics as detected by HW were recorded for VWP55 and VWP85 cows.
[2]Least squares means ± SEM reported. Treatment means within a row having unlike superscripts differ ($P < 0.05$)
[3]Includes all cows that exhibited natural or $PGF_{2\alpha}$ induced estrus detected by visual observation and/or HW
[4]Only includes cows exhibiting estrus in response to any $PGF_{2\alpha}$ administration within seven days
[5] As detected by HW

[6]Length of time between the first and last HW recorded mount

73 h to estrus in lactating cows (Renegar et al., 1978; Archbald et al., 1993; Walker et al., 1996). Survival curve analysis indicates that cows exhibited estrus at similar rates after a VWP of 55 day and 85 days for the initial $PGF_{2\alpha}$ administration and when all $PGF_{2\alpha}$ administrations were considered (P = 0.3848 and P = 0.2480, respectively; **Figure 4.2**). Similar to other reports, the largest increase in cows expressing estrus occurred between 40 and 80 hours following both administrations of $PGF_{2\alpha}$ [**Figure 4.2**; (Lauderdale et al., 1974; Walton et al., 1987)].

Number of mounts per estrus event, duration of each mount, and the total length of time that estrus was exhibited were similar when efforts to synchronize estrus began after a VWP of 55 or 85 days (**Table 4.1**). Independent of DIM when estrus was exhibited in cyclic cows, the number of mounts for both VWP55 and VWP85 cows is consistent with other reports in various settings and situations (Britt et al., 1986; Walker et al., 1996; Dransfield et al., 1998; Xu and Burton, 1999; Lopez et al., 2004). Our result for number of mounts contrasts with Peralta et al. (2005), who found that average number of mounts were higher for cows that were less than 80 DIM compared to cows ranging from 80 to more than 140 DIM. Average mount duration in our study was one to two seconds longer than reported by Lopez et al. (2004), Walker et al. (1996), or Xu and Burton (1999). Mounts of longer duration may have been detected due to facilities (deep sand bedded alley way) and because cows prefer more secure footing to exhibit estrual signs (Britt et al., 1986). Mounts of longer duration may be more easily observed when utilizing only visual estrus detection. Consistent with our findings, most studies evaluating the length of time cows exhibited estrus, independent of DIM, ranged from seven to nine hours (Walker et al., 1996; Dransfield et al., 1998; Xu and Burton, 1999; Lopez et al., 2004), though Britt et al. (1986) reported longer estrus lengths (~13 h). Similarities in estrus activity of cows when the VWP was either 55 days or 85 days is likely due




Figure 4.2. Percentage of estrus expression over time after $PGF_2\alpha$ administration
Depiction of survival curve for proportion of VWP55 and VWP85 cows expressing estrus by a given number of hours from the first $PGF_{2\alpha}$ administration (Panel A) or all $PGF_{2\alpha}$ administrations (Panel B) as recorded using the HW system (i.e., 30% of VWP85 cows had exhibited estrus by 60 hours after $PGF_{2\alpha}$ when all administrations were considered).

to the inherent similarities of our treatment groups for general production, cyclicity status, and overall management.

In our study, estrus activity was more influenced by average milk production in the three days around estrus (i.e., one day before, day of estrus, and one day after). Specifically, when estrus activity within seven days of the first $PGF_{2\alpha}$ was considered, larger percentages of cows with milk production ranging from 33.48 to 46.95 kg (medium) and from 46.95 to 62.78 kg (high) exhibited estrus compared to the cows producing 24.09 to 33.48 kg (low; 66.75% ± 11.13 and 85.03% ± 8.22 vs. 24.68% ± 8.81, respectively; P = 0.0014). Increased rates of estrus expression by higher producing cows were not necessarily expected because previous research has shown that higher producing cows have increased negative energy balance (Harrison et al., 1990; Butler, 2003; Wathes et al., 2007; Ospina et al., 2010) and higher rates of $PGF_{2\alpha}$ clearance (Davis et al., 1980; Shrestha et al., 2012), which may lead to reduction of estrus expression.

Artificial Insemination and Conception Rates

Artificial insemination score was not different among treatments (1.43 ± 0.13 vs. 1.68 ± 0.14; P = 0.2024), and neither was cervix score (1.30 ± 0.12 vs. 1.50 ± 0.12; P = 0.2520). No differences in sire conception rates were observed (35.29% ± 8.20 vs. 38.89% ± 8.13; P = 0.7567). Conception rates at approximately 30 days after AI were higher for VWP85 cows compared to VWP55 cows (P = 0.0181; **Figure 4.3, Panel A**), and were higher than reported conventional semen conception rates for lactating cows (Andersson et al., 2006; Norman et al., 2009; Chebel and Santos, 2010; DeJarnette et al., 2010; Karakaya et al., 2014). Benefits of extending the VWP by 30 days (VWP85) were still observed at 60 (P = 0.0166; **Figure 4.3, Panel B**), 80 (P = 0.0471; **Figure 4.3, Panel C**), and 150 days after AI (P = 0.0471; **Figure 4.3, Panel D**). Herein, pregnancy loss between 30 and 60 days after AI (11.5%; 3/26) was similar to




Figure 4.3. Conception rates for VWP55 and VWP85 cows
Conception rates at 30 d (Panel A), 60 d (Panel B), 80 d (Panel C), and 150 d (Panel D) after AI for lactating cows bred with sex-sorted semen after a 55 d or 85d VWP. Least squares means are reported, and bars with different letters were significantly different ($P \leq 0.05$).

reports by Schenk et al. (2009), but less than the 19.1% (9/47) reported in Holsteins inseminated with sex-sorted semen in Turkey (Karakaya et al., 2014). Cows that were not confirmed pregnant to first AI with sex-sorted semen after a 55 or 85 day VWP required a total of 2.5 and 1.5 services per conception, respectively (P = 0.0010).

Fertility of sex-sorted semen in our study using two different Holstein bulls having sire conception rates of +0.7 and +0.9 when the VWP was extended to 85 DIM in confirmed cyclic cows was comparable to studies using conventional semen at later DIM reporting conception rates (Hillers et al., 1984; Tenhagen et al., 2003; Niozas et al., 2015) and overall fertility (Whitmore et al., 1974; Britt, 1975; Stevenson and Phatak, 2005). Our results suggest that current practices for VWP (Caraviello et al., 2006; DeJarnette et al., 2007) may contribute to poor success rates in studies using sex-sorted semen after a VWP of 55 days, which is more consistent with industry standards. Though some sex-sorted semen studies reported inseminations at average DIM similar to our treatment groups, a VWP longer than the industry standard of ~55 DIM was not specifically controlled for because sperm dosages and comparison of estrus detection versus timed AI were the original endpoints, resulting in reduced conception rates by comparison [29.4% and 31.7%, respectively; (DeJarnette et al., 2008; Sá Filho et al., 2013)]. There are instances where extended VWPs or increased DIM at time of insemination did not impact pregnancy rates for conventional or sex-sorted semen, respectively (Arbel et al., 2001; Schenk et al., 2009). Interestingly in one trial, Schenk et al. (2009) selected cows for absence of uterine or ovarian abnormalities before starting the study, similar to our experimental design, though did not explicitly monitor cyclicity and still achieved similar fertility to our results after timed inseminations (average 80 DIM) following OvSynch synchronization. In a separate trial, Schenk et al. (2009) reported pregnancy rates of approximately 25% when cows

not selected for reproductive status were inseminated with sex-sorted semen, but upon retrospective analysis cows inseminated between 84 to 98 DIM had pregnancy rates that were 7.9 percentage points higher compared to those inseminated earlier in lactation.

After statistical analysis, only VWP length significantly influenced conception rates in our study. Although DeJarnette et al. (2008) reported that parity tended to impact conception rates of lactating cows inseminated with sex-sorted semen, we did not observe that effect (first: 36.67% ± 8.80, second: 36.00% ± 9.60, third or greater: 40.00% ± 12.65; P = 0.9660). This is not surprising since the majority of cows used in our study were first and second lactation due to recent repopulation of our dairy research herd. Additionally, conception rates were similar among reproductive tract scores of 1, 2, and 3 (40.63% ± 8.68, 33.33% ± 8.2, and 40.00% ± 21.91, respectively; P = 0.8242) and numerically higher than those reported by Young et al. (2010).

Independent of VWP55 and VWP85 treatment, milk production increased significantly as parity increased from first to third or greater at all stages of the study (P < 0.0001; **Table 4.2**), which is generally consistent with Tanaka et al. (2008) and Lee and Kim (2006). Moreover, parity impacted reproductive tract score (P ≤ 0.08) throughout the entire study (**Table 4.3**). First and second parity cows had smaller numerical reproductive tract scores compared to third and greater parity cows, similar to reports from Young et al. (2010).

Resumption of cyclicity was affected by parity and milk production at 55 DIM, independent of VWP55 or VWP85 treatment. The percentage of first parity cows that were cyclic by 55 DIM was significantly lower than second and third or greater parity cows (P = 0.0093; **Table 4.4**) and generally similar to other reports in the literature (Bulman and Lamming, 1978; Tanaka et al., 2008; Chebel and Santos, 2010). In our study, cows ranging in milk

Table 4.2. Influence of parity on milk production[1]

	Parity[2]			
	First	Second	Third or greater	P-Value
Treatment assignment[3]	$31.65^{c} \pm 0.90$	$42.75^{b} \pm 0.96$	$46.80^{a} \pm 1.22$	<0.0001
Cyclicity confirmation[4]	$33.34^{c} \pm 0.95$	$46.97^{b} \pm 0.92$	$51.76^{a} \pm 1.18$	<0.0001
Estrus synchronization[5]	$33.18^{c} \pm 0.94$	$45.97^{b} \pm 0.90$	$49.54^{a} \pm 1.16$	<0.0001
Time of AI	$32.99^{c} \pm 0.96$	$43.79^{b} \pm 1.06$	$50.16^{a} \pm 1.39$	<0.0001

[1]The influence of parity on milk production (kg), independent of VWP55/VWP85 treatment
[2]Least squares means ± SEM reported. Means within a row having unlike superscripts differ ($P < 0.0001$)
[3]At 21 DIM
[4]At 55 DIM
[5]Either 55 DIM or 85 DIM, depending on treatment designation

Table 4.3. Influence of parity on reproductive tract scores[1]

	Parity[2]			
	First	Second	Third or greater	P-Value
Treatment assignment[3]	$1.80^{b} \pm 0.08$	$1.96^{ab} \pm 0.08$	$2.14^{a} \pm 0.11$	0.0460
Cyclicity confirmation[4]	$1.54^{b} \pm 0.08$	$1.72^{b} \pm 0.09$	$2.05^{a} \pm 0.12$	0.0021
Estrus synchronization[5]	$1.56^{b} \pm 0.10$	$1.51^{b} \pm 0.10$	$2.02^{a} \pm 0.13$	0.0063
Time of AI	$1.53^{b} \pm 0.11$	$1.52^{b} \pm 0.12$	$1.93^{a} \pm 0.16$	0.0829

[1]The influence of parity on reproductive tract scores, independent of VWP55/VWP85 treatment

[2]Least squares means ± SEM reported. Means within a row having unlike superscripts differ ($P < 0.10$)

[3]At 21 DIM

[4]At 55 DIM

[5]Either 55 DIM or 85 DIM, depending on treatment designation

production from 24.95 to 35.90 kg (low) were more likely to be cyclic by 55 DIM than cows ranging from 35.90 to 51.10 kg (medium) or from 51.10 to 62.32 kg (high; P = 0.0006; **Table 4.4**).

Conclusions

Though other studies report information about VWPs and fertility increases at later DIM, few were controlled for this specific hypothesis, and to our knowledge, none were controlled experiments using sex-sorted semen (Whitmore et al., 1974; Britt, 1975; Hillers et al., 1984; Weller and Folman, 1990; Tenhagen et al., 2003; Stevenson and Phatak, 2005; Niozas et al., 2015). In circumstances where factors such as milk production, reproductive tract scores, ovarian cyclicity, parity, and BCS are similar, our study is the first that we are aware of to indicate improvements in fertility of sex-sorted semen after extending the VWP to 85 days. While the underlying mechanisms of these benefits are unclear, differences in conception rates were not a result of differing ovarian cyclicity, estrus response to $PGF_{2\alpha}$, or estrus characteristics. While difficult to explain, improved fertility of cows after extending the VWP to 85 days may be attributed to important cellular changes occurring in the uterus after the time of involution that may result in enhanced ability of the uterus to support a pregnancy (Sheldon et al., 2008). It is also possible for benefits of extending the VWP to be related to increased exposure to hormones produced by another 1 to 1.5 estrous cycles before insemination (Thatcher and Wilcox, 1973; Sakaguchi, 2010). The possibility also exists that cows may have returned to positive energy balance (Butler, 2003, 2005) before inseminations begin could be an influential factor.

Table 4.4. Influence of parity and milk production on resumption of ovarian cyclicity[1]

	Cycling by 55 DIM (%)	P-Value
Parity		0.0093
First	$52.89^{b} \pm 9.91$	
Second	$86.39^{a} \pm 4.65$	
Third or greater	$86.56^{a} \pm 6.17$	
Milk Production		0.0006
Low (24.95 – 35.90 kg)	$94.61^{A} \pm 3.15$	
Medium (35.90 – 51.10 kg)	$56.82^{B} \pm 7.08$	
High (51.10 – 62.32 kg)	$66.51^{B} \pm 10.67$	

[1]The influence of parity and milk production resumption of ovarian cyclicity by 55 DIM, independent of VWP55/VWP85 treatment

[2]Least squares means ± SEM reported. Means within a column having unlike superscripts of the same case differ ($P < 0.05$)

Implementing the use of sex-sorted semen for lactating cow inseminations may be possible under specific management conditions, specifically, when cows are managed to be healthy and cyclic, estrus detection is employed, and the VWP is extended to 85 DIM.

Implications

Extension of the VWP from 55 days to 85 days improved conception rates for sex-sorted semen in lactating Holstein cows that were previously determined cyclic and inseminated following detection of standing estrus. $PGF_{2\alpha}$ responsiveness and estrus characteristics were similar between VWP55 and VWP85 cows suggesting that cyclic cows under similar management are capable of exhibiting estrus at varying stages of lactation. Preliminary economic analysis suggests that extending the VWP to 85 days when using sex-sorted semen in lactating cows can be beneficial.

LITERATURE CITED

Amann, R. P. 1999. Issues affecting commercialization of sexed sperm. Theriogenology 52: 1441-1457.

Andersson, M., J. Taponen, M. Kommeri and M. Dahlbom 2006. Pregnancy rates in lactating Holstein–Friesian cows after artificial insemination with sexed sperm. Reproduction in Domestic Animals 41: 95-97.

Arbel, R., Y. Bigun, E. Ezra, H. Sturman and D. Hojman 2001. The effect of extended calving intervals in high-yielding lactating cows on milk production and profitability. Journal of Dairy Science 84: 600-608.

Archbald, L. F., C. Risco, P. Chavatte, S. Constant, T. Tran, E. Klapstein and J. Elliot 1993. Estrus and pregnancy rate of dairy cows given one or two doses of prostaglandin F2 alpha 8 or 24 hours apart. Theriogenology 40: 873-884.

Barker, A. R., F. N. Schrick, M. J. Lewis, H. H. Dowlen and S. P. Oliver 1998. Influence of clinical mastitis during early lactation on reproductive performance of Jersey cows. Journal of Dairy Science 81: 1285-1290.

Bauman, D. E. and W. B. Currie 1980. Partitioning of nutrients during pregnancy and lactation: A review of mechanisms involving homeostasis and homeorhesis. Journal of Dairy Science 63: 1514-1529.

Beam, S. W. and W. R. Butler 1999. Effects of energy balance on follicular development and first ovulation in postpartum dairy cows. Journal of Reproduction and Fertility 54: 411-424.

Bodmer, M., F. Janett, M. Hässig, N. den Daas, P. Reichert and R. Thun 2005. Fertility in heifers and cows after low dose insemination with sex-sorted and non-sorted sperm under field conditions. Theriogenology 64: 1647-1655.

Britt, J. H. 1975. Review- Early postpartum breeding in dairy cows Journal of Dairy Science 58: 266-271.

Britt, J. H., R. G. Scott, J. D. Armstrong and M. D. Whitacre 1986. Determinants of estrous behavior in lactating Holstein cows. Journal of Dairy Science 69: 2195-2202.

Bulman, D. C. and G. E. Lamming 1978. Milk progesterone levels in relation to conception, repeat breeding and factors influencing acyclicity in dairy cows. Journal of Reproduction and Fertility 54: 447-458.

Butler, S. T., I. A. Hutchinson, A. R. Cromie and L. Shalloo 2014. Applications and cost benefits of sexed semen in pasture-based dairy production systems. Animal 8: 165-172.

Butler, W. R. 2003. Energy balance relationships with follicular development, ovulation and fertility in postpartum dairy cows. Livestock Production Science 83: 211-218.

Butler, W. R. 2005. Nutrition, negative energy balance and fertility in the postpartum dairy cow. Cattle Practice 13: 13-18.

Butler, W. R. 1998. Review: Effect of protein nutrition on ovarian and uterine physiology in dairy cattle. Journal of Dairy Science 81: 2533-2539.

Caraviello, D. Z., K. A. Weigel, P. M. Fricke, M. C. Wiltbank, M. J. Florent, N. B. Cook, K. V. Nordlund, N. R. Zwald and C. L. Rawson 2006. Survey of management practices on reproductive performance of dairy cattle on large US commercial farms. Journal of Dairy Science 89: 4723-4735.

Cerri, R. L. A., J. E. P. Santos, S. O. Juchem, K. N. Galvão and R. C. Chebel 2004. Timed artificial insemination with estradiol cypionate or insemination at estrus in high-producing dairy cows. Journal of Dairy Science 87: 3704-3715.

Chebel, R. C., F. S. Guagnini, J. E. P. Santos, J. P. Fetrow and J. R. Lima 2010. Sex-sorted semen for dairy heifers: Effects on reproductive and lactational performances. Journal of Dairy Science 93: 2496-2507.

Chebel, R. C. and J. E. P. Santos 2010. Effect of inseminating cows in estrus following a presynchronization protocol on reproductive and lactation performances. Journal of Dairy Science 93: 4632-4643.

Crichton, E., S. Huffman, K. McSweeney and J. Schenk 2006. Artificial insemination of lactating Holstein cows with sexed sperm. Reproduction, Fertility and Development 18: 281-281.

Darwash, A. O., G. E. Lamming and J. A. Wooliams 1997. The phenotypic association between the interval to postpartum ovulation and traditional measures of fertility in dairy cattle. Animal Science 65: 9-16.

Davis, A. J., I. R. Fleet, F. A. Harrison and F. M. M. Walker 1980. Pulmonary metabolism of prostaglandin F2 alpha in the conscious non-pregnant ewe and sow. Journal of physiology 301: 86.

De Vries, A., M. Overton, J. Fetrow, K. Leslie, S. Eicker and G. Rogers 2008. Exploring the impact of sexed semen on the structure of the dairy industry. Journal of Dairy Science 91: 847-856.

DeJarnette, J. M., M. A. Leach, R. L. Nebel, C. E. Marshall, C. R. McCleary and J. F. Moreno 2011. Effects of sex-sorting and sperm dosage on conception rates of Holstein heifers: Is comparable fertility of sex-sorted and conventional semen plausible? Journal of Dairy Science 94: 3477-3483.

DeJarnette, J. M., C. R. McCleary, M. A. Leach, J. F. Moreno, R. L. Nebel and C. E. Marshall 2010. Effects of 2.1 and 3.5 x 10^6 sex-sorted sperm dosages on conception rates of Holstein cows and heifers. Journal of Dairy Science 93: 4079-4085.

DeJarnette, J. M., R. L. Nebel and C. E. Marshall 2009. Evaluating the success of sex-sorted semen in US dairy herds from on farm records. Theriogenology 71: 49-58.

DeJarnette, J. M., R. L. Nebel, C. E. Marshall, J. F. Moreno, C. R. McCleary and R. W. Lenz 2008. Effect of sex-sorted sperm dosage on conception rates in Holstein heifers and lactating cows. Journal of Dairy Science 91: 1778-1785.

DeJarnette, J. M., C. G. Sattler, C. E. Marshall and R. L. Nebel 2007. Voluntary waiting period management practices in dairy herds participating in a progeny test program. Journal of Dairy Science 90: 1073-1079.

Dematawena, C. M. B. and P. J. Berger 1997. Effect of dystocia on yield, fertility, and cow losses and an economic evaluation of dystocia scores for Holsteins. Journal of Dairy Science 80: 754-761.

Dhaliwal, G. S., R. D. Murray and H. Dobson 1996. Effects of milk yield, and calving to first service interval, in determining herd fertility in dairy cows. Animal Reproduction Science 41: 109-117.

Domecq, J. J., A. L. Skidmore, J. W. Lloyd and J. B. Kaneene 1997. Relationship between body condition scores and conception at first artificial insemination in a large dairy herd of high yielding Holstein cows. Journal of Dairy Science 80: 113-120.

Dransfield, M. B. G., R. L. Nebel, R. E. Pearson and L. D. Warnick 1998. Timing of insemination for dairy cows identified in estrus by a radiotelemetric estrus detection system. Journal of Dairy Science 81: 1874-1882.

El-Zarkouny, S. Z., J. A. Cartmill, B. A. Hensley and J. S. Stevenson 2004. Pregnancy in dairy cows after synchronized ovulation regimens with or without presynchronization and progesterone. Journal of Dairy Science 87: 1024-1037.

FAO 2011. World Livestock 2011-Livestock in food security. Rome, FAO.

Ferguson, J. D., D. T. Galligan and N. Thomsen 1994. Principal descriptors of body condition score in Holstein cows. Journal of Dairy Science 77: 2695-2703.

Gaafar, H. M. A., Sh M. Shamiah, M. A. Abu El-Hamd, A. A. Shitta and M. A. Tag El-Din 2011. Dystocia in Friesian cows and its effects on postpartum reproductive performance and milk production. Tropical Animal Health and Production 43: 229-234.

Gillund, P., O. Reksen, Y. T. Gröhn and K. Karlberg 2001. Body condition related to ketosis and reproductive performance in Norwegian dairy cows. Journal of Dairy Science 84: 1390-1396.

Gobikrushanth, M., A. De Vries, J. E. P. Santos, C. A. Risco and K. N. Galvão 2014. Effect of delayed breeding during the summer on profitability of dairy cows. Journal of Dairy Science 97: 4236-4246.

Gröhn, Y. T. and P. J. Rajala-Schultz 2000. Epidemiology of reproductive performance in dairy cows. Animal Reproduction Science 60–61: 605-614.

Harrison, R. O., S. P. Ford, J. W. Young, A. J. Conley and A. E. Freeman 1990. Increased milk production versus reproductive and energy status of high producing dairy cows. Journal of Dairy Science 73: 2749-2758.

Healy, A. A., J. K. House and P. C. Thomson 2013. Artificial insemination field data on the use of sexed and conventional semen in nulliparous Holstein heifers. Journal of Dairy Science 96: 1905-1914.

Hillers, J. K., P. L. Senger, R. L. Darlington and W. N. Fleming 1984. Effects of production, season, age of cow, days dry, and days in milk on conception to first service in large commercial dairy herds. Journal of Dairy Science 67: 861-867.

Hockett, M. E., R. A. Almeida, N. R. Rohrbach, S. P. Oliver, H. H. Dowlen and F. N. Schrick 2005. Effects of induced clinical mastitis during preovulation on endocrine and follicular function. Journal of Dairy Science 88: 2422-2431.

Hohenboken, W. D. 1999. Applications of sexed semen in cattle production. Theriogenology 52: 1421-1433.

Inchaisri, C., R. Jorritsma, P. L. A. M. Vos, G. C. van der Weijden and H. Hogeveen 2010. Economic consequences of reproductive performance in dairy cattle. Theriogenology 74: 835-846.

Jobst, S. M., R. L. Nebel, M. L. McGilliard and K. D. Pelzer 2000. Evaluation of reproductive performance in lactating dairy cows with prostaglandin F2 alpha gonadotropin-releasing hormone, and timed artificial insemination. Journal of Dairy Science 83: 2366-2372.

Karakaya, E., G. Yilmazbas-Mecitoglu, A. Keskin, A. Alkan, U. Tasdemir, J. E. P. Santos and A. Gumen 2014. Fertility in dairy cows after artificial insemination using sex-sorted sperm or conventional semen. Reproduction in Domestic Animals 49: 333-337.

Kasimanickam, R., J. M. Cornwell and R. L. Nebel 2005. Fertility following fixed-time AI or insemination at observed estrus in Ovsynch and Heatsynch programs in lactating dairy cows. Theriogenology 63: 2550-2559.

Lauderdale, J. W., B. E. Seguin, J. N. Stellflug, J. R. Chenault, W. W. Thatcher, C. K. Vincent and A. F. Loyancano 1974. Fertility of cattle following PGF 2 alpha injection. Journal of Animal Science 38: 964-967.

Lee, J. Y. and I. H. Kim 2006. Advancing parity is associated with high milk production at the cost of body condition and increased periparturient disorders in dairy herds. Journal of Veterinary Science 7: 161-166.

Linderoth, S. 2005. Don't cheat on your voluntary waiting period. Dairy Herd Management 42: 32-35.

Loeffler, S. H., M. J. de Vries and Y. H. Schukken 1999. The effects of time of disease occurrence, milk yield, and body condition on fertility of dairy cows. Journal of Dairy Science 82: 2589-2604.

Lopez, H., L. D. Satter and M. C. Wiltbank 2004. Relationship between level of milk production and estrous behavior of lactating dairy cows. Animal Reproduction Science 81: 209-223.

Lucy, M. C. 2001. ADSA Foundation Scholar Award - Reproductive loss in high-producing dairy cattle: Where will it end? Journal of Dairy Science 84: 1277-1293.

Macmillan, K. L., A. M. Day and J. F. Smith 1980. Onset of oestrus and fertility in lactating dairy cows injected with an analogue of prostaglandin F2 alpha, cloprostenol. Animal Reproduction Science 3: 171-180.

McCullock, K., D. L. K. Hoag, J. Parsons, M. Lacy, G. E. Seidel and W. Wailes 2013. Factors affecting economics of using sexed semen in dairy cattle. Journal of Dairy Science 96: 6366-6377.

Miller, R. H., H. D. Norman, M. T. Kuhn, J. S. Clay and J. L. Hutchison 2007. Voluntary waiting period and adoption of synchronized breeding in dairy herd improvement herds. Journal of Dairy Science 90: 1594-1606.

Momcilovic, D., L. F. Archbald, A. H. Walters, T. Tran, D. Kelbert, C. Risco and W. W. Thatcher 1998. Reproductive performance of lactating dairy cows treated with

gonadotrophin-releasing hormone and/or prostaglandin F2 alpha for synchronization of estrus and ovulation. Theriogenology 50: 1131-1139.

Moreira, F., C. Orlandi, C. A. Risco, R. Mattos, F. Lopes and W. W. Thatcher 2001. Effects of presynchronization and bovine somatotropin on pregnancy rates to a timed artificial insemination protocol in lactating dairy cows. Journal of Dairy Science 84: 1646-1659.

Niozas, G., G. Tsousis, C. Boscos, H. Bollwein and M. Kaske 2015. Effects of extension of voluntary waiting period on reproductive performance in high yielding dairy cows. Reproduction in Domestic Animals 50: 66-66.

Norman, H. D., J. L. Hutchison and R. H. Miller 2010. Use of sexed semen and its effect on conception rate, calf sex, dystocia, and stillbirth of Holsteins in the United States. Journal of Dairy Science 93: 3880-3890.

Norman, H. D., J. R. Wright, S. M. Hubbard, R. H. Miller and J. L. Hutchison 2009. Reproductive status of Holstein and Jersey cows in the United States. Journal of Dairy Science 92: 3517-3528.

NRC 2001. Nutrient requirements of dairy cattle. National Academies of Science 7th ed.

Okano, A. and T. Tomizuka 1987. Ultrasonic observation of postpartum uterine involution in the cow. Theriogenology 27: 369-376.

Ospina, P. A., D. V. Nydam, T. Stokol and T. R. Overton 2010. Associations of elevated nonesterified fatty acids and β-hydroxybutyrate concentrations with early lactation reproductive performance and milk production in transition dairy cattle in the Northeastern United States. Journal of Dairy Science 93: 1596-1603.

Peralta, O. A., R. E. Pearson and R. L. Nebel 2005. Comparison of three estrus detection systems during summer in a large commercial dairy herd. Animal Reproduction Science 87: 59-72.

Renegar, R. H., H. D. Hafs, J. H. Britt and T. D. Carruthers 1978. Luteolysis, growth-hormone, glucocorticoids, prolactin and milk-production in lactating dairy cows given prostaglandin F2 alpha. Journal of Animal Science 47: 532-537.

Roche, J. F. 2006. The effect of nutritional management of the dairy cow on reproductive efficiency. Animal Reproduction Science 96: 282-296.

Roche, J. R., N. C. Friggens, J. K. Kay, M. W. Fisher, K. J. Stafford and D. P. Berry 2009. Invited review: Body condition score and its association with dairy cow productivity, health, and welfare. Journal of Dairy Science 92: 5769-5801.

Royal, M. D., A. O. Darwash, A. P. F. Flint, R. Webb, J. A. Woolliams and G. E. Lamming 2000a. Declining fertility in dairy cattle: Changes in traditional and endocrine parameters of fertility. Animal Science 70: 487-501.

Royal, M., G. E. Mann and A. P. F. Flint 2000b. Strategies for reversing the trend towards subfertility in dairy cattle. The Veterinary Journal 160: 53-60.

Sa Filho, M. F., H. Ayres, R. M. Ferreira, M. Nichi, M. Fosado, E. P. Campos Filho and P. S. Baruselli 2010. Strategies to improve pregnancy per insemination using sex-sorted semen in dairy heifers detected in estrus. Theriogenology 74: 1636-1642.

Sá Filho, M. F., M. F. Mendanha, R. V. Sala, F. J. Carvalho, L. H. C. Guimarães and P. S. Baruselli 2013. Use of sex-sorted sperm in lactating dairy cows upon estrus detection or following timed artificial insemination. Animal Reproduction Science 143: 19-23.

Sakaguchi, M. 2010. Oestrous expression and relapse back into anoestrus at early postpartum ovulations in fertile dairy cows. Veterinary Record 167: 446-450.

Sakaguchi, M., Y. Sasamoto, T. Suzuki, Y. Takahashi and Y. Yamada 2004. Postpartum ovarian follicular dynamics and estrous activity in lactating dairy cows. Journal of Dairy Science 87: 2114-2121.

Sales, J. N. S., K. A. L. Neves, A. H. Souza, G. A. Crepaldi, R. V. Sala, M. Fosado, E. P. Campos, M. de Faria, M. F. Sa and P. S. Baruselli 2011. Timing of insemination and fertility in dairy and beef cattle receiving timed artificial insemination using sex-sorted sperm. Theriogenology 76: 427-435.

Santos, J. E. P., R. S. Bisinotto, E. S. Ribeiro, F. S. Lima, L. F. Greco, C. R. Staples and W. W. Thatcher 2010. Applying nutrition and physiology to improve reproduction in dairy cattle. Reproduction in Domestic Ruminants Vii 67: 387-403.

Santos, J. E. P., W. W. Thatcher, R. C. Chebel, R. L. A. Cerri and K. N. Galvão 2004. The effect of embryonic death rates in cattle on the efficacy of estrus synchronization programs. Animal Reproduction Science 82–83: 513-535.

Scenna, F. N., M. E. Hockett, T. M. Towns, A. M. Saxton, N. R. Rohrbach, M. E. Wehrman and F. N. Schrick 2005. Influence of a prostaglandin synthesis inhibitor administered at embryo transfer on pregnancy rates of recipient cows. Prostaglandins & Other Lipid Mediators 78: 38-45.

Schefers, J. M., K. A. Weigel, C. L. Rawson, N. R. Zwald and N. B. Cook 2010. Management practices associated with conception rate and service rate of lactating Holstein cows in large, commercial dairy herds. Journal of Dairy Science 93: 1459-1467.

Schenk, J. L., D. G. Cran, R. W. Everett and G. E. Seidel 2009. Pregnancy rates in heifers and cows with cryopreserved sexed sperm: Effects of sperm numbers per inseminate, sorting pressure and sperm storage before sorting. Theriogenology 71: 717-728.

Schrick, F. N., M. E. Hockett, A. M. Saxton, M. J. Lewis, H. H. Dowlen and S. P. Oliver 2001. Influence of subclinical mastitis during early lactation on reproductive parameters. Journal of Dairy Science 84: 1407-1412.

Seidel, G. E. 2003. Economics of selecting for sex: the most important genetic trait. Theriogenology 59: 585-598.

Seidel, G. E. 2007. Overview of sexing sperm. Theriogenology 68: 443-446.

Seidel, G. E. and D. L. Garner 2002. Current status of sexing mammalian spermatozoa. Reproduction 124: 733-743.

Seidel, G. E. and J. L. Schenk 2008. Pregnancy rates in cattle with cryopreserved sexed sperm: Effects of sperm numbers per inseminate and site of sperm deposition. Animal Reproduction Science 105: 129-138.

Sheldon, I. M., E. J. Williams, A. N. A. Miller, D. M. Nash and S. Herath 2008. Uterine diseases in cattle after parturition. The Veterinary Journal 176: 115-121.

Shrestha, H. K., M. A. Beg, R. R. Burnette and O.J. Ginther 2012. Plasma clearance and half-life of prostaglandin F2 alpha: A comparison between mares and heifers. Biology of Reproduction 87: 18, 11-16.

Shrestha, H. K., T. Nakao, T. Higaki, T. Suzuki and M. Akita 2004. Resumption of postpartum ovarian cyclicity in high-producing Holstein cows. Theriogenology 61: 637-649.

Simmons, K. R., S. C. Moses and B. L. Perkins 1979. Prostaglandin in milk, days open, and estrus detection in dairy cows treated with prostaglandin F2 alpha. Journal of Dairy Science 62: 1443-1448.

Stevenson, J. S. 2001. Reproductive management of dairy cows in high milk-producing herds. Journal of Dairy Science 84, Supplement: E128-E143.

Stevenson, J. S. and A. P. Phatak 2005. Inseminations at estrus induced by presynchronization before application of synchronized estrus and ovulation. Journal of Dairy Science 88: 399-405.

Tanaka, T., M. Arai, S. Ohtani, S. Uemura, T. Kuroiwa, S. Kim and H. Kamomae 2008. Influence of parity on follicular dynamics and resumption of ovarian cycle in postpartum dairy cows. Animal Reproduction Science 108: 134-143.

Tenhagen, B. A., A. Helmbold and W. Heuwieser 2007. Effect of various degrees of dystocia in dairy cattle on calf viability, milk production, fertility and culling. Journal of Veterinary Medicine Series A 54: 98-102.

Tenhagen, B. A., C. Vogel, M. Drillich, G. Thiele and W. Heuwieser 2003. Influence of stage of lactation and milk production on conception rates after timed artificial insemination following Ovsynch. Theriogenology 60: 1527-1537.

Thatcher, W. W., T. R. Bilby, J. A. Bartolome, F. Silvestre, C. R. Staples and J. E. P. Santos 2006. Strategies for improving fertility in the modern dairy cow. Theriogenology 65: 30-44.

Thatcher, W. W. and C. J. Wilcox 1973. Postpartum estrus as an indicator of reproductive status in dairy cows. Journal of Dairy Science 56: 608-610.

United Nations, Department Of Economic And Social Affairs, Population Division 2015. World population prospects: The 2015 revision, key findings and advance tables.

van Amburgh, M. E., D. M. Galton, D. E. Bauman and R. W. Everett 1997. Management and economics of extended calving intervals with use of bovine somatotropin. Livestock Production Science 50: 15-28.

Walker, W. L., R. L. Nebel and M. L. McGilliard 1996. Time of ovulation relative to mounting activity in dairy cattle. Journal of Dairy Science 79: 1555-1561.

Walton, J. S., L. P. Veenhuizen and G. J. King 1987. Relationships between time of day, estrous behavior, and the preovulatory luteinizing hormone surge in Holstein cows after treatment with cloprostenol. Journal of Dairy Science 70: 1652-1663.

Washburn, S. P., W. J. Silvia, C. H. Brown, B. T. McDaniel and A. J. McAllister 2002. Trends in reproductive performance in southeastern Holstein and Jersey DHI herds. Journal of Dairy Science 85: 244-251.

Wathes, D. C., M. Fenwick, Z. Cheng, N. Bourne, S. Llewellyn, D. G. Morris, D. Kenny, J. Murphy and R. Fitzpatrick 2007. Influence of negative energy balance on cyclicity and fertility in the high producing dairy cow. Theriogenology 68, Supplement 1: S232-S241.

Weigel, K. A. 2004. Exploring the role of sexed semen in dairy production systems. Journal of Dairy Science 87, Supplement: E120-E130.

Weller, J. I. and Y. Folman 1990. Effects of calf value and reproductive management on optimum days to first breeding. Journal of Dairy Science 73: 1318-1326.

Whitmore, H. L., W. J. Tyler and L. E. Casida 1974. Effects of early postpartum breeding in dairy cattle. Journal of Animal Science 38: 339-346.

Wiltbank, M. C., H. Lopez, R. Sartori, S. Sangsritavong and A. Gümen 2006. Changes in reproductive physiology of lactating dairy cows due to elevated steroid metabolism. Theriogenology 65: 17-29.

Wiltbank, M. C. and J. R. Pursley 2014. The cow as an induced ovulator: Timed AI after synchronization of ovulation. Theriogenology 81: 170-185.

Xu, Z. Z. and L. J. Burton 1999. Effects of oestrus synchronisation and fixed-time artificial insemination on the reproductive performance of dairy heifers. New Zealand Veterinary Journal 47: 101-104.

Young, C., F. A. Di Croce, D. Roper, J. Harris, N. Rohrbach, J. Wilkerson and F. N. Schrick 2010. Effect of reproductive tract size on conception rates in lactating dairy cows utilizing a reproductive tract scoring system. Reproduction, Fertility and Development 23: 119-119.

VITA

Sarah Elizabeth Orr was born on December 13, 1990 in Greeneville, Tennessee. She attended West Greene High School and graduated in the spring of 2009. She furthered her education by attending the University of Tennessee, Knoxville, where she earned a Bachelor's of Science in Animal Science in 2013, and continued on to graduate education under the guidance of Drs. J. Lannett Edwards and F. Neal Schrick. In December of 2015, Sarah graduated with a Master's of Science degree in Animal Science with a concentration in reproductive physiology.