



4-2019

Association of *Ceratobasidium theobromae* and *Lasiodiplodia* species with symptom variation in vascular streak disease of cacao

Johnny F. Balidion
North Carolina State University

Marc A. Cubeta
North Carolina State University

Follow this and additional works at: <https://trace.tennessee.edu/masmc>

Recommended Citation

Balidion, Johnny F. and Cubeta, Marc A., "Association of *Ceratobasidium theobromae* and *Lasiodiplodia* species with symptom variation in vascular streak disease of cacao" (2019). *Middle Atlantic States Mycological Conference 2019*.
<https://trace.tennessee.edu/masmc/11>

This Presentation is brought to you for free and open access by the Conferences at UT at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Middle Atlantic States Mycological Conference 2019 by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Mid-Atlantic States Mycological Conference (MASMC)
University of Tennessee – Knoxville
12-14 April 2019

ABSTRACTS - Oral Presentations

Association of *Ceratobasidium theobromae* and *Lasiodiplodia* species with symptom variation in vascular streak disease of cacao

Johnny F. Balidion, Marc A. Cubeta

Department of Entomology and Plant Pathology, North Carolina State University

Vascular streak dieback (VSD) is an endemic disease of cacao in Southeast Asia reported to be caused by *Ceratobasidium theobromae*. A survey was conducted in 2016-17 from 20 cacao-growing provinces in the Philippines to characterize VSD symptom expression. Fifty three percent of the leaves and petioles sampled (620/1170) exhibited marginal necrosis and chlorosis with epidermal cracks on the petiole or leaf midrib. Approximately 10% of the samples (115/1170) exhibited characteristic chlorotic spotting symptoms. Chlorosis with random necrotic spots and a chlorotic halo was observed in 19% of the sample, while varying mixtures of necrosis and chlorosis on the leaf represented approximately 18% of the samples collected. To validate the presence of fungi associated with the predominant symptom, a PCR-based detection assay using ITS species-specific primers was performed on 30 petiole samples exhibiting leaf symptoms and revealed 100% and 73% detection frequency for *Ceratobasidium theobromae* (Ct) and *Lasiodiplodia* spp., respectively. To examine localization of Ct and *Lasiodiplodia* spp. within a branch, we examined sections of 25 intact twigs grouped per symptom type to determine the frequency of occurrence. Approximately 22% of the segments sampled across symptom types were positive for Ct, 76.1% for *Lasiodiplodia* spp., including in asymptomatic samples, and 16.9% for both species. Morphological and sequence analysis of ITS, *tef1- α* , *tub2*, *rpb2*, and *cmdA* for 10 randomly selected fungal isolates sampled from VSD infected plants were identified as *L. theobromae* and *L. brasiliensis*. Our results suggest the possible role of *Lasiodiplodia* species in the observed variation in VSD symptoms.