



4-26-2018

Biodegradable Plastic Mulch Films: Impacts on Soil Microbial Communities and Ecosystem Functions

Sreejata Bandopadhyay
University of Tennessee, Knoxville

Lluís Martin-Closas
University of Lleida, Spain

Ana M. Pelacho
University of Lleida, Spain

Jennifer M. DeBruyn
University of Tennessee, Knoxville, jdebruyn@utk.edu

Follow this and additional works at: http://trace.tennessee.edu/utk_biospubs

Recommended Citation

Bandopadhyay, Sreejata, Lluís Martin-Closas, Ana M. Pelacho, and Jennifer M. DeBruyn. "Biodegradable Plastic Mulch Films: Impacts on Soil Microbial Communities and Ecosystem Functions." *Frontiers in Microbiology* 9 (2018). <https://doi.org/10.3389/fmicb.2018.00819>.

This Article is brought to you for free and open access by the Biosystems Engineering and Soil Science at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Biosystems Engineering and Soil Science Publications and Other Works by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.



Biodegradable Plastic Mulch Films: Impacts on Soil Microbial Communities and Ecosystem Functions

Sreejata Bandopadhyay¹, Lluís Martín-Closas², Ana M. Pelacho² and Jennifer M. DeBruyn^{1*}

¹ Department of Biosystems Engineering & Soil Science, The University of Tennessee, Knoxville, Knoxville, TN, United States,

² Department of Horticulture, Botany and Gardening, School of Agrifood and Forestry Science and Engineering, University of Lleida, Lleida, Spain

OPEN ACCESS

Edited by:

Paolina Garbeva,
Netherlands Institute of Ecology
(NIOO-KNAW), Netherlands

Reviewed by:

Richard Allen White III,
RAW Molecular Systems (RMS) LLC,
United States

Jim He,

University of Chinese Academy
of Sciences (UCAS), China
Esperanza Huerta Lwanga,
El Colegio de la Frontera Sur, Mexico

*Correspondence:

Jennifer M. DeBruyn
jdebruyn@utk.edu

Specialty section:

This article was submitted to
Terrestrial Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 10 January 2018

Accepted: 10 April 2018

Published: 26 April 2018

Citation:

Bandopadhyay S, Martín-Closas L,
Pelacho AM and DeBruyn JM (2018)
Biodegradable Plastic Mulch Films:
Impacts on Soil Microbial
Communities and Ecosystem
Functions. *Front. Microbiol.* 9:819.
doi: 10.3389/fmicb.2018.00819

Agricultural plastic mulch films are widely used in specialty crop production systems because of their agronomic benefits. Biodegradable plastic mulches (BDMs) offer an environmentally sustainable alternative to conventional polyethylene (PE) mulch. Unlike PE films, which need to be removed after use, BDMs are tilled into soil where they are expected to biodegrade. However, there remains considerable uncertainty about long-term impacts of BDM incorporation on soil ecosystems. BDMs potentially influence soil microbial communities in two ways: first, as a surface barrier prior to soil incorporation, indirectly affecting soil microclimate and atmosphere (similar to PE films) and second, after soil incorporation, as a direct input of physical fragments, which add carbon, microorganisms, additives, and adherent chemicals. This review summarizes the current literature on impacts of plastic mulches on soil biological and biogeochemical processes, with a special emphasis on BDMs. The combined findings indicated that when used as a surface barrier, plastic mulches altered soil microbial community composition and functioning via microclimate modification, though the nature of these alterations varied between studies. In addition, BDM incorporation into soil can result in enhanced microbial activity and enrichment of fungal taxa. This suggests that despite the fact that total carbon input from BDMs is minuscule, a stimulatory effect on microbial activity may ultimately affect soil organic matter dynamics. To address the current knowledge gaps, long term studies and a better understanding of impacts of BDMs on nutrient biogeochemistry are needed. These are critical to evaluating BDMs as they relate to soil health and agroecosystem sustainability.

Keywords: biodegradable plastic, plastic mulch, polyethylene, specialty crops, soil microbiology, soil microclimate, soil biogeochemistry, soil health

INTRODUCTION: AGRICULTURAL PLASTIC MULCH FILMS

Agricultural plastic mulch films are used in production of specialty crops to modify soil temperatures, conserve soil moisture (Kader et al., 2017) and reduce weed pressure (Martín-Closas et al., 2017), ultimately improving crop productivity. Low-density polyethylene (PE) is the most commonly used plastic mulch because it is inexpensive, easily processed, highly durable and flexible (Kasirajan and Ngouajio, 2012). However, widespread use of PE, which is not biodegradable, has resulted in serious environmental contamination (Teuten et al., 2009; Liu E.K. et al., 2014; He et al., 2015; Steinmetz et al., 2016).

A growing concern is that plastic mulches are never completely removed from a field, leaving remnants which remain in soil for decades (Feuilloley et al., 2005; Kyrikou and Briassoulis, 2007; Briassoulis et al., 2015; Ramos et al., 2015). In China, long term use of plastic film mulches has resulted in an estimated accumulation of 50–260 kg hm⁻² of residual plastics in topsoil (0–20 cm), which can inhibit plant growth (Liu E.K. et al., 2014). While PE is considered to be chemically inert, accumulated PE fragments can affect soil physically and may enter the food chain (Barnes et al., 2009; Teuten et al., 2009; Sivan, 2011; Rillig, 2012; Duis and Coors, 2016; Huerta Lwanga et al., 2016). Plastic mulches also introduce various additives such as plasticizing agents which may pollute soil (Van Wezel et al., 2000; Fu and Du, 2011; Kong et al., 2012; Magdouli et al., 2013; Wang et al., 2013, 2015).

Biodegradable plastic mulches (BDMs) have been developed as substitutes to PE mulch films and are designed to be tilled into soil after use where resident microorganisms degrade the plastic. BDMs can be prepared from biobased polymers derived from microbes or plants, or fossil-sourced materials (Marechal, 2003). Common biobased polymers used in BDMs include polylactic acid (PLA), starch, cellulose, and polyhydroxyalkanoates (PHA). Fossil-sourced polyesters used in BDMs include poly(butylene succinate) (PBS), poly(butylene succinate-co-adipate) (PBSA), and poly(butylene-adipate-co-terephthalate) (PBAT) (Kasirajan and Ngouajio, 2012). Polymers used in BDMs contain ester bonds or are polysaccharides, which are amenable to microbial hydrolysis (Brodhagen et al., 2015). In theory, BDMs should be completely catabolized by soil microorganisms, converted to microbial biomass, CO₂ and water (Malinconico et al., 2002; Feuilloley et al., 2005; Imam et al., 2005; Dintcheva and La Mantia, 2007; Kyrikou and Briassoulis, 2007; Kijchavengkul et al., 2008; Lucas et al., 2008). In practice, complete breakdown in a reasonable amount of time is not always observed (Li et al., 2014b). Regulators and growers cite concerns about unpredictable or incomplete breakdown and the ultimate fate of BDM constituents and their effect on soil ecosystems (Goldberger et al., 2015; Miles et al., 2017). Due to increased demand for eco-friendly substitutes to PE, the global market for BDMs is expected to continue to grow. Soil health is a key component of agroecosystem sustainability, thus there is a need to understand the effects of BDMs on both crop productivity and soils. To date, the majority of soil studies related to plastic mulching have focused on PE. The objective of this review is to highlight research concerning impacts of plastic mulches on soil microbial communities and their processes with an emphasis on BDMs. Gaps in our current understanding of how plastics affect soil ecosystems are highlighted.


INDIRECT EFFECTS OF PLASTIC MULCHES ON SOILS VIA MICROCLIMATE MODIFICATION

One way that plastic mulches (both BDMs and PE) may indirectly affect soil ecosystems and microbial community functioning is via modification of soil microclimate and atmosphere. As a

barrier on the soil surface, plastic mulches reduce evaporation and gas exchange, increase temperature and reduce light transmissivity (**Figure 1**; Kasirajan and Ngouajio, 2012). The extent of these modifications depends on their physicochemical properties; for example, PE mulches result in greater warming compared to BDMs (Moreno and Moreno, 2008; Kader et al., 2017) and are less vapor-permeable (Touchaleaume et al., 2016) resulting in accumulation of soil CO₂ (Zhang et al., 2015; Yu et al., 2016). By serving as a barrier to evaporation, plastic mulches can result in increased soil moisture levels (Qin et al., 2015) which can ultimately alter soil physical structure; for example by increasing the proportion of water stable aggregates (Siwek et al., 2015). Favorable moisture and temperature conditions under plastic mulches also affect plant roots, typically stimulating root development and increasing root exudation (Li et al., 2004b; Subrahmaniyan et al., 2006; Wang et al., 2016). This results in greater nutrient availability for rhizosphere microorganisms (Subrahmaniyan et al., 2006; Lin et al., 2008; Maul et al., 2014; Liu et al., 2015).

Since levels of soil moisture, temperature, vapor diffusivity and presence of roots modulate microbial activity, it follows that modifications to soil microclimate under plastic mulches affect soil microbial communities. Plastic mulching can also decrease populations of soil invertebrates (Schonbeck and Evanylo, 1998; Miñarro and Dapena, 2003), which may reduce top-down grazing pressures on soil microbes. A 28-year study in Shenyang, China, demonstrated that plastic film mulching increased relative abundances of *Proteobacteria* and *Actinobacteria* (Farmer et al., 2017). Other studies reported improved control of *Phytophthora capsici* (Núñez-Zofío et al., 2011) or increased mycotoxigenic fungi under plastic mulches (Munoz et al., 2015). From PE studies, we can infer that BDMs may have similar indirect effects and alter microbial community structure and diversity.

In addition to changes in microbial community structure, plastic film mulches affect microbial functioning. Some studies report increased microbial activity under mulches (Mu et al., 2014, 2016; Zhang et al., 2015; Chen H. et al., 2017), while others report decreased activity (Moreno and Moreno, 2008). The response is most likely dependent on the amount of warming under the mulches: where ambient temperatures are cool, mulches bring soil temperature closer to microbial optima and increase activity, whereas in warmer seasons, the mulches may push temperatures above optima, limiting soil microbial activity (Moreno and Moreno, 2008). The changes in microbial activity ultimately influence nutrient cycling and storage. The effect of plastic on soil organic carbon (SOC) is the result of the balance between increased root growth and exudate secretion, and microbial decomposition and loss to CO₂ (Wien et al., 1993; Nan et al., 2016). Thus, it is not surprising that studies examining SOC under plastic mulches have yielded mixed results, with some reporting increased microbial biomass carbon (Li et al., 2004a; An et al., 2015) and SOC (Munoz et al., 2017) and others no change (Wang et al., 2016) or decreased SOC (Cuello et al., 2015). It should be noted that changes in SOC take place over longer time intervals, so the short term (one or two seasons)



nature of most mulching experiments do not capture these longer term dynamics. Plastic mulching also affects cycling and losses of nitrogen in soils (Qin et al., 2015; Nan et al., 2016). Because plastic mulching improves water use efficiency (WUE), nitrate leaching is reduced (Romic et al., 2003). Indeed, Qin et al. (2015) estimated up to 60% increase in nitrogen use efficiency (NUE) under PE mulching compared to no-mulch controls. With respect to N₂O gas release, results are mixed, with some studies reporting decreases (Berger et al., 2013; Li et al., 2014d; Liu J.L. et al., 2014) and others, increases (Okuda et al., 2007; Arriaga et al., 2011; Nishimura et al., 2012; Cuello et al., 2015; Chen H. et al., 2017).

Together, these studies show that plastic mulching, independent of composition, has significant effects on soil microbes and their processes via environmental modification. In several cases, improved crop productivity with mulch was accompanied by a loss of soil organic matter and increased release of greenhouse gasses (Steinmetz et al., 2016). It is important to note that PE films often result in higher soil temperatures and are more effective in suppressing weeds compared to BDMs (Bonanomi et al., 2008). As a physical barrier, BDMs are expected to have similar, though not identical, indirect effects on soil microbes via microclimate modification; the outstanding question is how these effects play out when direct incorporation and biodegradation of BDMs are taken into consideration.

DIRECT EFFECTS OF BDMs VIA INCORPORATION INTO SOIL

While BDMs may have comparable effects as PE mulches when used as a surface barrier, they are distinctly different when considering their ultimate fate. After the growing season, PE films should be removed from the soil surface, while BDMs are meant to be tilled in and biodegraded by microorganisms. BDM fragments are both a physical and a biogeochemical input

(Figure 1). This aspect is unique to BDMs, and may have effects on soil ecology and functioning that cannot be predicted from studies of non-biodegradable plastics such as PE.

Biodegradable plastic mulch fragments may physically modify soil before they are fully biodegraded. For example, PE plastic fragments reduce soil infiltration and water absorption; their accumulation may affect soil ecosystems and ultimately plant germination and growth (Liu E.K. et al., 2014). Therefore, it is conceivable that under conditions restricting soil microbiological activity (e.g., water scarcity), BDM fragments may accumulate in soil with similar effects on soil and plants.

From a toxicology standpoint, the fragments of BDMs incorporated into the soil are generally considered to be safe. For example, tests of the starch-copolyester blend Mater-Bi® (Novamont, Novara, Italy) have shown no ecotoxic effects (Sforzini et al., 2016), nor adverse effect on nitrification potential (ISO 14238:2012) (Ardisson et al., 2014), *Enchytraeus albidus* reproduction (ISO/CD 16387), or *Vibrio fischeri* (ISO 11348 flash test) (Kapanen et al., 2008). Similarly, soil samples containing Ecoflex® (BASF), PHB, and PLA show no demonstrated visual phytotoxicity (ISO 11269-2) (Rychter et al., 2006, 2010). It should be noted that these studies focus on acute responses; possible effects of longer exposure is untested.

Plastic mulches are composed not only of the main polymers but also of small amounts of organic (e.g., additives, plasticizers, etc.) and inorganic (e.g., Cu, Ni, etc.) components, whose effects are largely unknown. Traditional plant tests for toxicity have not been adapted to identify effects of compounds released from BDMs. First, different compounds are released at different times during the biodegradation process. Second, frequently used tests fail to reckon the changing needs and responses throughout plant development by only focusing on germination. Finally, the diversity of plant responses in the ecosystem is narrowly represented by tests that analyze early growth in a few, mostly vigorous, plant species. Despite these constraints, some effects have emerged. A phytotoxicity test of several

chemicals used in bioplastics found that some exhibited a concentration-dependent inhibition of plant growth (Martin-Closas et al., 2014). Acrylate polymers used to maintain soil humidity damaged maize root and shoot development (Chen et al., 2016). Organic compounds released from mulch polymers have been found to be absorbed by crop plants (Du et al., 2009; Li et al., 2014c; Chen N. et al., 2017). Given some of the demonstrated effects on plants, these additives may also impact soil microbes and their functions, though these effects are largely unexplored.

Tilled into soil, BDMs are an input of carbon, albeit a very small one when taking into account the volume of soil into which they are incorporated. However, the growth of soil microbes in agricultural soil is usually carbon-limited and several studies have demonstrated responses by soil microbes to these small inputs. BDMs have caused increases in microbial biomass and enzyme activities (Li et al., 2014a; Yamamoto-Tamura et al., 2015) and changes in soil microbial community structures (Koitabashi et al., 2012; Li et al., 2014b; Muroi et al., 2016). There is evidence that BDMs enrich for certain taxa, for example, PBSA films preferentially selected for *Aspergillus*, *Penicillium*, and *Acanthamoeba* fungi (Koitabashi et al., 2012) and PBAT film surfaces were enriched in Ascomycota (*Apodus*, *Saccharicola*, *Setophoma*), and Proteobacteria (*Hyphomicrobium*, *Caenimonas*) (Muroi et al., 2016). Several studies have also noted increased fungal abundances in soil as a result of BDM incorporation (Rychter et al., 2006; Li et al., 2014b; Ma et al., 2016; Muroi et al., 2016). The majority of these studies examine only one soil type or location; one of the few studies to examine responses in multiple locations showed an enrichment of fungi in one location and Gram-positive bacteria in another (Li et al., 2014b) indicating that microbial responses to BDMs may be affected by environment, soil type and/or management legacies.

In order to tease apart whether observed changes in microbial communities are a result of microclimate effects (i.e., changes that would be expected regardless of the plastic material used) or are specific to BDMs tilled into soil, results from studies that directly compare microbial communities under PE and BDMs in the same experiment are required. The few studies available reported increased microbial abundances, respiration, and enzyme activities under BDMs compared to PE treatments (Moreno and Moreno, 2008; Li et al., 2014a; Yamamoto-Tamura et al., 2015; Barragán et al., 2016; Hajighasemi et al., 2016; Ma et al., 2016) suggesting that incorporation of BDMs does have some effect on microbial activity. Evidence of enhanced degradative activities by soil microbes suggests that BDMs may ultimately change carbon cycling and storage in soil. The total amount of carbon in BDMs is small, and much of it is expected to be respired as CO₂. However, repeated tilling of BDMs into soil may have an effect over time. In one study, use of BDMs resulted in increased microbial biomass carbon compared to PE mulches (Moreno and Moreno, 2008), suggesting an impact on soil carbon dynamics that may accumulate over time. It should also be considered whether enhanced BDM decomposition would impact cycling of other nutrients. Studies on nutrient transformation related to BDM use are limited; two

studies reported that BDMs, like PE films, had no measurable impact on nitrification potential of soils (Kapanen et al., 2008; Ardisson et al., 2014); effects on other nutrients remain unknown.

Taken together, the changes in microbial community structures, stimulated microbial decomposition, and increased microbial biomass suggest enhanced nutrient and carbon cycling under BDMs, which may result in long term effects on soil organic matter dynamics. However, with limited research on long term studies, it remains unknown if BDMs may impact soil functions differently than PE and what implications this has for sustainability of this technology for crop production.

FUTURE RESEARCH OPPORTUNITIES

Biodegradable plastic mulches are a promising alternative to PE plastic film mulches. However, there are considerable gaps in our understanding of how long-term use of BDMs affects soil ecosystems that are critical to crop productivity. Effects of conventional PE mulches on soil microclimate, microbial communities and biogeochemistry provide insight into how BDMs may be indirectly influencing soil. As a surface barrier, plastic mulches can alter soil microbial community composition and functioning in terms of carbon and nitrogen cycling via microclimate modification, though the nature of these alterations has varied between studies. Additionally, there is a lack of knowledge regarding the ecological consequences of BDM degradation products (Lambert and Wagner, 2017). Repeated tilling of BDM fragments into soil may alter the soil physical environment and act as a new source of carbon for microbes. In this regard, effects of BDMs on soils are unique compared to other plastics. The dearth of research directly comparing BDMs to PE renders it difficult to tease apart whether BDMs have an impact on soil microbes and their activities above and beyond what would be expected from a PE plastic film. The few available comparative studies show that microbial activity is enhanced under BDMs. This suggests that despite the fact that total carbon input from BDMs is minuscule, a stimulatory effect on microbial activity may contribute to soil microbial biomass and ultimately soil organic matter.

Several key gaps remain in our understanding of BDMs and their impacts on soil ecosystems. First, studies to date have focused on short term effects, generally one or two growing seasons, or acute toxicity, so long term effects are unknown. Second, the relationship between plastic composition and microbial responses needs exploration: different types of biodegradable plastics will likely differentially affect soil microbes, based on both the parent polymer composition and breakdown products. Third, additives have been demonstrated to leach out of plastic and affect plants; but their effects on soil microbes are unknown. Fourth, several studies have indicated that BDMs may stimulate decomposition; however, effects on nutrient biogeochemistry are largely unexplored. To address these knowledge gaps, long term studies are needed to assess soil health and sustainability impacts, particularly with respect to soil

carbon and/or chronic toxicity effects. In addition, studies should include a direct comparison of PE to BDMs to determine whether BDMs affect soils differently than conventional plastic mulches. Addressing these knowledge gaps will provide much-needed information to growers and regulators on the safety and sustainability of BDMs for agroecosystems.

AUTHOR CONTRIBUTIONS

SB and JD conceived of the review topic and were responsible for final editing. SB, LM-C, AP, and JD all wrote portions of this review.

REFERENCES

- An, T., Schaeffer, S., Li, S., Fu, S., Pei, J., Li, H., et al. (2015). Carbon fluxes from plants to soil and dynamics of microbial immobilization under plastic film mulching and fertilizer application using 13 C pulse-labeling. *Soil Biol. Biochem.* 80, 53–61. doi: 10.1016/j.soilbio.2014.09.024
- Ardissone, G. B., Tosin, M., Barbale, M., and Degli-Innocenti, F. (2014). Biodegradation of plastics in soil and effects on nitrification activity. A laboratory approach. *Front. Microbiol.* 5:710. doi: 10.3389/fmicb.2014.00710
- Arriaga, H., Nunez-Zofio, M., Larregla, S., and Merino, P. (2011). Gaseous emissions from soil biodesinfestation by animal manure on a greenhouse pepper crop. *Crop Prot.* 30, 412–419. doi: 10.1016/j.cropro.2010.12.012
- Barnes, D. K., Galgani, F., Thompson, R. C., and Barlaz, M. (2009). Accumulation and fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364, 1985–1998. doi: 10.1098/rstb.2008.0205
- Barragán, D., Pelacho, A., and Martín-Closas, L. (2016). Degradation of agricultural biodegradable plastics in the soil under laboratory conditions. *Soil Res.* 54, 216–224. doi: 10.1071/SR15034
- Berger, S., Kim, Y., Kettering, J., and Gebauer, G. (2013). Plastic mulching in agriculture-Friend or foe of N₂O emissions? *Agric. Ecosyst. Environ.* 167, 43–51. doi: 10.1016/j.agee.2013.01.010
- Bonanomi, G., Chiuazzini, M., Caporaso, S., Del Sorbo, G., Moschetti, G., and Felice, S. (2008). Soil solarization with biodegradable materials and its impact on soil microbial communities. *Soil Biol. Biochem.* 40, 1989–1998. doi: 10.1016/j.soilbio.2008.02.009
- Briassoulis, D., Babou, E., Hiskakis, M., and Kyriou, I. (2015). Analysis of long-term degradation behaviour of polyethylene mulching films with pro-oxidants under real cultivation and soil burial conditions. *Environ. Sci. Pollut. Res.* 22, 2584–2598. doi: 10.1007/s11356-014-3464-9
- Brodhagen, M., Peyron, M., Miles, C., and Inglis, D. A. (2015). Biodegradable plastic agricultural mulches and key features of microbial degradation. *Appl. Microbiol. Biotechnol.* 99, 1039–1056. doi: 10.1007/s00253-014-6267-5
- Chen, H., Liu, J., Zhang, A., Chen, J., Cheng, G., Sun, B., et al. (2017). Effects of straw and plastic film mulching on greenhouse gas emissions in Loess Plateau, China: a field study of 2 consecutive wheat-maize rotation cycles. *Sci. Total Environ.* 579, 814–824. doi: 10.1016/j.scitotenv.2016.11.022
- Chen, N., Shuai, W., Hao, X., Zhang, H., Zhou, D., and Gao, J. (2017). Contamination of phthalate esters in vegetable agriculture and human cumulative risk assessment. *Pedosphere* 27, 439–451. doi: 10.1016/S1002-0160(17)60340-0
- Chen, X., Mao, X., Lu, Q., Liao, Z., and He, Z. (2016). Characteristics and mechanisms of acrylate polymer damage to maize seedlings. *Ecotoxicol. Environ. Saf.* 129(Suppl. C), 228–234. doi: 10.1016/j.ecoenv.2016.03.018
- Cuello, J. P., Hwang, H. Y., Gutierrez, J., Kim, S. Y., and Kim, P. J. (2015). Impact of plastic film mulching on increasing greenhouse gas emissions in temperate upland soil during maize cultivation. *Appl. Soil Ecol.* 91, 48–57. doi: 10.1016/j.apsoil.2015.02.007
- Dintcheva, N. T., and La Mantia, F. P. (2007). Durability of a starch-based biodegradable polymer. *Polym. Degrad. Stab.* 92, 630–634. doi: 10.1016/j.polymdegradstab.2007.01.003

FUNDING

This work was supported by the United States Department of Agriculture (Award 2014-51181-22382 to JD) and the Spain Ministry of Education and Science (Ref. AGL2008-03733 to LM-C and AP).

ACKNOWLEDGMENTS

We are grateful to D. Hayes, S. Guerrini, D. Martens, and L. Tymon for providing valuable critical feedback on a draft of this manuscript.

- Du, Q. Z., Fu, X. W., and Xia, H. L. (2009). Uptake of di-(2-ethylhexyl)phthalate from plastic mulch film by vegetable plants. *Food Addit. Contam. Part A* 26, 1325–1329. doi: 10.1080/02652030903081952
- Duis, K., and Coors, A. (2016). Microplastics in the aquatic and terrestrial environment: sources (with a specific focus on personal care products), fate and effects. *Environ. Sci. Eur.* 28:2. doi: 10.1186/s12302-015-0069-y
- Farmer, J., Zhang, B., Jin, X. X., Zhang, P., and Wang, J. K. (2017). Long-term effect of plastic film mulching and fertilization on bacterial communities in a brown soil revealed by high through-put sequencing. *Arch. Agron. Soil Sci.* 63, 230–241. doi: 10.1080/03650340.2016.1193667
- Feuilletoy, P., Cesar, G., Benguigui, L., Grohens, Y., Pillin, I., Bewa, H., et al. (2005). Degradation of polyethylene designed for agricultural purposes. *J. Polym. Environ.* 13, 349–355. doi: 10.1007/s10924-005-5529-9
- Fu, X., and Du, Q. (2011). Uptake of di-(2-ethylhexyl) phthalate of vegetables from plastic film greenhouses. *J. Agric. Food Chem.* 59, 11585–11588. doi: 10.1021/jf203502e
- Goldberger, J. R., Jones, R. E., Miles, C. A., Wallace, R. W., and Inglis, D. A. (2015). Barriers and bridges to the adoption of biodegradable plastic mulches for US specialty crop production. *Renew. Agric. Food Syst.* 30, 143–153. doi: 10.1017/S1742170513000276
- Hajjighasemi, M., Nocek, B. P., Tchigvintsev, A., Brown, G., Flick, R., Xu, X., et al. (2016). Biochemical and structural insights into enzymatic depolymerization of polylactic acid and other polyesters by microbial carboxylesterases. *Biomacromolecules* 17, 2027–2039. doi: 10.1021/acs.biomac.6b00223
- He, L., Gielen, G., Bolan, N. S., Zhang, X., Qin, H., Huang, H., et al. (2015). Contamination and remediation of phthalic acid esters in agricultural soils in China: a review. *Agron. Sustain. Dev.* 35, 519–534. doi: 10.1007/s13593-014-0270-1
- Huerta Lwanga, E., Gertsen, H., Gooren, H., Peters, P., Salánki, T., van der Ploeg, M., et al. (2016). Microplastics in the terrestrial ecosystem: implications for *Lumbricus terrestris* (Oligochaeta, Lumbricidae). *Environ. Sci. Technol.* 50, 2685–2691. doi: 10.1021/acs.est.5b05478
- Imam, S. H., Cinelli, P., Gordon, S. H., and Chiellini, E. (2005). Characterization of biodegradable composite films prepared from blends of poly(vinyl alcohol), cornstarch, and lignocellulosic fiber. *J. Polym. Environ.* 13, 47–55. doi: 10.1007/s10924-004-1215-6
- Kader, M. A., Senge, M., Mojid, M. A., and Ito, K. (2017). Recent advances in mulching materials and methods for modifying soil environment. *Soil Tillage Res.* 168, 155–166. doi: 10.1016/j.still.2017.01.001
- Kapanen, A., Schettini, E., Vox, G., and Itävaara, M. (2008). Performance and environmental impact of biodegradable films in agriculture: a field study on protected cultivation. *J. Polym. Environ.* 16, 109–122. doi: 10.1007/s10924-008-0091-x
- Kasirajan, S., and Ngouajio, M. (2012). Polyethylene and biodegradable mulches for agricultural applications: a review. *Agron. Sustain. Dev.* 32, 501–529. doi: 10.1007/s13593-011-0068-3
- Kijchavengkul, T., Auras, R., Rubino, M., Ngouajio, M., and Fernandez, R. T. (2008). Assessment of aliphatic-aromatic copolyester biodegradable mulch films. Part I: field study. *Chemosphere* 71, 942–953. doi: 10.1016/j.chemosphere.2007.10.074

- Koitaishi, M., Noguchi, M. T., Sameshima-Yamashita, Y., Hiradate, S., Suzuki, K., Yoshida, S., et al. (2012). Degradation of biodegradable plastic mulch films in soil environment by phylloplane fungi isolated from gramineous plants. *AMB Express* 2:40. doi: 10.1186/2191-0855-2-40
- Kong, S., Ji, Y., Liu, L., Chen, L., Zhao, X., Wang, J., et al. (2012). Diversities of phthalate esters in suburban agricultural soils and wasteland soil appeared with urbanization in China. *Environ. Pollut.* 170, 161–168. doi: 10.1016/j.envpol.2012.06.017
- Kyrrikou, I., and Briassoulis, D. (2007). Biodegradation of agricultural plastic films: a critical review. *J. Polym. Environ.* 15, 125–150. doi: 10.1007/s10924-007-0053-8
- Lambert, S., and Wagner, M. (2017). Environmental performance of bio-based and biodegradable plastics: the road ahead. *Chem. Soc. Rev.* 46, 6855–6871. doi: 10.1039/c7cs00149e
- Li, C., Moore-Kucera, J., Lee, J., Corbin, A., Brodhagen, M., Miles, C., et al. (2014a). Effects of biodegradable mulch on soil quality. *Appl. Soil Ecol.* 79, 59–69. doi: 10.1016/j.apsoil.2014.02.012
- Li, C., Moore-Kucera, J., Miles, C., Leonas, K., Lee, J., Corbin, A., et al. (2014b). Degradation of potentially biodegradable plastic mulch films at three diverse US locations. *Agroecol. Sustain. Food Syst.* 38, 861–889. doi: 10.1080/21683565.2014.884515
- Li, F.-M., Song, Q. H., Jjemba, P. K., and Shi, Y.-C. (2004a). Dynamics of soil microbial biomass C and soil fertility in cropland mulched with plastic film in a semiarid agro-ecosystem. *Soil Biol. Biochem.* 36, 1893–1902. doi: 10.1016/j.soilbio.2004.04.040
- Li, F.-M., Wang, J., Xu, J.-Z., and Xu, H.-L. (2004b). Productivity and soil response to plastic film mulching durations for spring wheat on entisols in the semiarid Loess Plateau of China. *Soil Tillage Res.* 78, 9–20. doi: 10.1016/j.still.2003.12.009
- Li, Y. W., Cai, Q. Y., Mo, C. H., Zeng, Q. Y., Lü, H., Li, Q. S., et al. (2014c). Plant uptake and enhanced dissipation of di(2-ethylhexyl) phthalate (DEHP) in spiked soils by different plant species. *Int. J. Phytoremediation* 16, 609–620.
- Li, Z., Zhang, R., Wang, X., Chen, F., Lai, D., and Tian, C. (2014d). Effects of plastic film mulching with drip irrigation on N₂O and CH₄ emissions from cotton fields in arid land. *J. Agric. Sci.* 152, 534–542. doi: 10.1017/S0021859613000701
- Lin, Y.-B., Xue, Q.-H., and Yan, X. (2008). Effect of mulching mode and wheat root on soil microbial flora. *Chin. J. Eco Agric.* 16, 1389–1393. doi: 10.3724/SP.J.1011.2008.01389
- Liu, E. K., He, W. Q., and Yan, C. R. (2014). ‘White revolution’ to ‘white pollution’- agricultural plastic film mulch in China. *Environ. Res. Lett.* 9:3. doi: 10.1088/1748-9326/9/9/091001
- Liu, J., Zhan, A., Chen, H., Luo, S., Bu, L., Chen, X., et al. (2015). Response of nitrogen use efficiency and soil nitrate dynamics to soil mulching in dryland maize (*Zea mays* L.) fields. *Nutr. Cycling Agroecosyst.* 101, 271–283. doi: 10.1007/s10705-015-9678-5
- Liu, J. L., Zhu, L., Luo, S. S., Bu, L. D., Chen, X. P., Yue, S. C., et al. (2014). Response of nitrous oxide emission to soil mulching and nitrogen fertilization in semiarid farmland. *Agric. Ecosyst. Environ.* 188, 20–28. doi: 10.1016/j.agee.2014.02.010
- Lucas, N., Bienaime, C., Bello, C., Queneudec, M., Silvestre, F., and Nava-Saucedo, J.-E. (2008). Polymer biodegradation: mechanisms and estimation techniques—A review. *Chemosphere* 73, 429–442. doi: 10.1016/j.chemosphere.2008.06.064
- Ma, Z. F., Ma, Y. B., Qin, L. Z., Liu, J. X., and Su, H. J. (2016). Preparation and characteristics of biodegradable mulching films based on fermentation industry wastes. *Int. Biodeterior. Biodegradation* 111, 54–61. doi: 10.1016/j.ibiod.2016.04.024
- Magdouli, S., Dagherir, R., Brar, S. K., Drogui, P., and Tyagi, R. D. (2013). Di 2-ethylhexylphthalate in the aquatic and terrestrial environment: a critical review. *J. Environ. Manage.* 127, 36–49. doi: 10.1016/j.jenvman.2013.04.013
- Malinconico, M., Immirzi, B., Massenti, S., La Mantia, F. P., Mormile, P., and Petti, L. (2002). Blends of polyvinylalcohol and functionalised polycaprolactone. A study on the melt extrusion and post-cure of films suitable for protected cultivation. *J. Mater. Sci.* 37, 4973–4978. doi: 10.1023/A:1021058810774
- Marechal, F. (2003). “Biodegradable plastics,” in *Biodegradable Polymers and Plastics*, eds E. Chiellini and R. Solaro (New York, NY: Springer).
- Martin-Closas, L., Botet, R., and Pelacho, A. (2014). An in vitro crop plant ecotoxicity test for agricultural bioplastic constituents. *Polym. Degrad. Stab.* 108, 250–256. doi: 10.1016/j.polymdegradstab.2014.03.037
- Martin-Closas, L., Costa, J., and Pelacho, A. M. (2017). “Agronomic effects of biodegradable films on crop and field environment,” in *Soil Degradable Bioplastics for a Sustainable Modern Agriculture*, ed. M. Malinconico (Berlin: Springer), 67–104. doi: 10.1007/978-3-662-54130-2_4
- Maul, J. E., Buyer, J. S., Lehman, R. M., Culman, S., Blackwood, C. B., Roberts, D. P., et al. (2014). Microbial community structure and abundance in the rhizosphere and bulk soil of a tomato cropping system that includes cover crops. *Appl. Soil Ecol.* 77, 42–50. doi: 10.1016/j.apsoil.2014.01.002
- Miles, C., DeVetter, L., Ghimire, S., and Hayes, D. G. (2017). Suitability of biodegradable plastic mulches for organic and sustainable agricultural production systems. *HortScience* 52, 10–15. doi: 10.21273/HORTSCI11249-16
- Miñarro, M., and Dapena, E. (2003). Effects of groundcover management on ground beetles (Coleoptera: Carabidae) in an apple orchard. *Appl. Soil Ecol.* 23, 111–117. doi: 10.1016/S0929-1393(03)00025-8
- Moreno, M. M., and Moreno, A. (2008). Effect of different biodegradable and polyethylene mulches on soil properties and production in a tomato crop. *Sci. Hortic.* 116, 256–263. doi: 10.1016/j.scienta.2008.01.007
- Mu, L., Fang, L., and Liang, Y. (2016). Temporal and spatial variation of soil respiration under mulching in a greenhouse cucumber cultivation. *Pesqui. Agropecuár. Bras.* 51, 869–879. doi: 10.1590/S0100-204X2016000700010
- Mu, L., Liang, Y., Zhang, C., Wang, K., and Shi, G. (2014). Soil respiration of hot pepper (*Capsicum annuum* L.) under different mulching practices in a greenhouse, including controlling factors in China. *Acta Agric. Scand. B Soil Plant Sci.* 64, 85–95. doi: 10.1080/09064710.2014.887141
- Munoz, K., Buchmann, C., Meyer, M., Schmidt-Heydt, M., Steinmetz, Z., Diehl, D., et al. (2017). Physicochemical and microbial soil quality indicators as affected by the agricultural management system in strawberry cultivation using straw or black polyethylene mulching. *Appl. Soil Ecol.* 113, 36–44. doi: 10.1016/j.apsoil.2017.01.014
- Munoz, K., Schmidt-Heydt, M., Stoll, D., Diehl, D., Ziegler, J., Geisen, R., et al. (2015). Effect of plastic mulching on mycotoxin occurrence and mycobiome abundance in soil samples from asparagus crops. *Mycotoxin Res.* 31, 191–201. doi: 10.1007/s12550-015-0231-9
- Muroi, F., Tachibana, Y., Kobayashi, Y., Sakurai, T., and Kasuya, K. (2016). Influences of poly(butylene adipate-co-terephthalate) on soil microbiota and plant growth. *Polym. Degrad. Stab.* 129, 338–346. doi: 10.1016/j.polymdegradstab.2016.05.018
- Nan, W. G., Yue, S. C., Huang, H. Z., Li, S. Q., and Shen, Y. F. (2016). Effects of plastic film mulching on soil greenhouse gases (CO₂, CH₄ and N₂O) concentration within soil profiles in maize fields on the Loess Plateau, China. *J. Integr. Agric.* 15, 451–464. doi: 10.1016/S2095-3119(15)61106-6
- Nishimura, S., Komada, M., Takebe, M., Yonemura, S., and Kato, N. (2012). Nitrous oxide evolved from soil covered with plastic mulch film in horticultural field. *Biol. Fertil. Soils* 48, 787–795. doi: 10.1007/s00374-012-0672-7
- Núñez-Zofio, M., Larregla, S., and Garbisu, C. (2011). Application of organic amendments followed by soil plastic mulching reduces the incidence of *Phytophthora capsici* in pepper crops under temperate climate. *Crop Prot.* 30, 1563–1572. doi: 10.1016/j.cropro.2011.08.020
- Okuda, H., Noda, K., Sawamoto, T., Tsuruta, H., Hirabayashi, T., Yonemoto, J. Y., et al. (2007). Emission of N₂O and CO₂ and uptake of CH₄ in soil from a satsuma mandarin orchard under mulching cultivation in central Japan. *J. Jpn. Soc. Hortic. Sci.* 76, 279–287. doi: 10.2503/jjshs.76.279
- Qin, W., Hu, C. S., and Oenema, O. (2015). Soil mulching significantly enhances yields and water and nitrogen use efficiencies of maize and wheat: a meta-analysis. *Sci. Rep.* 5:16210. doi: 10.1038/srep16210
- Ramos, L., Berenstein, G., Hughes, E. A., Zalts, A., and Montserrat, J. M. (2015). Polyethylene film incorporation into the horticultural soil of small periurban production units in Argentina. *Sci. Total Environ.* 523, 74–81. doi: 10.1016/j.scitotenv.2015.03.142
- Rillig, M. C. (2012). *Microplastic in Terrestrial Ecosystems and the Soil?* Washington, DC: ACS Publications.
- Romic, D., Romic, M., Borosic, J., and Poljak, M. (2003). Mulching decreases nitrate leaching in bell pepper (*Capsicum annuum* L.) cultivation. *Agric. Water Manag.* 60, 87–97. doi: 10.1016/S0378-3774(02)00168-3

- Rychter, P., Biczak, R., Herman, B., Smylla, A., Kurcok, P., Adamus, G., et al. (2006). Environmental degradation of polyester blends containing atactic poly(3-hydroxybutyrate). Biodegradation in soil and ecotoxicological impact. *Biomacromolecules* 7, 3125–3131. doi: 10.1021/bm060708r
- Rychter, P., Kawalec, M., Sobota, M., Kurcok, P., and Kowalczyk, M. (2010). Study of aliphatic-aromatic copolyester degradation in sandy soil and its ecotoxicological impact. *Biomacromolecules* 11, 839–847. doi: 10.1021/bm901331t
- Schonbeck, M. W., and Evanylo, G. K. (1998). Effects of mulches on soil properties and tomato production II. Plant-available nitrogen, organic matter input, and tillage-related properties. *J. Sustain. Agric.* 13, 83–100. doi: 10.1300/J064v13n01_07
- Sforzini, S., Oliveri, L., Chinaglia, S., and Viarengo, A. (2016). Application of biotests for the determination of soil ecotoxicity after exposure to biodegradable plastics. *Front. Environ. Sci.* 4:68. doi: 10.3389/fenvs.2016.00068
- Sivan, A. (2011). New perspectives in plastic biodegradation. *Curr. Opin. Biotechnol.* 22, 422–426. doi: 10.1016/j.copbio.2011.01.013
- Siwek, P., Domagala-Swiatkiewicz, I., and Kalisz, A. (2015). The influence of degradable polymer mulches on soil properties and cucumber yield. *Agrochimica* 59, 108–123.
- Steinmetz, Z., Wollmann, C., Schaefer, M., Buchmann, C., David, J., Troger, J., et al. (2016). Plastic mulching in agriculture. Trading short-term agronomic benefits for long-term soil degradation? *Sci. Total Environ.* 550, 690–705. doi: 10.1016/j.scitotenv.2016.01.153
- Subrahmaniyan, K., Kalaiselvan, P., Balasubramanian, T., and Zhou, W. (2006). Crop productivity and soil properties as affected by polyethylene film mulch and land configurations in groundnut (*Arachis hypogaea* L.) (Einfluss von polyethylenfilm-mulch und feldbeschaffenheit auf ertrag und bodeneigenschaften im erdnussanbau [*Arachis hypogaea* L.]). *Arch. Agron. Soil Sci.* 52, 79–103. doi: 10.1080/03650340500421786
- Teuten, E. L., Saquing, J. M., Knappe, D. R., Barlaz, M. A., Jonsson, S., Björn, A., et al. (2009). Transport and release of chemicals from plastics to the environment and to wildlife. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364, 2027–2045. doi: 10.1098/rstb.2008.0284
- Touchaleaume, F., Martin-Closas, L., Angellier-Coussy, H., Chevillard, A., Cesar, G., Gontard, N., et al. (2016). Performance and environmental impact of biodegradable polymers as agricultural mulching films. *Chemosphere* 144, 433–439. doi: 10.1016/j.chemosphere.2015.09.006
- Van Wezel, A., Van Vlaardingen, P., Posthumus, R., Crommentuijn, G., and Sijm, D. (2000). Environmental risk limits for two phthalates, with special emphasis on endocrine disruptive properties. *Ecotoxicol. Environ. Saf.* 46, 305–321. doi: 10.1006/eesa.2000.1930
- Wang, J., Chen, G., Christie, P., Zhang, M., Luo, Y., and Teng, Y. (2015). Occurrence and risk assessment of phthalate esters (PAEs) in vegetables and soils of suburban plastic film greenhouses. *Sci. Total Environ.* 523, 129–137. doi: 10.1016/j.scitotenv.2015.02.101
- Wang, J., Luo, Y., Teng, Y., Ma, W., Christie, P., and Li, Z. (2013). Soil contamination by phthalate esters in Chinese intensive vegetable production systems with different modes of use of plastic film. *Environ. Pollut.* 180, 265–273. doi: 10.1016/j.envpol.2013.05.036
- Wang, Y. P., Li, X. G., Fu, T. T., Wang, L., Turner, N. C., Siddique, K. H. M., et al. (2016). Multi-site assessment of the effects of plastic-film mulch on the soil organic carbon balance in semiarid areas of China. *Agric. For. Meteorol.* 228, 42–51. doi: 10.1016/j.agrformet.2016.06.016
- Wien, H. C., Minotti, P. L., and Grubinger, V. P. (1993). Polyethylene mulch stimulates early root growth and nutrient uptake of transplanted tomatoes. *J. Am. Soc. Hortic. Sci.* 118, 207–211.
- Yamamoto-Tamura, K., Hiradate, S., Watanabe, T., Koitabashi, M., Sameshima-Yamashita, Y., Yarimizu, T., et al. (2015). Contribution of soil esterase to biodegradation of aliphatic polyester agricultural mulch film in cultivated soils. *AMB Express* 5:10. doi: 10.1186/s13568-014-0088-x
- Yu, Y., Zhao, C., Stahr, K., Zhao, X., and Jia, H. (2016). Plastic mulching increased soil CO₂ concentration and emissions from an oasis cotton field in Central Asia. *Soil Use Manag.* 32, 230–239. doi: 10.1111/sum.12266
- Zhang, F., Li, M., Qi, J. H., Li, F. M., and Sun, G. J. (2015). Plastic film mulching increases soil respiration in ridge-furrow maize management. *Arid Land Res. Manag.* 29, 432–453. doi: 10.1080/15324982.2015.1018456

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Bandopadhyay, Martin-Closas, Pelacho and DeBruyn. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.