



4-2019

## **Endosymbiotic bacteria of the causative agent of white nose syndrome (*Pseudogymnoascus destructans*) of bats**

Ashton Reece  
*Middle Tennessee State University*

Kylie Bowe  
*Middle Tennessee State University*

Matthew Grisnik  
*Middle Tennessee State University*

Donald Walker  
*Middle Tennessee State University*

Follow this and additional works at: <https://trace.tennessee.edu/masmc>

---

### **Recommended Citation**

Reece, Ashton; Bowe, Kylie; Grisnik, Matthew; and Walker, Donald, "Endosymbiotic bacteria of the causative agent of white nose syndrome (*Pseudogymnoascus destructans*) of bats" (2019). *Middle Atlantic States Mycological Conference 2019*.  
<https://trace.tennessee.edu/masmc/6>

This Poster is brought to you for free and open access by the Conferences at UT at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Middle Atlantic States Mycological Conference 2019 by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

Mid-Atlantic States Mycological Conference (MASMC)  
University of Tennessee – Knoxville  
12-14 April 2019

ABSTRACTS – Posters

**Endosymbiotic bacteria of the causative agent of white nose syndrome (*Pseudogymnoascus destructans*) of bats**

Ashton Reece, Kylie Bowe, Matthew Grisnik, Donald Walker

Biology Department, Toxicology and Disease Group, Middle Tennessee State University

Since its initial isolation in 2008, *Pseudogymnoascus destructans* has been identified as the causative agent of the disease known as White Nose Syndrome. This emerging fungal pathogen has resulted in a drastic decline in many North American bat populations. Extensive research has focused on methods to ameliorate disease-based declines due to the necessary ecological role that bats play in pollination and pest control throughout the country. We have a current understanding of *P. destructans*' interactions with the host and environment, and recent work has shown a correlation between the bat microbiome and susceptibility to infection. Endosymbiotic bacteria are found in most major lineages of plant-associated fungi and have a biological/ecological influence on host phenotypic expression. This study screened for endosymbiotic bacteria within fungal cells of *P. destructans* since these symbionts may play a role in fungal pathogenicity. We isolated *P. destructans* from 28 bat individuals and have detected bacterial DNA in three isolates of *P. destructans*. We generated full length bacterial 16S rRNA sequences for two of three samples that identify as the genus *Nocardia* (Actinobacteria). Future work will use both imaging and high-throughput DNA sequencing to characterize the tentative endosymbiotic bacteria. Bacterial symbionts have thus far only been observed in endophytic fungi of plants, therefore, detection of endosymbiotic bacteria in *P. destructans* is the first wildlife pathogen to show this biological relationship.