



4-2019

## **An enhanced strategy for molecular detection of thousand cankers disease**

Tammy Stackhouse  
*University of Tennessee, Knoxville*

Emel Oren  
*University of Tennessee, Knoxville*

Sarah Boggess  
*University of Tennessee, Knoxville*

Matthew Ginzel  
*Purdue University*

Denita Hadziabdic  
*University of Tennessee, Knoxville*

*See next page for additional authors*

Follow this and additional works at: <https://trace.tennessee.edu/masmc>

---

### **Recommended Citation**

Stackhouse, Tammy; Oren, Emel; Boggess, Sarah; Ginzel, Matthew; Hadziabdic, Denita; Trigiano, Robert; and Klingeman, William, "An enhanced strategy for molecular detection of thousand cankers disease" (2019). *Middle Atlantic States Mycological Conference 2019*.  
<https://trace.tennessee.edu/masmc/5>

This Poster is brought to you for free and open access by the Conferences at UT at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Middle Atlantic States Mycological Conference 2019 by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

---

**Authors**

Tammy Stackhouse, Emel Oren, Sarah Boggess, Matthew Ginzel, Denita Hadziabdic, Robert Trigiano, and William Klingeman

Mid-Atlantic States Mycological Conference (MASMC)  
University of Tennessee – Knoxville  
12-14 April 2019

ABSTRACTS – Posters

**An enhanced strategy for molecular detection of thousand cankers disease**

Tammy Stackhouse<sup>1</sup>, Emel Oren<sup>2,4</sup>, Sarah Boggess<sup>2</sup>, Matthew Ginzel<sup>3</sup>, Denita Hadziabdic<sup>2</sup>, Robert Trigiano<sup>2</sup>, William Klingeman<sup>1</sup>

<sup>1</sup>Department of Plant Sciences, University of Tennessee, <sup>2</sup>Department of Entomology and Plant Pathology, University of Tennessee, <sup>3</sup>Departments of Entomology and Forestry & Natural Resources, Purdue University, <sup>4</sup>Current Address: Diyarbakir Plant Protection Research Institute, Diyarbakir, Turkey

*Juglans nigra* (black walnut) is native throughout the eastern U.S. and used in the western U.S. as an ornamental plant and rootstock. The current estimated value of standing *Juglans nigra* timber exceeds \$539 billion dollars (USD) across its native range. All members of the *Juglans* genus, as well as some *Pterocarya* spp., are threatened by Thousand Cankers Disease (TCD). This disease complex includes the fungal pathogen, *Geosmithia morbida*, the primary insect vector, *Pityophthorus juglandis*, and susceptible host trees. We have used microsatellite loci in a rapid molecular detection protocol that confirms presence of either pathogen or vector DNA in suspect samples. However, the existing protocol requires expensive specialized equipment and technically trained personnel. Our goal was to optimize and simplify the existing molecular detection protocol to enable faster adoption by end-users. The modifications will allow easier visualization of positive samples by utilizing a conventional gel protocol, thus reducing the equipment cost by about \$36,000. Results presented here demonstrate successful amplification of *G. morbida* and *P. juglandis* DNA from drilled bark shaving samples and bark sample archetypes. Assays were also performed to determine the sensitivity of the new conventional gel protocol, yielding detectability for positive identification with very low amounts of target DNA; the equivalent of about 6 conidia or hyphal cells. State and federal regulators and disease diagnosticians will be able to confirm disease presence in suspect samples in less than 12 hours greatly enhancing the ability for preventing further spread of TCD.