



University of Tennessee, Knoxville
**TRACE: Tennessee Research and Creative
Exchange**

Forestry, Trees, and Timber

UT Extension Publications

12-2003

SP290-F Maple Leaf Pouch Galls

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: https://trace.tennessee.edu/utk_agexfores



Part of the [Entomology Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

"SP290-F Maple Leaf Pouch Galls," The University of Tennessee Agricultural Extension Service, 04-0164 SP290F 150 12/03(Rev) E12-4615-00-008-04, https://trace.tennessee.edu/utk_agexfores/27

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Insects, Pests, Diseases & Weeds is brought to you for free and open access by the UT Extension Publications at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Forestry, Trees, and Timber by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Insects

Maple Leaf Pouch Galls

Frank A. Hale, Professor

Originally Developed by Harry E. Williams, Professor Emeritus
Entomology and Plant Pathology

Eriophyid mite species cause a variety of pouch galls on red, sugar and silver maples. Pouch gall formation, a localized growth reaction of the infested plant, occurs as a result of mite feeding activity. The species of mite can be identified by the specific characteristics of the gall on the host plant.

Maple Bladder Gall

Feeding by the maple bladder gall mite, *Vasates quadripedes*, causes a pouch-like growth known as bladder gall on the upper leaf surface of red and silver maples. The galls are variable in shape, rounded or elongated. They are usually crowded and numerous at the base part of the leaf between the larger veins.

The exterior of the galls appears wrinkled and glossy. They change from yellowish green or dark green progressively to pink, brown and finally black. The interior of the gall is hollow. The exit for the new generation of mites is from the underside of the leaf.

Maple Spindle or Finger Gall

The spindle or finger gall mite, *Vasates aceriscrumena*, is common on sugar maples. The galls are elongated, pointed or spindle-shaped. They are variable in size up to 0.2 inch in length and tend to crowd at the tip end of the leaf blade. The galls vary from greenish tinged with yellow to pink to crimson. The interior of the gall is thin-walled and the exit hole is on the underside of the leaf.

Life History of Eriophyid Mites

The mites exit the galls in the fall and migrate to terminal buds of the host tree. The mites enter the buds and overwinter under the bark scales.

In the spring, the mites ride out on the growing leaves. Mite feeding results in the formation of galls. The mites then enter the galls, lay eggs and die. The summer generation of mites occupies the galls during the summer.

Injury to the Maple Trees

Heavy mite infestation and gall formation causes discoloration and distortion. Usually the infestation levels do not cause severe stress in the tree.

Control Measures

In early spring before leaf buds break, spray bark with dormant oil. Some varieties of maples are susceptible to oil injury. Early in the spring when the leaves are about one-fourth expanded, spray with one of the listed insecticide/miticides (Table 1). Mites are exposed on the leaf surface at this time. Cover the upper and lower leaf surfaces with the spray, and repeat the application in 10 days.




Table 1. Chemical Control for Gall Mites on Maple:

Insecticide (Brand Name)	Formulation	100 Gallons*	1 Gallon*
carbaryl (Sevin)	80% S	1¼ lb	1¼ Tbs
dicofol (Kelthane)*	50 WSP	½ - 1 lb (high-volume sprayers) ½ - 1 lb per acre (low-volume sprayers)	— —
insecticidal soap (M-Pede)	49% a.i.	1-2% insecticidal soap in finished spray mixture (i.e., 2 gal soap in 98 gal water)	
(Safer Insecticidal Soap)	49% a.i.	—	5 Tbs
horticultural oil (SunSpray Ultra-Fine Oil)	98.8% paraffinic oil	1-2% oil in finished spray mixture	2.5-5 Tbs

* Amount of formulation/water volume

* Not for use on residential home lawns or residential ornamentals.



Maple bladder gall

This factsheet is available in full color on the University of Tennessee Extension Website at <http://www.utextension.edu/publications/pests/default.asp>.

04-0164 SP290F 150 12/03(Rev) E12-4615-00-008-04

The Agricultural Extension Service offers its programs to all eligible persons regardless of race, color, national origin, sex, age, disability, religion or veteran status and is an Equal Opportunity Employer. COOPERATIVE EXTENSION WORK IN AGRICULTURE AND HOME ECONOMICS

The University of Tennessee Institute of Agriculture, U.S. Department of Agriculture, and county governments cooperating in furtherance of Acts of May 8 and June 30, 1914.

Agricultural Extension Service Charles L. Norman, Dean