



2019

Nutrient Source and Tillage Effects on Maize: II. Yield, Soil Carbon, and Carbon Dioxide Emissions

Deb O'Dell
University of Tennessee, Knoxville

Neal S. Eash
University of Tennessee, Knoxville

Bruce B. Hicks
MetCorps, P.O. Box 1510, Norris, TN 37828

James A. Zahn
DuPont Tate & Lyle Bio Products, LLC, Loudon, TN

Joel N. Oetting
University of Tennessee, Knoxville

See next page for additional authors

Follow this and additional works at: https://trace.tennessee.edu/utk_biospubs

Recommended Citation

O'Dell, Deb; Eash, Neal S.; Hicks, Bruce B.; Zahn, James A.; Oetting, Joel N.; Sauer, Thomas J.; Lambert, Dayton M.; Logan, Joanne; and Goddard, John J., "Nutrient Source and Tillage Effects on Maize: II. Yield, Soil Carbon, and Carbon Dioxide Emissions" (2019). *Biosystems Engineering and Soil Science Publications and Other Works*.
https://trace.tennessee.edu/utk_biospubs/20

This Article is brought to you for free and open access by the Biosystems Engineering and Soil Science at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Biosystems Engineering and Soil Science Publications and Other Works by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Authors

Deb O'Dell, Neal S. Eash, Bruce B. Hicks, James A. Zahn, Joel N. Oetting, Thomas J. Sauer, Dayton M. Lambert, Joanne Logan, and John J. Goddard

Published June 20, 2019

Environment

Nutrient Source and Tillage Effects on Maize: I. Micrometeorological Methods for Measuring Carbon Dioxide Emissions

Deb O'Dell,* Neal S. Eash, Bruce B. Hicks, James A. Zahn, Joel N. Oetting, Thomas J. Sauer, Dayton M. Lambert, Joanne Logan, and John J. Goddard

Core Ideas

- Aerodynamic methods can be used to gap-fill Bowen ratio energy balance micrometeorological measurements.
- Eddy covariance and Bowen ratio energy balance methods agree during turbulent daytime conditions.
- Measuring nighttime net ecosystem exchange is challenging using turbulence-based micrometeorology.

D. O'Dell, N.S. Eash, J.N. Oetting, and J. Logan, Dep. of Biosystems Engineering and Soil Science, Univ. of Tennessee Institute of

ABSTRACT

There is a need to understand the potential benefits of using the biotechnology waste by-product from manufacturing as a fertilizer replacement in agriculture, by quantifying the economic value for the farmer and measuring the environmental impact. Measuring CO₂ emissions can be used to assess environmental impact, including three widely used micrometeorological methodologies: (i) the Bowen Ratio Energy Balance (BREB), (ii) aerodynamic flux-gradient theory, and (iii) eddy covariance (EC). As a first step in quantifying benefits of applying biotechnology waste in agriculture, a detailed examination of these three methods was conducted to understand their effectiveness in quantifying CO₂ emissions for this specific circumstance. The study measured micrometeorological properties over a field planted to maize (*Zea mays* L. var. *indentata*), one plot treated with biotechnology waste applied as a nutrient amendment, and one plot treated with a typical farmer fertilizer practice. Carbon dioxide flux measurements took place over 1 yr, using both BREB and EC systems. The aerodynamic method was used to gap-fill BREB system measurements, and those flux estimates were compared with estimates produced separately by the aerodynamic and EC methods. All methods found greater emissions over the biotechnology waste application. The aerodynamic method CO₂ flux estimates were considerably greater than both the EC and a combined BREB-aerodynamic approach. During the day, the EC and BREB methods agree. At night, the aerodynamic approach detects and accounts for buildup of CO₂ at the surface during