



University of Tennessee, Knoxville
**TRACE: Tennessee Research and Creative
Exchange**

Commercial Horticulture

UT Extension Publications

11-22-2011

SP341-C Insects: Fungus Gnats

Frank A. Hale

Follow this and additional works at: https://trace.tennessee.edu/utk_agexcomhort



Part of the [Agriculture Commons](#)

Recommended Citation

"SP341-C Insects: Fungus Gnats," Frank A. Hale,
SP341-C

, https://trace.tennessee.edu/utk_agexcomhort/35

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Greenhouse Production is brought to you for free and open access by the UT Extension Publications at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Commercial Horticulture by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Insects



Fungus Gnats

Frank A. Hale, Professor

*Originally developed by Harry E. Williams, Professor Emeritus
and Jaime Yanes, Jr., former Assistant Professor
Entomology and Plant Pathology*

The fungus gnat has recently been recognized as an important pest in greenhouses. Although greenhouse plants are commonly attacked, house plants may also become infested. Several genera in the family Sciaridae are of economic concern, such as *Sciara*, *Orfelia*, *Lycoriella* and *Bradysia* species.

Damage

Damage is caused when the larvae, which feed in highly organic soils, infest the roots of African violets, poinsettias, carnations, Easter lilies, geraniums, cyclamens, bedding plants and foliage plants. Symptoms first appear as a loss in plant vigor. As the damage progresses, the plants may fade, begin to wilt suddenly and finally lose foliage.

Description and life cycle

The adult fungus gnat (Fig. A) is about 1/8 inch long and is grayish-black. Its slender body with delicate, long legs and antennae resembles a mosquito. The adult has one pair of clear wings. In the winter and spring, this insect becomes most abundant in greenhouses. Fungus gnats complete development in moist, shady areas in decaying organic matter, such as leaf litter. The adults, which live for seven to 10 days, may deposit up to 150 eggs on the moist soil surface or adjacent to the plant stem. Eggs, laid in strings of three to 40, begin to hatch in four days.

The larvae or maggots (Fig. B) have shiny black head capsules, thread-like white bodies and are up to 1/4 inch in length. Larvae feed primarily on fungi and decaying organic matter. After their normal food



Fig. A: Adult fungus gnat



Fig. B: Fungus gnat larva

supply is exhausted and/or populations become very high, larvae may begin to feed on fine root hairs or other tender tissues. Chewed roots and underground parts of the injured stem possess brown scars. Major problems occur when root rot organisms have begun to damage roots. Fungus gnat larvae will then begin to attack the rotting tissues and can greatly increase the amount of damage by their feeding. After 14 days, they construct a pupal case made of silk and debris in the soil.

The pupal stage lasts about three days. Adults are weak fliers and are normally seen running rapidly across the surface of the soil when disturbed. All stages of the life cycle may be present because of overlapping generations.

Control

Prevention and sanitation are two of the best control measures. Avoid overwatering since fungus gnats prefer abundant moisture. Poor drainage and water leaks may also increase populations. Plants should be inspected carefully for signs of infestation before purchasing them. Sterile potting mix should be used to prevent introduction of fungus gnats. Houseplants taken outdoors when the weather is warm may become infested before being brought indoors. Inspect plants carefully, destroying those plants that cannot be saved. Old plant material and debris in and around the greenhouse should be removed.

Decoy pots of sprouting grain may be used to attract adult females, which lay their eggs in these pots. Remove and dispose of infested decoy pots every two weeks and replace with new decoy pots of sprouting grain.

Insecticides may be necessary if populations become unmanageable. For effective control, it will be necessary to control both the adults and the larvae. Adult fungus gnats are easily killed with aerosols or sprays. To control larvae, soil drenches or coarse

sprays should be applied to the soil surface. It is best to make applications to moist potting media so that at least the top 2 inches of media are treated.

Home Use Indoors:

azadirachtin (Safer Bioneem), insecticidal soap (Safer Insecticidal Soap), horticultural oil (various brand names), or pyrethrins plus piperonyl butoxide (Pyrenone, PT 1600 X-clude A)

Greenhouse Use:

<http://eppserver.ag.utk.edu/redbook/pdf/ghinsects.pdf>

Interior plantscapes such as hotels, shopping malls, office buildings, etc.:

<http://eppserver.ag.utk.edu/redbook/pdf/interiorscapeinsects.pdf>

Precautionary Statement

To protect people and the environment, pesticides should be used safely. This is everyone's responsibility, especially the user. Read and follow label directions carefully before you buy, mix, apply, store, or dispose of a pesticide. According to laws regulating pesticides, they must be used only as directed by the label. Persons who do not obey the law will be subject to penalties.

Disclaimer Statement

Pesticides recommended in this publication were registered for the prescribed uses when printed. Pesticides registrations are continuously reviewed. Should registration of a recommended pesticide be canceled, it would no longer be recommended by the University of Tennessee.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others which may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product.

THE UNIVERSITY of TENNESSEE 
INSTITUTE of AGRICULTURE

Visit the UT Extension Web site at
<http://utextension.tennessee.edu>

SP341-C 1/11 11-0085

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.
UT Extension provides equal opportunities in programs and employment.