



7-31-2012

## The influence of sample biases on estimations of marine microbial diversity

Caroline Rempe

*University of Tennessee - Knoxville, crempe@utk.edu*

Charles R. Budinoff

*University of Tennessee - Knoxville*

T. C. Effler

*University of Tennessee - Knoxville*

Alison Buchan

*University of Tennessee - Knoxville*

Follow this and additional works at: [https://trace.tennessee.edu/utk\\_micrpubs](https://trace.tennessee.edu/utk_micrpubs)



Part of the [Microbiology Commons](#)

---

### Recommended Citation

BMC Bioinformatics 2012, 13(Suppl 12):A5 doi:10.1186/1471-2105-13-S12-A5

This Article is brought to you for free and open access by the Microbiology at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Microbiology Publications and Other Works by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

MEETING ABSTRACT

Open Access

# The influence of sample biases on estimations of marine microbial diversity

Caroline Rempe<sup>1\*</sup>, Charles R Budinoff<sup>2</sup>, T Chad Effler<sup>2</sup>, Alison Buchan<sup>1,2</sup>

From 11th Annual UT-ORNL-KBRIN Bioinformatics Summit 2012  
Louisville, KY, USA. 30 March - 1 April 2012

## Background

The *Roseobacter* clade is a widespread, metabolically versatile, and easily cultured lineage of marine microbes. Furthermore, cultured isolates are generally considered to be representative of the naturally occurring diversity in the environment. *Roseobacter* is thus considered an ideal clade for exploring and understanding microbial function in a representative model system. Nevertheless, recent studies question whether cultured strains are truly representative of natural populations. Support for one of these studies was based on the potentially biased samples of a Global Ocean Sampling (GOS) expedition [1]. The marine metagenomic studies of the GOS expeditions initially focused on microbes collected from a single filter range (0.1µm – 0.8µm) and it has been previously suggested that this excludes many *Roseobacters*, thus biasing the dataset towards smaller cells.

## Materials and methods

Here, we take advantage of a GOS metagenomic dataset from a 2006-2008 Antarctic expedition that included community analyses over three different size classes (0.1µm-0.8µm; 0.8µm-3µm; and 3µm-200µm). We examined correlations between bacterial taxonomic diversity (based on 16S rRNA gene analyses) and the three size classes to understand the potential for diversity biases between filter ranges. The 16S rRNA gene sequences from these datasets were extracted using Metaxa and subsequently analyzed using MOTHUR, which provided fine-level resolution taxonomic assignments.

## Results

This analysis reveals a potential bias as the ranges of *Roseobacter* subgroup diversity in this dataset varied between size classes. From this analysis we identified specific *Roseobacter* subgroups that were predominantly found in specific size classes. For instance, the OCT and *Roseovarius* subgroups were most prevalent in the 0.8µm-3µm filter range, while the DC5-80 and NAC11-7 subgroups were well represented in both 0.1µm-0.8µm and 0.8µm-3µm filter ranges. These findings reveal the value of analyzing multiple size class fractions when assessing microbial diversity.

## Author details

<sup>1</sup>Genome Science and Technology, The University of Tennessee, Knoxville, TN 37916, USA. <sup>2</sup>Microbiology, The University of Tennessee, Knoxville, TN 37916, USA.

Published: 31 July 2012

## Reference

1. Luo H, Loytynoja A, Moran MA: **Genome content of uncultivated marine *Roseobacters* in the surface ocean.** *Environmental Microbiology* 2012, **14**(1):41-51.

doi:10.1186/1471-2105-13-S12-A5

**Cite this article as:** Rempe et al.: The influence of sample biases on estimations of marine microbial diversity. *BMC Bioinformatics* 2012 **13** (Suppl 12):A5.

\* Correspondence: crempe@utk.edu

<sup>1</sup>Genome Science and Technology, The University of Tennessee, Knoxville, TN 37916, USA

Full list of author information is available at the end of the article