



8-2013

## Complete Genome Sequence of Cyanobacterial Siphovirus KBS2A

Alise J. Ponsoero

Feng Chen

Jay T. Lennon

Steven W. Wilhelm

*University of Tennessee - Knoxville*

Follow this and additional works at: [https://trace.tennessee.edu/utk\\_micrpubs](https://trace.tennessee.edu/utk_micrpubs)

 Part of the [Microbiology Commons](#)

---

### Recommended Citation

Genome Announc. 2013 Jul-Aug; 1(4): e00472-13. Published online 2013 August 22. doi: 10.1128/genomeA.00472-13

This Article is brought to you for free and open access by the Microbiology at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Microbiology Publications and Other Works by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

# Complete Genome Sequence of Cyanobacterial Siphovirus KBS2A

Alise J. Ponsero,<sup>a</sup> Feng Chen,<sup>b</sup> Jay T. Lennon,<sup>c</sup> Steven W. Wilhelm<sup>a</sup>

Department of Microbiology, University of Tennessee, Knoxville, Tennessee, USA<sup>a</sup>; Institute of Marine and Environmental Technology, University of Maryland Center for Environmental Science, Baltimore, Maryland, USA<sup>b</sup>; Department of Biology, Indiana University, Bloomington, Indiana, USA<sup>c</sup>

**We present the genome of a cyanosiphovirus (KBS2A) that infects a marine *Synechococcus* sp. (strain WH7803). Unique to this genome, relative to other sequenced cyanosiphoviruses, is the absence of elements associated with integration into the host chromosome, suggesting this virus may not be able to establish a lysogenic relationship.**

Received 31 May 2013 Accepted 31 July 2013 Published 22 August 2013

Citation Ponsero AJ, Chen F, Lennon JT, Wilhelm SW. 2013. Complete genome sequence of cyanobacterial siphovirus KBS2A. *Genome Announc.* 1(4):e00472-13. doi:10.1128/genomeA.00472-13.

Copyright © 2013 Ponsero et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Steven W. Wilhelm, wilhelm@utk.edu.

As obligate parasites, viruses can regulate their host population dynamics but also influence the structure and productivity of microbial communities (1, 2). *Synechococcus* species are an abundant and ecologically important group of *Cyanobacteria* found in freshwater and marine ecosystems worldwide. Virus-cyanobacterium interactions may have important implications for global biogeochemical cycles. The most commonly isolated cyanophages are myoviruses and podoviruses (3, 4). Siphoviruses are a third group of viruses that infect cyanobacteria, but they have received less attention (5).

The genomes of 5 cyanosiphoviruses have recently become available: that of P-SS2, a siphovirus infecting *Prochlorococcus* (MIT9313) (6), followed by the cyanosiphoviruses S-CBS1, S-CBS2, S-CBS3, and S-CBS4, isolated from the Chesapeake Bay Estuary, all infecting *Synechococcus* populations (5). Here, we present the complete genome of cyanosiphophage (KBS2A, originally named KBS-S-2A), a virus that infects *Synechococcus* sp. strain WH7803.

The virus was isolated by plaque assay from the Chesapeake Bay by plating on *Synechococcus* sp. WH7803. Purified virus DNA was submitted to the Broad Institute as part of the Marine Phage Sequencing Project, where it was sequenced to ~30-fold coverage using 454 pyrosequencing. Translated open reading frames (ORFs) were compared with known protein sequences using the BLASTp program. ORF annotation was aided by the use of PSI-BLAST, HHpred, gene size, and domain conservation.

The genome size of KBS2A is 40,658 bp. In total, 64 ORFs have been predicted in this genome; of these, 43 have homologues in databases, and among them, 33 have been assigned to a putative function. For most (88%) predicted ORFs with homologues, homology has been found with the other cyanosiphovirus genomes. We compared the genomic arrangements of the 6 sequenced cyanosiphoviruses using dot plot and global gene homology and found no common genomic organization, suggesting strong mosaicism in the cyanosiphoviruses.

In cyanophages, cyanobacterium-related proteins can be found and are often associated with photosynthesis and transcriptional regulation (6). In previously sequenced cyanosiphovirus genomes (5, 6), numerous viral genes (6 to 40 per genome) possess

homology with host genes. In the case of the KBS2A genome, only 3 ORFs (coding for RNA polymerase sigma factor RpoD, HNH endonuclease, and a putative DNA polymerase) show such homology, implying less exchange (and potentially interaction) with the host genome.

The first annotated cyanosiphovirus genome (that of P-SS2) showed the presence of genes identified as encoding an integrase and excisionase, which are enzymes that allow for phage integration into the host's genome (6). Moreover, the annotation of cyanosiphoviruses S-CBS1 and S-CBS3 led to the discovery of a prophage-like structure in two sequenced *Synechococcus elongatus* strains (5). In phage genomes, tRNA genes serve as indicators of potential phage integration by site-specific recombination (7, 8), although recent models have offered alternative suggestions for the role of these genes (9). Sequences of this nature can, however, be found in the P-SS2 and S-CBS4 genomes. No such features (tRNAs, integrases, etc.) were found in the genome of KBS2A, suggesting that this siphovirus might be an exclusively lytic phage rather than a temperate phage.

**Nucleotide sequence accession number.** The complete sequence of the *Synechococcus* phage KBS2A genome can be accessed under the GenBank accession no. [HQ634187](https://www.ncbi.nlm.nih.gov/nuclseq/HQ634187).

## ACKNOWLEDGMENTS

Sequencing and assembly were funded in part by the Gordon and Betty Moore Foundation through a grant to the Broad Institute, where this work was completed. A.J.P. was supported by an NSF grant to S.W.W. (OCE-0851113).

## REFERENCES

1. Suttle CA, Chan AM. 1994. Dynamics and distribution of cyanophages and their effect on marine *Synechococcus* spp. *Appl. Environ. Microbiol.* 60:3167–3174.
2. Weinbauer MG, Bonilla-Finji O, Chan AM, Agis M, Dolan JR, Short SM, Simek K, Wilhelm SW, Suttle CA. 2011. *Synechococcus* growth in the ocean may depend on the lysis of heterotrophic bacteria. *J. Plankton Res.* 33:1465–1476.
3. Sullivan MB, Coleman ML, Quinlivan V, Rosenkranz JE, DeFrancesco AS, Tan G, Fu R, Lee JA, Waterbury JB, Bielawski JP, Chisholm SW. 2008. Portal protein diversity and phage ecology. *Environ. Microbiol.* 10: 2810–2823.

4. Fuller NJ, Wilson WH, Joint IR, Mann NH. 1998. Occurrence of a sequence in marine cyanophages similar to that of T4 g20 and its application to PCR-based detection and quantification techniques. *Appl. Environ. Microbiol.* **64**:2051–2060.
5. Huang S, Wang K, Jiao N, Chen F. 2012. Genome sequences of siphoviruses infecting marine *Synechococcus* unveil a diverse cyanophage group and extensive phage-host genetic exchanges. *Environ. Microbiol.* **14**:540–558.
6. Sullivan MB, Krastins B, Hughes JL, Kelly L, Chase M, Sarracino D, Chisholm SW. 2009. The genome and structural proteome of an ocean siphovirus: a new window into the cyanobacterial ‘mobilome.’ *Environ. Microbiol.* **11**:2935–2951.
7. Campbell A. 2003. Prophage insertion sites. *Res. Microbiol.* **154**:277–282.
8. Canchaya C, Fournous G, Brussow H. 2004. The impact of prophages on bacterial chromosomes. *Mol. Microbiol.* **53**:9–18.
9. Bailly-Bechet M, Vergassola M, Rocha E. 2007. Causes for the intriguing presence of tRNAs in phages. *Genome Res.* **17**:1486–1495.