



5-14-2013

W289-D IPM QuickFacts Series: Phytophthora Root Rot

Amy Fulcher

Heather Bowers

Follow this and additional works at: https://trace.tennessee.edu/utk_agexdise



Part of the [Agricultural Education Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

"W289-D IPM QuickFacts Series: Phytophthora Root Rot," Amy Fulcher and Heather Bowers,
W 289-D
, https://trace.tennessee.edu/utk_agexdise/82

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Ornamental insects, Pests & Diseases is brought to you for free and open access by the UT Extension Publications at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Insects, Pests, Plant Diseases and Weeds by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Phytophthora Root Rot

Phytophthora spp.

Host Plants

Arborvitae	Laurel
Barberry	Leucothoe
Birch	Leyland cypress
Blackgum	Lilac
Blueberry	Magnolia
Boxwood	Maple
Buckeye	Nandina
Butterfly-bush	Oak
Camellia	Ornamental cherry
Cedar	Pear
Chestnut	Pieris
Cotoneaster	Pine
Crabapple	Privet
Dogwood	Rhododendron
Elm	Spruce
Euonymus	Sweetgum
Fir	Sycamore
Forsythia	Willow
Hemlock	Wisteria
Holly	Yew
Honeylocust	Zelkova
Juniper	and more
Larch	

Introduction



The name *Phytophthora* means plant destroyer. *Phytophthora* is a soil-borne, fungus-like organism that infects and kills a range of nursery and landscape plants.

Pathogen and Disease Cycle

Phytophthora overwinters in soil. Extended periods of high soil moisture favor infection. Sporangia (sacs of spores) cause new infections by germinating directly and colonizing roots or by releasing many zoospores (motile spores) that formed inside the sporangium into water. *Phytophthora* spreads via contaminated water (amounts as small as droplets) or soil (including reused media and pots). Recent studies have found that the pathogen can also be spread by fungus, gnats and shore flies.

Monitoring

Look for symptoms during the growing season, when temperatures are high and plants are actively growing. Infected roots may be tan to brown to cinnamon; conifers may lack white root-tips. Many of the aboveground symptoms may be confused with a nutritional disorder or over or under watering. Have a diagnostic laboratory run tests to confirm the presence of this pathogen.



Symptoms

Phytophthora kills the roots and crown of infected plants. It reduces the volume of the roots, which limits the uptake of water and nutrients. A common symptom is wilted foliage, even when adequate moisture is present. Wilting causes a need to water more. Other symptoms include yellow or bronze foliage, branch dieback and poor plant vigor; death is often common.



Integrated Pest Management

CULTURAL CONTROL

Plant in raised beds with well-drained soil. Avoid over irrigating. Avoid planting too deep. Remove organic debris and soak containers in disinfectant or steam before reuse. See Adkins et al. (below) for disinfestation guidelines. Media should be stored on a concrete pad. Store containers on gravel or concrete. Recycled irrigation water should be tested and/or sanitized before reusing.

CHEMICAL CONTROL

Please refer to http://eppserver.ag.utk.edu/redbook/sections/trees_flowers.htm for the most up-to-date recommendations.

Resources

Photo credits: Amy Fulcher and Alan Windham, University of Tennessee

Adkins, C., G. Armel, M. Chappell, J.C. Chong, S. Frank, A. Fulcher, F. Hale, K. Ivors, W. Klingeman III, A. LeBude, J. Neal, A. Senesac, S. White, and A. Windham. 2010. Pest management strategic plan for container and field-produced nursery crops in GA, KY, NC, SC, TN. A. Fulcher, ed. Southern Region IPM Center.

Vincelli, P. and D. Hershman. 2005. Controlling phytophthora root rot in greenhouse ornamentals. University of Kentucky Extension publication PPF5-OR-H-09. http://www.ca.uky.edu/aqcollege/plantpathology/ext_files/PPFShtml/PPFS-OR-H-9.pdf

Windham, A. 2007. Phytophthora diseases in ornamentals. The University of Tennessee/Agricultural Extension Service. What's Happening Newsletter (11). <http://eppserver.ag.utk.edu/Whats/wh2007/Issue-11-2007.pdf>

THE UNIVERSITY of TENNESSEE 
INSTITUTE of AGRICULTURE

Prepared by Heather Bowers, Intern, Department of Plant Sciences
and Dr. Amy Fulcher, Assistant Professor, Department of Plant Sciences
Publication funded by USDA Extension IPM Grant in partnership with University of Kentucky Integrated Pest Management Program.
The authors thank M. Halcomb, W. Russell and A. Windham for their careful review.

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development.
University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.
UT Extension provides equal opportunities in programs and employment.