



3-14-2019

Cryopreservation of Paramecium bursaria Chlorella Virus-1 during an active infection cycle of its host

Samantha R. Coy
University of Tennessee, Knoxville

Alyssa N. Alsante
University of Tennessee, Knoxville

James L. Van Etten
University of Nebraska, Lincoln

Steven W. Wilhelm
University of Tennessee, Knoxville, wilhelm@utk.edu

Follow this and additional works at: https://trace.tennessee.edu/utk_micrpubs

Recommended Citation

Coy SR, Alsante AN, Van Etten JL, Wilhelm SW (2019) Cryopreservation of Paramecium bursaria Chlorella Virus-1 during an active infection cycle of its host. PLoS ONE 14(3): e0211755. <https://doi.org/10.1371/journal.pone.0211755>

This Article is brought to you for free and open access by the Microbiology at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Microbiology Publications and Other Works by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

RESEARCH ARTICLE

Cryopreservation of *Paramecium bursaria* *Chlorella Virus-1* during an active infection cycle of its host

Samantha R. Coy¹, Alyssa N. Alsante^{1†}, James L. Van Etten², Steven W. Wilhelm^{1*}

1 Department of Microbiology, University of Tennessee, Knoxville, Tennessee, United States of America,

2 Department of Plant Pathology and Nebraska Center for Virology, University of Nebraska, Lincoln, Nebraska, United States of America

† Current address: Department of Oceanography, Texas A&M University, College Station, Texas, United States of America

* wilhelm@utk.edu



OPEN ACCESS

Citation: Coy SR, Alsante AN, Van Etten JL, Wilhelm SW (2019) Cryopreservation of *Paramecium bursaria* *Chlorella Virus-1* during an active infection cycle of its host. PLoS ONE 14(3): e0211755. <https://doi.org/10.1371/journal.pone.0211755>

Editor: Susanna A Wood, Cawthron Institute, NEW ZEALAND

Received: January 18, 2019

Accepted: March 1, 2019

Published: March 14, 2019

Copyright: © 2019 Coy et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript.

Funding: This work was supported by grants from the National Science Foundation (NSF-OCE 1829641) and the Gordon & Betty Moore Foundation (#4971) to SWW. Funding for open access to this research was provided by University of Tennessee's Open Publishing Support Fund. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Best practices in laboratory culture management often include cryopreservation of microbiota, but this can be challenging with some virus particles. By preserving viral isolates researchers can mitigate genetic drift and laboratory-induced selection, thereby maintaining genetically consistent strains between experiments. To this end, we developed a method to cryopreserve the model, green-alga infecting virus, *Paramecium bursaria* *Chlorella virus 1* (PBCV-1). We explored cryotolerance of the infectivity of this virus particle, whereby freezing without cryoprotectants was found to maintain the highest infectivity (~2.5%). We then assessed the cryopreservation potential of PBCV-1 during an active infection cycle in its *Chlorella variabilis* NC64A host, and found that virus survivorship was highest (69.5 ± 16.5%) when the infected host is cryopreserved during mid-late stages of infection (*i.e.*, coinciding with virion assembly). The most optimal condition for cryopreservation was observed at 240 minutes post-infection. Overall, utilizing the cell as a vehicle for viral cryopreservation resulted in 24.9–30.1 fold increases in PBCV-1 survival based on 95% confidence intervals of frozen virus particles and virus cryopreserved at 240 minutes post-infection. Given that cryoprotectants are often naturally produced by psychrophilic organisms, we suspect that cryopreservation of infected hosts may be a reliable mechanism for virus persistence in non-growth permitting circumstances in the environment, such as ancient permafrosts.

Introduction

Viruses are abundant components of all biological systems and they likely infect every lineage of eukaryotic algae. Their impact is most readily noticed following infection and lysis of abundant bloom forming algae [1–3], though lytic activity of all algal viruses contributes to significant biomass recycling *via* the ‘viral shunt’ [4]. To date, 65 eukaryotic algal viruses have been isolated and developed as laboratory strains [5, 6]. Most of these are maintained through serial

Competing interests: The authors have declared that no competing interests exist.

propagation on their respective hosts. Though this has been effective for culturing many strains over the last few decades [7, 8], each passage allows for genetic mutations that can accumulate in a population [9], leading to a deviation from a standard ‘wild-type.’ Moreover, it is imperative to control evolution following the development of genetically tractable algal hosts [10] and (ultimately) virus systems. Although seed-stock systems can be developed without cryopreservation, many systems are not amenable to this either because the virus particles are degraded during purification efforts or lose their infectivity during storage. Moreover, it can take time to achieve axenic status with new virus isolates, thus making contaminating bacterial activity a significant source of degradation. Thus, a protocol for successful virus cryobiological preservation that is applicable to a wide variety of algae-virus systems would offer an opportunity to universally improve virus management and distribution in the laboratory.

Cryopreservation is not a new concept in biological sciences. For most protocols, it involves controlled cooling of biota to sub-freezing temperatures to achieve biological cessation while preserving viability. This most often manifests as slow-cooling at a rate of 1°C / min in the presence of osmoprotectant(s) (e.g., dimethylsulfoxide (DMSO), glycerol) for long-term storage at -130°C or below [11]. Too slow a cooling rate can result in higher intracellular concentration of osmoprotectants, resulting in toxicity, whereas too fast a cooling rate allows the formation of intracellular ice crystals which can rupture cell membranes [12]. The thawing process is typically quick, as microbial death is commonly associated with slow thaw rates. Though cryopreservation is a standard method for maintaining cellular organisms, it has rarely been utilized for the preservation of algal viruses.

One eukaryotic algal virus cryopreservation protocol is in existence. It was developed for HaV, a dsDNA virus that infects the red tide forming dinoflagellate *Heterosigma akashiwo* [13]. Researchers investigated a combination of cryoprotectants and storage temperatures with the highest recovery (8.3% of infectious virus) employing flash freezing of HaV particles suspended in 20% DMSO. This protocol has been adapted for a handful of other algal viruses with viable recovery ranging from < 1% to 27% [14–16]. The typical low recovery in these procedures is likely due to physiological differences between viruses and cells including differences in permeability, osmolarity tolerance, and toxicity to osmoprotectants. It is also clear that these protocols deviate from the standard method which controls the cooling rate; to our knowledge this has not been tested as a matter of improving virus particle survival. Owing to these complications, we decided to take a new approach by investigating cryopreservation recovery and stability of actively infecting, cell-associated algal viruses.

Chloroviruses are large (> 300 kb) dsDNA viruses in the family *Phycodnaviridae* [17]. They are members of the proposed order the Megavirales [18], also known as “giant” viruses, and remain the best characterized algal-virus system to date. Isolated in the early 1980’s [7], the prototype chlorovirus *Paramecium bursaria Chlorella virus 1* (PBCV-1) has been maintained through serial propagation on its host, *Chlorella variabilis* NC64A. PBCV-1 is inactivated by freezing, though other closely related virus strains, including other chloroviruses, persist through freeze/thaw events [19, 20]. As a great deal of research has centered on PBCV-1, including genomics [21], transcriptomics [22, 23], and proteomics [21], it is important to develop a successful cryopreservation protocol for this strain that may serve as a model for preserving algal viruses. There are several reports of cryopreservation techniques for eukaryotic algae [24–28] which might be adapted for the preservation of actively replicating chloroviruses.

Here, we tested the cryo-potential of chlorovirus PBCV-1 using a protocol that yielded consistent recovery (~50% viable cells) of four strains of algae over 15 years: *Chlorella vulgaris* C-27, *Chlorella vulgaris* M-207A7, *Nannochloropsis oculata* ST-4, and *Tetraselmis tetraethle* T-501 [29]. Owing to the close relationship between *C. vulgaris* and *C. variabilis*, as well as the

consistent results across unique algae, we elected to determine if these results could be recapitulated in PBCV-1. To test this, we attempted cryopreservation of both the virus particle as well as the virus replicating in its host.

Materials and methods

Virus particle cryopreservation

Chlorella variabilis NC64A was infected with PBCV-1 during mid-logarithmic growth at standard culturing conditions (25°C; continuous light exposure at 30 μ Ein/m²/s) using Modified Bold's Basal Medium [30]. Following complete lysis, the viral lysate was pre-filtered through a sterile, 0.45 μ m polycarbonate syringe filter and titered by plaque assay [31, 32] for initial infectivity assessments. Cryoprotectant choice was guided by Nakanishi *et al.* [29], in which a combination of 5% DMSO (v/v), 5% ethylene glycol (v/v), and 5% proline (w/v) was found to consistently produce the highest algal recoveries. Stock solutions of each cryoprotectant were made at a concentration of 30% with sterilized Milli-Q water and combined in a 1:1:1 ratio to yield a final concentration of 10% for each compound. For virus particle cryopreservation, 1 mL of PBCV-1 particles (7.82x 10⁸ plaque forming units (PFUs) per ml) was added to 1 mL of ice-chilled cryoprotectant solution contained in a 2-mL cryovial. The cryovials were incubated on ice for 45 min, then transferred to a freeze-rate controlled container (Mr. Frosty, Thermo Fisher Scientific Inc., USA) filled with isopropanol for overnight incubation at -80°C. The next morning, cryovials were transferred to a -150°C freezer. At the designated recovery times, vials were removed from the freezer and set in a 40°C water bath. After thawing, the samples were serially diluted ten-fold in 50 mM Tris-HCl (pH = 7.8) and virus infectivity was determined by plaque assay [31]. Virus viability was calculated as a percentage by comparison to the initial virus particle stock titer before cryopreservation. Long-term experiments assessed the stability of virus infectivity in particles stored at -150°C.

Infected *Chlorella* cryopreservation

Chlorovirus PBCV-1 was propagated as described above and titered to obtain infectious PFUs/ml. This virus particle stock was used to infect late-logarithmically growing *C. variabilis* NC64A at an M.O.I. of 5, at which point infected cultures were returned to standard incubation conditions. At 1, 10, 30, 60, 120, 180, 240, 300, and 360 min post-infection (PI), 1 mL aliquots of infected cells were mixed with 1 mL of ice-chilled cryoprotectants [final concentration: 5% DMSO (v/v), 5% ethylene glycol (v/v), and 5% proline (w/v)] in duplicates. The mixture was incubated on ice for 45 min, then transferred to a freeze-rate controlled container (Mr. Frosty, Thermo Fisher Scientific Inc., USA has a -1C/min cooling rate) filled with isopropanol for overnight incubation at -80°C. The next morning, cryovials were immediately transferred to a -150°C freezer. At the designated recovery times, vials were removed from the freezer and placed in a 40°C water bath. After thawing, the infected cells were pelleted in a Sorvall Legend RT Benchtop Centrifuge at 3,700 rpm (~3,000 rcf) for 10 min: (free virus requires higher speeds for pelleting). Cell pellets were re-suspended in 2 mL of 0.01M HEPES solution (pH = 6.5). Suspensions were immediately diluted and plaque assayed, plating late-infection treatments first. Viability was determined as a percentage of the pre-frozen cellular concentration (3.57 x 10⁶ cells/mL), as only surviving infected cells would be capable of producing plaques. Long-term experiments were conducted in the same manner, though only time points 10, 180, and 240 min PI were collected and assayed. The complete step-by-step method can be found at protocols.io [33].

Results

Following the cryopreservation procedures of other algal virus researchers [13–16], we investigated the cryo-potential of the PBCV-1 particle. Cryoprotectant alone treatments elicited a lethal effect: ~87% of the infectious virus particles were inactivated in the presence of these chemicals following 24 hr exposure at 4°C. Given this effect, we decided to freeze PBCV-1 particles at -150°C without any cryoprotectants. This resulted in ~2.5% recovery of the infectious virus population, which was stable for storage periods of up to one year (Fig 1). Seeing room for improvement, we tested the cryo-potential of PBCV-1 in an infected, cell-associated state.

The PBCV-1 replication cycle requires about 6–8 h to release nascent virus particles [34]. Post-infection sampling times for cryopreservation (10, 30, 60, 120, 180, 240, 300, 360 min PI) followed similar sampling strategies used in PBCV-1 transcription studies [22, 23]. Specifically, these time points were collected across distinct physiological phases in the PBCV-1 lifecycle and thus represent likely unique conditions for cryopreservation. Following 24-h storage of cryopreserved, infected cells, we found that late stages of infection were more conducive to virus survival than early stages (Fig 2). Thus, we followed cryo-stability for one year in one early (10 min PI) and two late infection stages (180 and 240 min PI) (Fig 3). Small day-to-day fluctuations in virus titers were common, but were typically consistent among treatments, suggesting human error. Despite these fluctuations, the virus particle stock control, 180-min, and 240-min PI treatment yielded an acceptable relative standard deviation (RSD) for these plate counts [35] across all recovery assessments, indicating cryo-stability (Table 1). Cryo-stability




Fig 1. Cryo-stability of the PBCV-1 particle. Viability of chlorovirus PBCV-1 was determined by plaque assaying viruses that had been stored as particles either at 4°C or -150°C. Green circles represent virus particles stored at 4°C, while red squares denote virus particles stored at -150°C. Error bars are represented as the standard deviation of biological and technical replicates.

<https://doi.org/10.1371/journal.pone.0211755.g001>




Fig 2. Recovery of infectious PBCV-1 frozen at various times after PBCV-1 infection of the *C. variabilis* NC64A host. Viability of chlorovirus PBCV-1 was assayed by monitoring plaque formation of cell-associated viruses that were collected at different times during an active infection cycle of the NC64A host. Open circles denote replicate plaque titers, with the average represented by the solid line.

<https://doi.org/10.1371/journal.pone.0211755.g002>

was not observed in the 10 min PI samples (Table 1). In comparison to virus particle cryopreservation, the cell-associated method yielded significant improvement in survivorship for the optimal 240-minute treatment (24.9–30.1 fold increases).

Discussion

The current maintenance strategy for chloroviruses involves serial propagation on the alga host followed by lysate particle storage at 4°C. Chloroviruses are relatively stable under these conditions, though even PBCV-1 is known to degrade after several years of storage. In any case, many algae-virus systems are less amenable to long-term storage at 4°C. For example, new algae-virus systems are not always quickly made axenic, and are thus susceptible to degradation from contaminating bacteria. On the other hand, viruses propagated on axenic hosts can still degrade. For reasons unknown, chloroviruses are more stable in lysates (bacterial-free) than in particle stocks purified by sucrose density gradients [36], but they always eventually lose their infectivity. Serial propagation of viruses is therefore often required. Even if this is done infrequently, it can still promote genetic drift and result in deviation from wild-type status. This is concerning for all virus types, though RNA viruses, which have the fastest mutation rates, would be most susceptible [9, 37]. Beyond considering spontaneous, replication-associated errors, chloroviruses encode putative enzymes involved in genomic rearrangements. For




Fig 3. Long-term cryo-stability of PBCV-1 frozen in host cells at various times after infection of its host *C. variabilis* NC64A. Infectious chlorovirus PBCV-1 was monitored by plaque assay in virus particle stocks stored at 4°C (green circles) and in cryopreserved, PBCV-1-infected host cultures. Blue triangles, yellow squares, and orange diamonds represent virus viability following storage of infected cells cryopreserved after 240, 180, and 10 minutes PI. Error bars represent the standard deviation among biological and technical replicates.

<https://doi.org/10.1371/journal.pone.0211755.g003>

example, GIY-YIG mobile endonucleases and an IS607 transposon may be involved in insertions/deletions and/or gene loss/duplications observed in genomic comparisons of chloroviruses [38, 39]. Thus, maintenance of wild-type strains is important for consistency between experiments. Virology labs could follow the microbial culture collection strategy, which typically uses a cryo-banking/seed-stock system for the dissemination of microbial specimens. The purpose of the seed-stock system is to minimize serial propagation of microbiota. The American Type Culture Collection (ATCC) suggests that consumers transfer their cultures no more than five-times after propagation from the thawed culture collection stock. Though a

Table 1. Statistical assessment of PBCV-1 infectivity across storage treatments for ~1 year.

Treatment	N	Average	SD	RSD	95%CI
Virus Particle Stock (4°C)	67	75.1	16.9	22.5*	71.1–79.2
Virus Particle Stock (-150°C)	124	2.53	0.61	24.0*	2.42–2.64
Cell-associated virus 10 minutes PI (-150°C, +CPA)	79	7.56	3.38	44.7	6.81–8.31
Cell-associated virus 180 minutes PI (-150°C, +CPA)	82	31.9	10.9	34.2*	29.5–34.3
Cell-associated virus 240 minutes PI (-150°C, +CPA)	82	69.5	16.5	23.8*	65.9–73.0

+CPA, cryoprotectants present as described in materials and methods section. Asterisks (*) denote an acceptable RSD (i.e., Coefficient of Variation) for plaque assays based on a 35% threshold used in bacterial plating standards set from chapter 1223 by the U.S. Pharmacopeia and National Formulary.

<https://doi.org/10.1371/journal.pone.0211755.t001>

seemingly strict standard, it is not difficult to imagine the consequences of violating this. For example, the United States Pharmacopeia and National Formulary requires test organisms to be maintained this way for routine antibiotic efficacy screens, and non-compliance can undermine therapeutic treatment [35]. Although there is no direct clinical link to maintaining algal viruses this way, the logic is consistent with any research requirements. The cryopreservation protocol described here can help researchers better set up these cryo-banking/seed stock systems.

Standard cryopreservation techniques are not designed for the unique structure and physiology of virus particles. Indeed, cryoprotectants are classified by their permeability across cell membranes, which often coincides with their molecular weight [24]. Smaller compounds, such as ethylene glycol and DMSO, are considered penetrating cryoprotectants, while larger compounds (e.g. amino acids; L-proline) are typically non-penetrating. That said, the exclusion size threshold has not been established for most viruses so it is not clear which, if any of these compounds penetrate the viral capsid. It is generally thought that virus capsids are permeable to water and ions, though the latter diffuses much slower; this mechanism has been used to osmotically rupture capsids [40, 41], including PBCV-1 [42]. The final cryoprotectant solution used for PBCV-1 particle cryopreservation has an estimated osmolarity of ~150 mOsmoles/L, which is comparable to the storage buffer used for this virus. In light of this, we propose that the lethal effect the cryoprotectants have on the PBCV-1 particle is not the result of osmotic stress, and that inactivation instead occurred by toxicity of cryoprotectants or oxidative stress. This would be consistent with viruses not being metabolically active and therefore unable to repair damage caused by this treatment. It is also consistent with the observation that Mimivirus, a giant virus relative which also contains an internal lipid membrane, is said to be inactivated by lipophilic compounds such as DMSO [43]. That said, DMSO is often used as a stabilizer for freezing of enveloped virus particles [44]. This discrepancy may be due to unique properties between external and internal membranes, or even system differences between animal and plant viruses, which imparts resistance in some cases over others. Regardless, the mechanism of inactivation may be better ascertained by looking at survivorship of virion particles via epifluorescent microscopy, flow cytometry [45–47], or using bioassays to quantify oxidative stress.

Although the algal cell is in a sub-optimal physiological state during infection, it is apparently robust enough to survive and maintain an active infection during cryopreservation. That said, fewer infectious virus were recovered when the cell was cryopreserved during early infection stages. This might be explained by differences in adsorption rates and synchronicity of infection, resulting in fewer infected cells at the start of the experiment. Most, if not all cells are infected at the later stages of infection (3–4 hr PI). Regardless of any differences in synchronicity, the algal cell will be completely arrested during cryopreservation, and will only continue the infection cycle after thawing. Internal, mature viruses that have not yet lysed their host cell might still be inactivated by cryoprotectants, thus reducing viral burst size, but our experiments did not account for this. We also did not account for inefficiencies in infection rates; though we infected at M.O.I. values based on infectious particle counts, it is possible that all the cells were not infected. Had we plated the infected cell population prior to cryoprotection we could have corrected for this in our results. In any case, accounting for infection inefficiency can only improve PBCV-1 survivorship and the success of our method.

The general classification of cryoprotectants based on membrane permeability is consistent in the infected cell treatment. Although the *C. variabilis* NC64A genome encodes a secondary active transporter for the uptake of proline, radio-labeled solute uptake experiments revealed that PBCV-1 infection abolishes its activity [48]. With that in mind, the tonicity of the cryoprotectant mixture would equate to ~90 mOsmoles/L, as only DMSO and ethylene glycol are

penetrating, and many of the components in the MBBM media would be spent by late-logarithmic growth. This concentration is comparable to buffers routinely used in our lab for handling *C. variabilis* (40 mOsmoles/L), so there is little concern of osmotic stress. The chances of osmotic stress were also low considering the consistent success associated with this cryopreservation formula across eukaryotic algae, including two *Chlorella* spp. [29]. Our results are likely applicable to any algal virus whose host can be cryopreserved. That said, we expect that researchers may still have to adjust their cryoprotectant mixture to account for system differences related to osmolarity tolerance and cryoprotectant toxicity. There has also been research indicating that axenicity impacts cryopreservation survival in microalgae. In this light, it is possible that the bacterial community produces secondary metabolites which promote survival [49]. In another scenario, organisms with psychrophilic tendencies might be adapted to freeze situations and cryoprotectant additives may not be necessary.

The goal of this study was to develop a long-term cryopreservation method for chlorovirus PBCV-1, but there are also interesting ecological implications of this research. Recent metagenomic and isolation efforts indicate that giant viruses of microeukaryotes (e.g., *Phycodnaviridae* and *Mimiviridae*) are widely distributed in nature [50, 51], but it is not well understood how these viruses persist in the environment. Freezing events represent a potential mechanism of inactivation for some algal viruses, though chlorovirus ATCV-1 is stable during these conditions [19]. In two other studies, a closely related giant virus of the family *Mimiviridae* [52], as well as a second giant virus in the family *Molliviridae* [53], were revived from 30,000 year old permafrost. Both of these viruses were revived using *Acanthamoeba* spp., one of the main hosts for many giant viruses. That said, there have been questions about whether *Acanthamoeba* and other protists used for laboratory viral propagation are the natural or primary hosts of these ancient viruses [54]. Although these viruses might be able to withstand freezing temperatures on their own, the results of this study suggest that a natural host might serve as a better vehicle for surviving freezing. Indeed, many microbes produce natural cryoprotectants (e.g. L-proline, trehalose, betaine, etc.) or encode machinery to transport these osmoprotectants into the cell. Following this thought process, it is possible that environments containing frozen, infected cells might contain naturally cryopreserved algal-virus systems. These systems may be deciphered following advances in single-cell sorting and sequencing techniques. Indeed, a similar approach has been successfully utilized to identify and sequence single virus genomes in the ocean [55]. Though this latter study sorted virus particles, flow-cytometry sorting of viral infected cells may be achieved using fluorescent probes specific for viral marker genes (e.g., major capsid protein) or dyes to detect viral-induced host phenotypes (e.g., membrane blebbing). As a proof of concept, viral genetic sequences recovered from Siberian permafrost could be used to probe for still frozen viral-infected host cells, thereby testing the natural host range of these viruses.

To our knowledge, this is the first report of successful cryopreservation of a eukaryotic algal virus during its infection cycle. We expect that respective cellular hosts will provide more suitable physiological conditions for cryopreservation and storage of algal viruses that infect eukaryotic algae. We also recommend that laboratories working with algal viruses establish cryopreserved seed-stock systems to better preserve wild-type controls for future experimentation, especially in lieu of future modification of these viral systems.

Acknowledgments

We thank Professor David A. Hutchins (University of Southern California) for his thoughtful discussion on this matter over 2 decades ago. We would also like to thank Dr. Irina Agarkova (University of Nebraska, Lincoln) for her insights on chlorovirus stability over many years

longer than our experiments accounted for. This work was supported by grants from the National Science Foundation (NSF-OCE 1829641) and the Gordon & Betty Moore Foundation (#4971) to SWW. Funding for open access to this research was provided by University of Tennessee's Open Publishing Support Fund.

Author Contributions

Conceptualization: Samantha R. Coy, Steven W. Wilhelm.

Data curation: Samantha R. Coy, Alyssa N. Alsante.

Formal analysis: Samantha R. Coy.

Funding acquisition: Steven W. Wilhelm.

Investigation: Samantha R. Coy, Alyssa N. Alsante.

Methodology: Samantha R. Coy, Alyssa N. Alsante, James L. Van Etten.

Project administration: Steven W. Wilhelm.

Resources: James L. Van Etten.

Supervision: Steven W. Wilhelm.

Validation: Samantha R. Coy.

Visualization: Samantha R. Coy.

Writing – original draft: Samantha R. Coy, Alyssa N. Alsante, James L. Van Etten, Steven W. Wilhelm.

Writing – review & editing: Samantha R. Coy, Alyssa N. Alsante, James L. Van Etten, Steven W. Wilhelm.

References

1. Bratbak G, Egge JK, Heldal M. Viral mortality of the marine alga *Emiliania-huxleyi* (Haptophyceae) and termination of algal blooms. *Mar Ecol Prog Ser*. 1993; 93(1–2):39–48. <https://doi.org/10.3354/meps093039>
2. Rowe JM, Dunlap JR, Gobler CJ, Anderson OR, Gastrich MD, Wilhelm SW. Isolation of a non-phage-like lytic virus infecting *Aureococcus anophagefferens*. *J Phycol*. 2008; 44(1):71–6. <https://doi.org/10.1111/j.1529-8817.2007.00453.x> PMID: 27041042
3. Nagasaki K, Tomaru Y, Nakanishi K, Hata N, Katanozaka N, Yamaguchi M. Dynamics of *Heterocapsa circularisquama* (Dinophyceae) and its viruses in Ago Bay, Japan. *Aquat Microb Ecol*. 2004; 34(3):219–26. <https://doi.org/10.3354/ame034219>
4. Wilhelm SW, Suttle CA. Viruses and nutrient cycles in the sea—viruses play critical roles in the structure and function of aquatic food webs. *BioScience*. 1999; 49(10):781–8. <https://doi.org/10.2307/1313569>
5. Coy SR, Gann ER, Pound HL, Short SM, Wilhelm SW. Viruses of eukaryotic algae: diversity, methods for detection, and future directions. *Viruses*. 2018; 10(9). Epub 2018/09/14. <https://doi.org/10.3390/v10090487> PMID: 30208617.
6. Short SM, Staniewski MA, Chaban YV, Long AM, Wang D. Diversity of viruses infecting eukaryotic algae. In: H P., Abedon ST, editors. *Viruses of microorganisms*. Poole, UK: Caister Academic Press; 2018. p. 211–44.
7. Van Etten JL, Burbank DE, Xia Y, Meints RH. Growth-cycle of a virus, PBCV-1, that infects *Chlorella*-like algae. *Virology*. 1983; 126(1):117–25. [https://doi.org/10.1016/0042-6822\(83\)90466-x](https://doi.org/10.1016/0042-6822(83)90466-x) PMID: 18638936
8. Castberg T, Thyraug R, Larsen A, Sandaa RA, Heldal M, Van Etten JL, et al. Isolation and characterization of a virus that infects *Emiliania huxleyi* (Haptophyta). *J Phycol*. 2002; 38(4):767–74.
9. Peck KM, Lauring AS. Complexities of viral mutation rates. *J Virol*. 2018; 92(14):8. <https://doi.org/10.1128/jvi.01031-17> PMID: 29720522

10. Waller RF, Cleves PA, Rubio-Brotons M, Woods A, Bender SJ, Edgcomb V, et al. Strength in numbers: collaborative science for new experimental model systems. *PLoS Biol.* 2018; 16(7):10. <https://doi.org/10.1371/journal.pbio.2006333> PMID: 29965960
11. Mazur P. Freezing of living cells—mechanisms and implications. *Am J Physiol.* 1984; 247(3):C125–C42
12. Mazur P, Leibo SP, Chu EHY. A two-factor hypothesis of freezing injury—evidence from chineses-hamster tissue-culture cells. *Exp Cell Res.* 1972; 71(2):345–55. [https://doi.org/10.1016/0014-4827\(72\)90303-5](https://doi.org/10.1016/0014-4827(72)90303-5) PMID: 5045639
13. Nagasaki K, Yamaguchi M. Cryopreservation of a virus (HaV) infecting a harmful bloom causing micro-alga, *Heterosigma akashiwo* (Raphidophyceae). *Fish Sci.* 1999; 65(2):319–20. <https://doi.org/10.2331/fishsci.65.319>
14. Kim J, Kim CH, Youn SH, Choi TJ. Isolation and physiological characterization of a novel algicidal virus infecting the marine diatom *Skeletonema costatum*. *Plant Pathol J.* 2015; 31(2):186–91. <https://doi.org/10.5423/PPJ.NT.03.2015.0029> PMID: 26060438
15. Kim J, Yoon SH, Choi TJ. Isolation and physiological characterization of a novel virus infecting *Stephanopyxis palmeriana* (Bacillariophyta). *Algae.* 2015; 30(2):81–7. <https://doi.org/10.4490/algae.2015.30.2.081>
16. Kim J, Kim CH, Takano Y, Jang IK, Kim SW, Choi TJ. Isolation and physiological characterization of a new algicidal virus infecting the harmful dinoflagellate *Heterocapsa pygmaea*. *Plant Pathol J.* 2012; 28(4):433–8. <https://doi.org/10.5423/ppj.nt.07.2012.0093>
17. Jeanniard A, Dunigan DD, Gurnon JR, Agarkova IV, Kang M, Vitek J, et al. Towards defining the chloroviruses: a genomic journey through a genus of large DNA viruses. *BMC Genomics.* 2013; 14. <https://doi.org/10.1186/1471-2164-14-158> PMID: 23497343
18. Colson P, De Lamballerie X, Yutin N, Asgari S, Bigot Y, Bideshi DK, et al. "Megavirales", a proposed new order for eukaryotic nucleocytoplasmic large DNA viruses. *Arch Virol.* 2013; 158(12):2517–21. <https://doi.org/10.1007/s00705-013-1768-6> PMID: 23812617
19. Long AM, Short SM. Seasonal determinations of algal virus decay rates reveal overwintering in a temperate freshwater pond. *ISME J.* 2016; 10(7):1602–12. <https://doi.org/10.1038/ismej.2015.240> PMID: 26943625
20. Bubeck JA, Pfitzner AJP. Isolation and characterization of a new type of chlorovirus that infects an endosymbiotic *Chlorella* strain of the heliozoon *Acanthocystis turfacea*. *J Gen Virol.* 2005; 86:2871–7. <https://doi.org/10.1099/vir.0.81068-0> PMID: 16186243
21. Dunigan DD, Cerny RL, Bauman AT, Roach JC, Lane LC, Agarkova IV, et al. Paramecium bursaria chlorella virus 1 proteome reveals novel architectural and regulatory features of a giant virus. *J Virol.* 2012; 86(16):8821–34. <https://doi.org/10.1128/JVI.00907-12> PMID: 22696644
22. Yanai-Balser GM, Duncan GA, Eudy JD, Wang D, Li X, Agarkova IV, et al. Microarray analysis of paramecium bursaria chlorella virus 1 transcription. *J Virol.* 2010; 84(1):532–42. <https://doi.org/10.1128/JVI.01698-09> PMID: 19828609
23. Blanc G, Mozar M, Agarkova IV, Gurnon JR, Yanai-Balser G, Rowe JM, et al. Deep RNA sequencing reveals hidden features and dynamics of early gene transcription in paramecium bursaria chlorella virus 1. *PLoS One.* 2014; 9(3):10. <https://doi.org/10.1371/journal.pone.0090989> PMID: 24608750
24. Hubalek Z. Protectants used in the cryopreservation of microorganisms. *Cryobiology.* 2003; 46(3):205–29. [https://doi.org/10.1016/s0011-2240\(03\)00046-4](https://doi.org/10.1016/s0011-2240(03)00046-4) PMID: 12818211
25. Benson EE. Cryopreservation of phytodiversity: A critical appraisal of theory practice. *Crit Rev Plant Sci.* 2008; 27(3):141–219. <https://doi.org/10.1080/07352680802202034>
26. Day JG, Watanabe MM, Morris GJ, Fleck RA, McLellan MR. Long-term viability of preserved eukaryotic algae. *J Appl Phycol.* 1997; 9(2):121–7. <https://doi.org/10.1023/a:1007991507314>
27. Taylor R, Fletcher RL. Cryopreservation of eukaryotic algae—a review of methodologies. *J Appl Phycol.* 1998; 10(5):481–501. <https://doi.org/10.1023/a:1008094622412>
28. Rhodes L, Smith J, Tervit R, Roberts R, Adamson J, Adams S, et al. Cryopreservation of economically valuable marine micro-algae in the classes *Bacillariophyceae*, *Chlorophyceae*, *Cyanophyceae*, *Dinophyceae*, *Haptophyceae*, *Prasinophyceae*, and *Rhodophyceae*. *Cryobiology.* 2006; 52(1):152–6. <https://doi.org/10.1016/j.cryobiol.2005.10.003> PMID: 16321370
29. Nakanishi K, Deuchi K, Kuwano K. Cryopreservation of four valuable strains of microalgae, including viability and characteristics during 15 years of cryostorage. *J Appl Phycol.* 2012; 24(6):1381–5. <https://doi.org/10.1007/s10811-012-9790-8>
30. Dunigan DD, Agarkova I. Formulation of MBBM (modified Bold's Basal medium). *protocols.io.* 2016; <https://doi.org/10.17504/protocols.io.etwbepe>

31. Van Etten JL, Burbank DE, Kuczmarski D, Meints RH. Virus-infection of culturable *Chlorella*-like algae and development of a plaque assay. *Science*. 1983; 219(4587):994–6. <https://doi.org/10.1126/science.219.4587.994> PMID: 17817937
32. Dunigan DD, Agarkova I. PBCV-1 virus plaque assay. *protocols.io*. 2016: <https://doi.org/10.17504/protocols.io.estbeen>
33. Coy SR, Alsante A, Wilhelm SW. Long term cryopreservation of chloroviruses by infection of *Chlorella*. *protocols.io*. 2018: <https://doi.org/10.17504/protocols.io.wa2fage>
34. Dunigan DD, Fitzgerald LA, Van Etten JL. Phycodnaviruses: a peek at genetic diversity. *Virus Res*. 2006; 117(1):119–32. <https://doi.org/10.1016/j.virusres.2006.01.024> PMID: 16516998
35. Convention. USP. U.S. Pharmacopeia National Formulary 2018: UPS41-NF36: Nielsen bookata; 2018.
36. Agarkova I, Hertel B, Zhang XZ, Lane L, Tchourbanov A, Dunigan DD, et al. Dynamic attachment of Chlorovirus PBCV-1 to *Chlorella variabilis*. *Virology*. 2014; 466:95–102. <https://doi.org/10.1016/j.virol.2014.07.002> PMID: 25240455
37. Sanjuan R, Nebot MR, Chirico N, Mansky LM, Belshaw R. Viral mutation rates. *J Virol*. 2010; 84(19):9733–48. <https://doi.org/10.1128/JVI.00694-10> PMID: 20660197
38. Filee J, Pouget N, Chandler M. Phylogenetic evidence for extensive lateral acquisition of cellular genes by nucleocytoplasmic large DNA viruses. *BMC Evol Biol*. 2008; 8(320). <https://doi.org/10.1186/1471-2148-8-320> PMID: 19036122
39. Filee J, Siguier P, Chandler M. I am what I eat and I eat what I am: acquisition of bacterial genes by giant viruses. *Trends Genet*. 2007; 23(1):10–5. <https://doi.org/10.1016/j.tig.2006.11.002> PMID: 17109990
40. Cordova A, Deserno M, Gelbart WM, Ben-Shaul A. Osmotic shock and the strength of viral capsids. *Biophys J*. 2003; 85(1):70–4. [https://doi.org/10.1016/S0006-3495\(03\)74455-5](https://doi.org/10.1016/S0006-3495(03)74455-5) PMID: 12829465
41. Roos WH, Ivanovska IL, Evilevitch A, Wuite GJL. Viral capsids: mechanical characteristics, genome packaging and delivery mechanisms. *Cell Mol Life Sci*. 2007; 64(12):1484–97. <https://doi.org/10.1007/s00018-007-6451-1> PMID: 17440680
42. Wulfmeyer T, Polzer C, Hiepler G, Hamacher K, Shoeman R, Dunigan DD, et al. Structural organization of DNA in *Chlorella* viruses. *PLoS One*. 2012; 7(2). <https://doi.org/10.1371/journal.pone.0030133> PMID: 22359540
43. Claverie JM, Abergel C. *Virus Taxonomy Ninth Report of the International Committee on Taxonomy of Viruses*: Elsevier Inc.; 2012. p. 223–8.
44. Wallis C, Melnick JL. Stabilization of enveloped viruses by dimethyl sulfoxide. *J Virol*. 1968; 2(9):953–4 PMID: 4302192
45. Noble RT, Fuhrman JA. Use of SYBR Green I for rapid epifluorescence counts of marine viruses and bacteria. *Aquat Microb Ecol*. 1998; 14(2):113–8. <https://doi.org/10.3354/ame014113>
46. Brussaard CPD, Marie D, Bratbak G. Flow cytometric detection of viruses. *J Virol Methods*. 2000; 85(1–2):175–82. [https://doi.org/10.1016/s0166-0934\(99\)00167-6](https://doi.org/10.1016/s0166-0934(99)00167-6) PMID: 10716350
47. Brussaard CPD. Optimization of procedures for counting viruses by flow cytometry. *Appl Environ Microbiol*. 2004; 70(3):1506–13. <https://doi.org/10.1128/AEM.70.3.1506-1513.2004> PMID: 15006772
48. Agarkova I, Dunigan D, Gurnon J, Greiner T, Barres J, Thiel G, et al. Chlorovirus-mediated membrane depolarization of *Chlorella* alters secondary active transport of solutes. *J Virol*. 2008; 82(24):12181–90. <https://doi.org/10.1128/JVI.01687-08> PMID: 18842725
49. Amaral R, Pereira JC, Pais A, Santos LMA. Is axenicity crucial to cryopreserve microalgae? *Cryobiology*. 2013; 67(3):312–20. <https://doi.org/10.1016/j.cryobiol.2013.09.006> PMID: 24055827
50. Wilhelm SW, Coy SR, Gann ER, Moniruzzaman M, Stough JMA. Standing on the shoulders of giant viruses: five lessons learned about large viruses infecting small eukaryotes and the opportunities they create. *PLoS Pathog*. 2016; 12(8). <https://doi.org/10.1371/journal.ppat.1005752> PMID: 27559742
51. Kerepesi C, Grolmusz V. The "giant virus finder" discovers an abundance of giant viruses in the Antarctic dry valleys. *Arch Virol*. 2017; 162(6):1671–6. <https://doi.org/10.1007/s00705-017-3286-4> PMID: 28247094
52. Legendre M, Bartoli J, Shmakova L, Jeudy S, Labadie K, Adrait A, et al. Thirty-thousand-year-old distant relative of giant icosahedral DNA viruses with a pandoravirus morphology. *Proc Natl Acad Sci USA*. 2014; 111(11):4274–9. <https://doi.org/10.1073/pnas.1320670111> PMID: 24591590
53. Legendre M, Lartigue A, Bertaux L, Jeudy S, Bartoli J, Lescot M, et al. In-depth study of *Mollivirus sibiricum*, a new 30,000-y-old giant virus infecting *Acanthamoeba*. *Proc Natl Acad Sci USA*. 2015; 112(38): E5327–E35. <https://doi.org/10.1073/pnas.1510795112> PMID: 26351664

54. Wilhelm SW, Bird JT, Bonifer KS, Calfee BC, Chen T, Coy SR, et al. A student's guide to giant viruses infecting small eukaryotes: from *Acanthamoeba* to *Zooxanthellae*. *Viruses*. 2017; 9(3). <https://doi.org/10.3390/v9030046> PMID: [28304329](https://pubmed.ncbi.nlm.nih.gov/28304329/)
55. Wilson WH, Gilg IC, Moniruzzaman M, Field EK, Koren S, LeCleir GR, et al. Genomic exploration of individual giant ocean viruses. *ISME J*. 2017; 11(8):1736–45. <https://doi.org/10.1038/ismej.2017.61> PMID: [28498373](https://pubmed.ncbi.nlm.nih.gov/28498373/)