



5-2011

The Role of 5-HT2A and 5-HT2C Receptors in Conditioned Defeat

Marquinta Juvon Lee
mlee4@utk.edu

Follow this and additional works at: https://trace.tennessee.edu/utk_gradthes

Recommended Citation

Lee, Marquinta Juvon, "The Role of 5-HT2A and 5-HT2C Receptors in Conditioned Defeat. " Master's Thesis, University of Tennessee, 2011.
https://trace.tennessee.edu/utk_gradthes/891

This Thesis is brought to you for free and open access by the Graduate School at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Masters Theses by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

To the Graduate Council:

I am submitting herewith a thesis written by Marquinta Juvon Lee entitled "The Role of 5-HT2A and 5-HT2C Receptors in Conditioned Defeat." I have examined the final electronic copy of this thesis for form and content and recommend that it be accepted in partial fulfillment of the requirements for the degree of Master of Arts, with a major in Psychology.

Matthew A. Cooper, Major Professor

We have read this thesis and recommend its acceptance:

Todd Freeberg, Rebecca Prosser

Accepted for the Council:

Carolyn R. Hodges

Vice Provost and Dean of the Graduate School

(Original signatures are on file with official student records.)

**The Role of 5-HT_{2A} and 5-HT_{2C} Receptors in
Conditioned Defeat**

A Thesis Presented for the
Master of Arts
Degree
The University of Tennessee, Knoxville

Marquinta Juvon Lee
May 2011

Copyright © 2011 by Marquinta Lee Harvey

All rights reserved

DEDICATION

To my husband
Brandon M. Harvey

my mother
Regina L. Dorsey

my father
Henry W. Lee

my step-father
Dwayne L. Dorsey

my grandparents
Jasper G. Hatcher
Thelma T. Hatcher
Josephine Lee

and cousins
Aundrea Anderson
Kamaria Anderson
Sharonda Anderson
Katrina Perry 'my ace'

ACKNOWLEDGEMENTS

While writing my thesis at the University of Tennessee, Knoxville I have confirmed my passion for scientific research. I have sincere gratitude to everyone who has helped me throughout my graduate school experience. I am heartily thankful to Dr. Matthew Cooper, whose encouragement, supervision and support from the preliminary to the concluding level enabled me to develop critical thinking skills and an understanding of the subject. I would also like to thank Dr. Todd Freeberg and Dr. Rebecca Prosser for their direction, assistance, and guidance.

I would also like to thank Kathleen Morrison for all of her advice, help with my experiments, and laboratory tasks and Lauren Bader for all of her assistance. Thank you to all the undergraduate students in the Cooper Lab, my pocket hamsters, whom I had the pleasure of working with. Your help was essential to the completion of my research and I am deeply grateful for your contribution.

I would like to thank all the faculty, staff and colleagues in this department and the BCMB department who have helped me through these past couple of years. Their assistance made my tenure at the University of Tennessee, Knoxville an enjoyable one.

Lastly, thank you to my family and friends, namely my Uncle Charles Dorsey, whom have continued to encourage me to pursue my academic goals. I would especially like to thank my husband, parents, grandparents, aunts, uncles, in-laws, and cousins.

ABSTRACT

Previous research indicates that serotonin (5-HT) enhances the acquisition of stress-induced changes in behavior; although it is unclear which serotonin receptors mediate this enhancement. 5-HT₂ receptors are potential candidates because activation at these receptors is associated with increased fear and anxiety. In this study we investigated whether pharmacological treatments targeting 5-HT_{2A} and 5-HT_{2C} receptors modulated the acquisition and expression of conditioned defeat. Conditioned defeat is a social defeat model in Syrian hamsters (*Mesocricetus auratus*) that is characterized by increased submissive and defensive behavior and a loss of territorial aggression following social defeat. In experiment 1, we injected the 5-HT_{2C} receptor agonist mCPP (0.3, 1.0, or 3.0 mg/kg) or vehicle prior to social defeat and tested subjects for conditioned defeat behavior in a social interaction test 24 hours later. In experiment 2, subjects received a social defeat, and 24 hours later we injected mCPP (0.3, 1.0, or 3.0 mg/kg) or vehicle prior to a social interaction test. We found that injection of mCPP increased the expression, but not acquisition, of conditioned defeat. In experiment 3, we injected the 5-HT_{2A} receptor antagonist MDL 11,939 (0.5 or 2.0 mg/kg) or vehicle prior to a social defeat and tested subjects for conditioned defeat behavior. In experiment 4, subjects received a social defeat, and 24 hours later we injected MDL 11,939 (0.5 or 2.0 mg/kg) or vehicle prior to a social interaction test. We found that injection of MDL 11,939 significantly decreased the acquisition, but not expression, of conditioned defeat. These data suggest that pharmacological activation of 5-HT_{2C} receptors

enhances the expression of conditioned defeat, while pharmacological blockade of 5-HT_{2A} receptors impairs the acquisition of conditioned defeat. These data extend other studies indicating that 5-HT signaling at 5-HT_{2A} receptors facilitate memories for aversive events and 5-HT signaling at 5-HT_{2C} receptors enhance stress-induced anxiety.

TABLE OF CONTENTS

Section	Page
Introduction.....	1
Methods.....	6
Results.....	12
Discussion.....	17
List of References.....	24
Vita.....	38

LIST OF FIGURES

Figure	Page
Figure 1.....	33
Figure 2.....	34
Figure 3.....	35
Figure 4.....	36
Figure 5.....	37

INTRODUCTION

Psychosocial stress in humans can lead to a variety of psychiatric disorders, including major depression, acute stress disorder, and post-traumatic stress disorder (Ramboz et al., 1998, Naughton et al., 2000, Middlemiss et al., 2002, Davidson, 2003). The biological basis for stress-related mental illness remains poorly understood. Animal models of fear and anxiety have been used to understand the neural mechanisms underlying these psychiatric disorders. Because most of the stressors that are experienced by humans are social in nature (Brown and Prudo, 1981, Kessler, 1997, Bjorkqvist, 2001), ethologically relevant animal models that examine social conflict are particularly useful for determining how social experience alters the brain and subsequent behavior. Previous research has used physical stressors such as forced swim, foot shock and immobilization test. Although these are potent stressors that activate the stress response, physical stressors activate slightly different neural circuitry compared to psychosocial stressors (Canteras and Blanchard, 2008).

Social defeat is a robust stressor that activates the HPA-axis (Blanchard et al., 1995, Koolhaas et al., 1997). Social defeat also leads to several long-lasting behavioral and physiological changes, such as decreased locomotor activity /exploratory behavior (Meerlo et al., 1996a, Koolhaas et al., 1997, Rygula et al., 2005), changes in circadian rhythmicity (Meerlo et al., 1996a, Meerlo et al., 2002) and altered feeding and body weight (van de Poll et al., 1982, Meerlo et al., 1996b, Bartolomucci et al., 2004, Foster et al., 2006). The behavioral

effects produced by social defeat stress are noticeably similar to symptoms of depression, and many of these effects are reversed with antidepressant treatments (e.g., drugs or controlled sleep deprivation) (Fuchs et al., 1996, Meerlo et al., 1996a, Berton et al., 1999). Social defeat also produces changes in the serotonergic system. Serotonin's (5-HT's) specific role in fear and anxiety-like behavior is mixed. Conflicting data from human and animal research support both an anxiolytic and anxiogenic role for 5-HT (Gordon and Hen, 2004). More research is being conducted on various 5-HT receptors that may be mediating the changes in anxiety behavior.

Siberian and Syrian hamsters have been used as rodent models in circadian rhythms, obesity, and agonistic behavior (Wade and Bartness, 1984a, b). Syrian hamsters are especially useful for studying changes in agonistic behavior because they are solitary aggressive animals that will defend their territory from conspecifics (Nowack and Paradiso 1983). In a laboratory setting, singly housed hamsters defend their territory from intruders who are placed in their home cage (Albers 2002). However if a Syrian hamster loses an aggressive encounter, it will fail to display its' natural territorial aggression in future encounters and instead display submissive and defensive behavior towards a novel intruder (Huhman et al., 2003). This switch in agonistic behavior has been called conditioned defeat and has been used as a model for stress-induced anxiety disorders (Huhman, 2006).

Stressful events are known to increase fear and anxiety. Exposure to a predator is an ethologically relevant stressor that causes an increase in flight,

avoidance, and risk assessment in a mouse defensive test battery (Blanchard et al., 1990, Griebel et al., 1995). Predator odor increases different types of defensive behavior in both mice and rats depending on whether the threat is uncertain, distal, or proximal (Blanchard and Blanchard 2008). For instance, rats perform cautious exploration, such as risk assessment, in novel environments, when danger is certain. When a predator is perceived at a distance, tense and attentive immobility (freezing) ensues. Finally, when a predator is near or in actual contact with the rat, the animal flees whenever possible or otherwise threatens back or even attacks the predator defensively (Blanchard and Blanchard, 1988). Our lab has attempted to differentiate these defensive behaviors in hamsters by quantifying flight as a fear-like response, and stretch attends as an anxiety-like response.

Serotonin (5-HT) is a neurochemical increased during stressful events and known to modulate fear and anxiety. Previous research suggests that disruption of 5-HT is linked to anxiety disorders and serotonergic drugs are used as pharmacological treatment for many anxiety disorders (Owens and Nemeroff, 1998, Ballenger, 1999). The majority of 5-HT neurons that innervate stress-sensitive regions of the forebrain project from the dorsal raphe nucleus (DRN). Increases in 5-HT concentrations in the DRN are associated with exposure to stressful stimuli such as forced swim (Kirby et al., 1995) and foot shock (Yoshioka et al., 1995). 5-HT has been shown to increase anxiety in conflict tests in rats tested in an elevated t-maze (Graeff, 2002). Our lab has shown that social defeat activates 5-HT neurons in the DRN (Cooper et al., 2009). Additionally, we

have shown that blocking 5-HT activity by activating 5-HT_{1A} autoreceptors in the DRN disrupts the acquisition and expression of conditioned defeat (Cooper et al., 2008).

It is unclear which 5-HT receptors in the forebrain facilitate stress-induced changes in fear and anxiety. 5-HT₂ receptors are potential candidates for translating stress-induced increases in 5-HT into increased anxiety-like behavior. 5-HT₂ receptors are postsynaptic, G-protein coupled receptors, that elevate cytosolic Ca⁺⁺ (Conn and Sanders-Bush, 1986). The three subtypes of 5-HT₂ receptors (2A, 2B, and 2C) have different distributions in the brain. While 5-HT_{2B} receptors are found mainly in the periphery, 5-HT_{2A} and 5-HT_{2C} receptors are widely distributed throughout the brain. 5-HT_{2A} receptors occur in high densities in the frontal cortex, piriform cortex, ventro-caudal part of the hippocampus (CA3), medial mammillary nucleus, the pontine nuclei, the motor cranial nerve nuclei in the brainstem, and the ventral horn of the spinal cord (Pompeiano et al., 1994). High densities of 5-HT_{2C} receptors are found in retrosplenial, piriform and entorhinal cortex, anterior olfactory nucleus, lateral septal nucleus, subthalamic nucleus, amygdala, subiculum and ventral part of CA3, lateral habenula, substantia nigra pars compacta, several brainstem nuclei and the whole grey matter of the spinal cord (Pompeiano et al., 1994). 5-HT_{2C} receptors contribute to the expression of fear and anxiety. Pharmacological activation of 5-HT_{2C} receptors has induced panic attacks in humans (Kahn et al., 1988). Administration of a 5-HT_{2C} receptor agonist before testing has been shown to increase the expression of learned helplessness behavior, such as reduced

social exploration, in rats (Strong et al., 2009). 5-HT_{2A} receptors are important for the formation of emotional memories. Injection of 5-HT_{2A} receptor agonists prior to training has been shown to facilitate eye blink conditioning in rabbits (Harvey, 2003).

The purpose of this study was to examine the role of 5-HT_{2A} and 5-HT_{2C} receptors in conditioned defeat. We chose the drug mCPP, a 5-HT_{2C} receptor agonist, because previous research in animal and human studies has shown that mCPP increases anxiety. We choose MDL 11,939, a 5-HT_{2A} receptor antagonist because of its' high affinity for 5-HT_{2A} receptors over other 5-HT receptors. We hypothesized that 5-HT_{2C} receptor activation prior to testing would increase the production of conditioned defeat behavior. Also we hypothesized that 5-HT_{2A} receptor blockade prior to social defeat training would impair the formation of conditioned defeat.

METHODS

Animals

Subjects were adult male Syrian hamsters (*Mesocricetus auratus*) that weighed 130–190 g (3–4 months) at the start of the study, and were individually housed for 10–14 days prior to testing. Older hamsters that weighed 180–200 g (>6 months) were individually housed and used as resident aggressors for social defeat training. Immature hamsters that weighed 90–120 g (2 months) were group-housed (three or four animals per cage) and used as non-aggressive opponents for conditioned defeat testing. All animals were housed in polycarbonate cages (20 cm × 40 cm × 20 cm) with corncob bedding, cotton nesting materials, and wire mesh tops. Animal cages were not changed for at least 1 week prior to testing to allow individuals to scent mark their territory. Animals were housed in a temperature-controlled colony room (20 ± 2 °C) and maintained on a 14:10 h light-dark cycle with food and water available ad libitum. All procedures were approved by the University of Tennessee Animal Care and Use Committee and are in accordance with the US National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Conditioned defeat protocol

During social defeat training subjects experienced either one 15-min or one sub-optimal 10-min aggressive encounter in a resident aggressor's home-cage. The 10-min aggressive encounters were used to avoid a ceiling effect when we expected drug treatment to increase conditioned defeat behavior.

During social defeat training subjects were reliably attacked and defeated by resident aggressors. To standardize the amount of aggression received and the duration of social defeat, timing of aggressive encounters began at the first attack by the resident aggressor, which usually occurred within the first 60 s. During social defeat training we recorded the total duration of aggression and the number of attacks displayed by resident aggressors. No-defeat control subjects did not receive a social defeat. To investigate whether drug treatments affected agonistic behavior in the absence of social defeat experience, we included no-defeat control groups that were exposed to a resident aggressor's empty cage. We performed all training and subsequent testing under red or dim light (< 40 lux) during the first 3 h of the dark phase of the light-dark cycle.

Behavioral testing occurred 24 h after training and consisted of one, 5-min encounter with a novel, non-aggressive opponent in the subject's home cage. Testing sessions were digitally recorded and later scored by researchers blind to the experimental conditions using an ethogram adapted from Albers et al.(2002). A second researcher scored a subset of testing sessions; inter-observer reliability was 91% with a kappa of coefficient of .292. We recorded the total duration of four classes of behavior during the 5-min tests: (a) non-agonistic social (approach, investigate, sniff, and nose touch); (b) nonsocial (locomotion, exploration, self-groom, nest build, and feed); (c) submissive and defensive (flight, avoid, tail up, upright and side defense, full submissive posture, stretch-attend, head flag, attempt to escape from cage); and (d) aggressive (upright and side offense, chase, and attack including bite). For a more detailed analysis of

the subject's agonistic behavior, we also quantified the frequency of flight, stretch-attend, and attack.

Drugs

We dissolved 1-(3-Chlorophenyl)piperazine (mCPP; Sigma-Aldrich) in sterile saline as per previous research (Fox et al. 2008). mCPP is a non-selective 5-HT₂ receptor agonist which shows a preferential affinity at 5-HT_{2C} receptors (Kimura et al., 2009). We dissolved α -phenyl-1-(2-phenylethyl)-4-piperidine methanol (MDL 11,939; Tocris) in sterile saline with 1% of acetic acid and adjusted the pH to 5.5 with NaOH as per previous research (Welsh et al., 1998). MDL 11,939 is a highly selective 5-HT_{2A} receptor antagonist (Welsh et al., 1998). All drugs were administered in a 0.3 ml volume using an intraperitoneal injection (i.p.) with a 1ml syringe.

Experiments 1 and 2: 5-HT_{2C} receptor agonist

We designed experiment 1 to test whether injection of a 5-HT_{2C} receptor agonist would enhance the acquisition of conditioned defeat. We injected mCPP (0.3 mg/kg, N=11; 1.0 mg/kg, N=11; or 3.0 mg/kg, N=11) or vehicle (N = 11) 15 min prior to a 10-min social defeat. Animals were tested for conditioned defeat behavior 24 h later.

We designed experiment 2 to test whether injection of a 5-HT_{2C} receptor agonist would enhance the expression of conditioned defeat. Hamsters received a 10-min social defeat, and 24 hours later we injected mCPP (0.3 mg/kg, N=10;

1.0 mg/kg, N=11; or 3.0 mg/kg, N=10) or vehicle (N=11) 15-minutes prior to conditioned defeat testing. No defeat controls received exposure to a resident aggressor's empty cage during training, and 24 h later we injected mCPP (1.0 mg/kg, N=8) or vehicle (0.0 mg/kg, N=8) 15-minutes prior to conditioned defeat testing.

Experiments 3 and 4: 5-HT_{2A} receptor antagonist

We designed Experiment 3 to test whether injection of a selective 5-HT_{2A} receptor antagonist would reduce the acquisition of conditioned defeat. We injected MDL 11,939 (0.5 mg/kg, N=11; or 2.0 mg/kg, N=10) or vehicle (N=10) 30 min prior to a 15-min social defeat. For no defeat controls, we injected MDL 11,939 (2.0 mg/kg, N=8) or vehicle (0.0 mg/kg, N=9) 30 min prior to exposure to a resident aggressor's empty cage. Animals were tested for conditioned defeat behavior 24 h later as described above.

We designed Experiment 4 to test whether injection of a selective 5-HT_{2A} receptor antagonist would reduce the expression of conditioned defeat. Hamsters receive a 15-min social defeat, and 24 h later we injected MDL 11,939 (0.5 mg/kg, N=11; or 2.0 mg/kg, N=10) or vehicle (N=10) 30 min prior conditioned defeat testing. Likewise, no defeat controls received exposure to a resident aggressor's empty cage during training, and 24 h later we injected MDL 11,939 (2.0 mg/kg, N=8) or vehicle (0.0 mg/kg, N=8) 30 min prior to conditioned defeat testing.

Data analysis

Several subjects were not included in statistical analysis because of difficulties with the conditioned defeat protocol. Nine animals were excluded because they were attacked by intruders during testing, two were excluded because of insufficient defeats, and 25 animals were excluded because two cohorts of subjects failed to show conditioned defeat behavior. The 2 excluded cohorts were bred in the Walters Life Science building at the University of Tennessee, Knoxville, born in June and July, and had no obvious reason for not showing conditioned defeat. Our lab is currently investigating individual variation in our subjects to explain why Syrian hamsters can widely vary in their display of conditioned defeat behavior. We analyzed the data with both cohorts included, dropping all animals that did not show conditioned defeat, and by dropping cohorts when vehicle control subjects did not show conditioned defeat and the statistical results were similar for all three types of analyses.

For social defeat training, we analyzed the total duration of aggression received by subjects and the frequency of attacks received by subjects. For conditioned defeat testing, we separately analyzed the total durations of submissive and defensive, non-agonistic social, nonsocial, and aggressive behavior, as well as the frequencies of attack, flight, and stretch-attend posture. Conditioned defeat data were analyzed using two-way ANOVAs with defeat experience (defeat; no defeat) and drug dose as independent variables. To investigate a dose-response relationship for drug treatments in defeated subjects we performed one-way ANOVAs with Tukey or LSD post-hoc tests. We used t-tests to further investigate the effect of drug treatment in no defeat controls. All

statistical tests were two-tailed and the alpha level was $p < 0.05$. Data are presented as mean \pm S.E.

RESULTS

Experiment 1: mCPP and acquisition of conditioned defeat

Injection of mCPP prior to social defeat did not significantly alter the acquisition of conditioned defeat (Fig 1). mCPP treatment did not significantly alter the duration of submissive and defensive behavior ($F_{(3,43)} = 0.279$, $p = 0.840$). Likewise, animals injected with 0.3, 1.0 or 3.0 mg/kg of mCPP did not show significant changes in frequency of flight ($F_{(3,43)} = 1.177$, $p = 0.330$) or stretch-attend postures ($F_{(3,43)} = .320$, $p = 0.811$) compared to vehicle controls (Table 1). Injection of mCPP prior to social defeat did not alter the duration of non-agonistic social behavior ($F_{(3,43)} = .390$, $p = 0.760$), nonsocial behavior ($F_{(3,43)} = .346$, $p = 0.792$), or aggressive behavior ($F_{(3,43)} = 1.306$, $p = 0.286$).

Injection of mCPP prior to social defeat training did not alter the amount of aggression resident aggressors directed toward subjects. Vehicle controls received 200.5 s (± 33.3) of aggression during social defeat and individuals injected with 0.3, 1.0, or 3.0 mg/kg of mCPP received 250.5 s (± 28.9), 291.5 (± 24.6) and 296.0 s (± 28.4) of aggression, respectively ($F_{(3,43)} = 0.55$, $p = 0.567$). Vehicle controls received 12.3 (± 1.1) attacks during social defeat and individuals injected with 0.3, 1.0 and 3.0 mg/kg of mCPP received 10.8 (± 1.6), 13.7 (± 2.1) and 12.7 (± 1.9), attacks, respectively ($F_{(3,43)} = 1.41$, $p = 0.796$).

Experiment 2: mCPP and expression of conditioned defeat

Injection of mCPP prior to behavioral testing dose dependently increased the expression of conditioned defeat (Fig 2). A nearly significant drug by defeat

interaction ($F_{(1,36)} = 4.116, p = 0.051$) was found for the duration of submissive and defensive behavior. This result indicates that 1.0 mg/kg of mCPP increased submissive and defensive behavior in defeated subjects, but not in no defeat controls. Also, a one-way ANOVA on defeated subjects showed a significant increase in submissive and defensive behavior at 1.0 mg/kg mCPP ($F_{(3,40)} = 4.204, p = 0.012$, Tukey, $p = .011$) but not at 3.0 mg/kg (Tukey, $p = ???$). However, mCPP did not significantly change the frequency of flight ($F_{(1,36)} = .216, p = 0.645$) or stretch-attend postures ($F_{(1,36)} = .430, p = 0.517$) compared to vehicle controls (Table 2). To measure the selectivity of mCPP's effect on conditioned defeat behavior we also quantified three other classes of behavior, non-agonistic social, nonsocial, aggressive behavior. We expect a selective effect of mCPP on submissive/defensive behavior but not on the other classes of behavior. Our results were as expected, injection of mCPP prior to behavioral testing did not alter the duration of other classes of behavior such as non-agonistic social ($F_{(1,36)} = 0.081, p = 0.778$), nonsocial ($F_{(1,36)} = 1.270, p = 0.268$), or aggression ($F_{(1,36)} = 0.515, p = 0.478$).

No defeat controls did not show greater aggression ($F_{(1,36)} = .515, p = 0.478$) but did show less submissive and defensive behavior ($F_{(1,36)} = 17.382, p < 0.001$) compared to defeated subjects (Fig. 2). Also, injection of mCPP in no defeat control animals did not alter the duration of submissive and defensive ($t = -1.260, p = 0.248$), aggressive ($t = 1.000, p = .351$), non-agonistic social ($t = .476, p = .649$), or nonsocial behavior ($t = -.643, p = .541$). Similarly, no defeat controls injected with mCPP or vehicle did not significantly differ in the number of

attacks initiated during conditioned defeat testing ($t = 1.000$, $p = .351$; Table 2).

Experiment 3: MDL 11,939 and acquisition of conditioned defeat

Injection of MDL 11,939 prior to social defeat dose dependently decreased the acquisition of conditioned defeat (Fig 3). We found significant drug by defeat interaction ($F_{(1,36)} = 4.793$, $p = 0.036$) was found for the duration of submissive and defensive behavior. This result indicates that 2.0 mg/kg of MDL 11,939 reduced submissive and defensive behavior in defeated subjects, but not in no defeat controls. However, a one-way ANOVA on defeated subjects showed a marginally significant decrease in submissive and defensive behavior at 2.0 mg/kg ($F_{(1,30)} = 2.594$, $p = 0.093$, LSD, $p = .05$). MDL 11,939 did not significantly change the frequency of flight ($F_{(1,36)} = .378$, $p = 0.543$) or stretch-attend postures ($F_{(1,36)} = .757$, $p = 0.391$) compared to vehicle controls (Table 3). Also, injection of MDL 11,939 prior to social defeat did not alter the duration of other classes of behavior such as non-agonistic social ($F_{(1,36)} = 0.204$, $p = 0.661$), nonsocial ($F_{(1,36)} = 0.012$, $p = 0.912$), or aggression ($F_{(1,36)} = 0.196$, $p = 0.661$).

No defeat controls showed greater aggression ($F_{(1,36)} = 9.412$, $p = 0.004$) and less submissive and defensive behavior at testing ($F_{(1,36)} = 17.945$, $p = 0.000$) compared to defeated subjects (Fig. 3). However, injection of MDL 11,939 in no defeat control animals did not alter the duration of submissive and defensive ($t_{(1,36)} = 1.174$, $p = 0.279$), aggressive ($t_{(1,36)} = .417$, $p = .689$), non-agonistic social ($t_{(1,36)} = -.361$, $p = .729$), or nonsocial behavior ($t_{(1,36)} = -.203$, $p = .845$). Similarly, no defeat controls injected with MDL 11,939 or vehicle did not

significantly differ in the number of attacks displayed during conditioned defeat testing ($t_{(1,36)} = .403$, $p = .699$; Table 3).

Injection of MDL 11,939 did not alter the level of aggression subjects received during social defeat training. Vehicle controls received 307 s (± 57.9) of aggression during social defeat and individuals injected with 0.5 or 2.0 mg/kg of MDL 11,939 received 303 s (± 47.5) and 277.3 s (± 53.3), respectively ($F_{(2,28)} = 0.095$, $p = 0.909$). Vehicle controls received 21.7 (± 1.2) attacks during social defeat and individuals injected with 0.5 or 2.0 mg/kg of MDL 11,939 received 19.5 (± 2.5) and 16.4 (± 3.3), attacks, respectively ($F_{(2,28)} = 0.857$, $p = 0.436$).

Experiment 4: MDL 11,939 and expression of conditioned defeat

Injection of MDL 11,939 prior to behavioral testing did not significantly alter the expression of conditioned defeat (Fig 4). We did not find a significant drug by defeat interaction for the duration of submissive and defensive behavior ($F_{(1,37)} = 0.98$, $p = 0.757$). Also, a one-way ANOVA on defeated subjects did not reveal a significant difference in submissive and defensive behavior ($F_{(1,31)} = 0.248$, $p = 0.782$). Likewise, animals injected with 0.5 or 2.0 mg/kg of MDL 11,939 did not show significant changes in frequency of flight ($F_{(1,37)} = 0.287$, $p = 0.837$) or stretch-attend postures ($F_{(1,37)} = 1.400$, $p = 0.245$) compared to vehicle controls (Table 4). Injection of MDL 11,939 prior to conditioned defeat testing did not produce changes in the duration of non-agonistic social behavior ($F_{(1,37)} = 1.227$, $p = 0.276$) or aggressive behavior

($F_{(1,37)} = 0.314, p = 0.579$). However we found a significant drug by defeat interaction for nonsocial behavior ($F_{(1,37)} = 4.672, p = 0.038$), but a one-way ANOVA on defeated subjects did not show a significant effect of drug treatment on nonsocial behavior ($F_{(2,31)} = 2.619, p = 0.090$). The modest increase in nonsocial behavior appears related to increased cage climbing, nest building, and self-grooming and is not directly attributed to drug-induced hyper-locomotion.

No defeat controls showed elevated aggression ($F_{(1,37)} = 5.309, p = 0.027$) and reduced submissive and defensive behavior ($F_{(1,37)} = 10.354, p = 0.003$) compared to defeated subjects (Fig. 4). Also, injection of MDL 11,939 in no defeat control subjects did not alter the duration of aggression ($t_{(1,36)} = 1.052, p = .328$), submission ($t_{(1,36)} = .037, p = .971$), non-agonistic social ($t_{(1,36)} = -.361, p = .729$), or nonsocial behavior ($t_{(1,36)} = -.203, p = .845$). Similarly, no defeat controls attacked non-aggressive opponents at testing more often than did defeat animals ($F_{(1,37)} = 5.309, p = 0.027$), although MDL 11,939 injection did not alter frequency of attacks ($t_{(1,36)} = 1.055, p = .326$; Table 4).

DISCUSSION

In each experiments we found an effect of social defeat on conditioned defeat behavior. Specifically, defeated animals show increased submissive and defensive behavior and decrease in aggressive behavior, when compared to no defeat control subjects. Also, in all experiments pharmacological manipulations did not produce a significant change in the behavior of no defeat subjects. Because the effects of drug treatment were limited to defeated subjects, we concluded the prior psychosocial stress is required for the 5-HT₂ ligands used here to modulate agonistic behavior. Our study shows that administration of mCPP, a nonselective 5-HT_{2C} receptor agonist, increases the expression but not acquisition of conditioned defeat behavior. These results suggest that activation of 5-HT_{2C} receptors are important for the production of submissive and defensive behavior at testing but not the development of conditioned defeat behavior. We found that injection of MDL 11,939, a selective 5-HT_{2A} receptor antagonist, reduces the acquisition but not expression of conditioned defeat behavior. These results suggest that 5-HT_{2A} receptor blockade impairs the development of conditioned defeat but is not critical for the production of submissive and defensive behavior at testing. Together these data suggest that 5-HT may act at 5-HT_{2C} and 5-HT_{2A} receptors to facilitate the expression and acquisition of conditioned defeat, respectively. In sum, these results support our overarching hypothesis that defeat-induced increases in serotonin act at 5-HT₂ receptors in the forebrain to promote conditioned defeat behavior.

Administration of mCPP exacerbates panic attacks in humans with panic disorder causing behavioral effects such as increased anxiety, depression and panic attacks (Kahn et al., 1988). Also, mCPP increases the expression of anxiety-like behavior in several animal models including the social interaction test (Bagdy et al., 2001), light/dark transition box test (Bilkei-Gorzo et al., 1998), and open field test (Campbell and Merchant, 2003). Although mCPP often is used as a 5-HT_{2C} receptor agonist, the drug pharmacology is complex. mCPP activates several other receptors and binds with equal affinity to 5-HT_{2C} and 5-HT_{2B} receptors. It binds to 5-HT_{2C} receptors with a ten-fold greater selectivity than 5-HT_{2A} receptors and a two fold greater selectivity than 5-HT_{1A} (Roth et al., 1995, Campbell and Merchant, 2003). The non-selective binding of mCPP at 5-HT receptors could explain the non-linear dose response curve in our results. We found that 1.0 mg/kg of mCPP increased conditioned defeat, whereas 3.0 mg/kg was less effective. Our data is consistent with other research showing inverted U-shaped dose response curves for mCPP effects. For example, mCPP treatment increases anxiety in an open field test at doses between 3 and 300 pmol but not at 3000 pmol (Campbell and Merchant, 2003). One possibility is that mCPP fails to increase conditioned defeat at high doses because it binds to other receptors, such as the 5-HT_{1A} receptor. This possibility would be consistent with our previous finding that activation of 5-HT_{1A} receptors in the basolateral amygdala (BLA) decreases conditioned defeat (Morrison et al., 2011). In a learned helplessness model, another 5-HT_{2C} receptor agonist, CP-809101, has been shown to impair escape behavior in the absence of prior stress (Strong et al.,

2009). Unlike with Strong et al. (2009), activation of 5-HT_{2C} receptors in our study did not create conditioned defeat behavior in our no defeat subjects. Thus, activation of 5-HT_{2C} receptors appears to interact with social defeat to enhance the display of submissive and defensive behavior, however it does not produce conditioned defeat itself.

We also quantified flees and stretch attends in an attempt to differentiate fear and anxiety. Threat stimuli, like predator odor, increases different types of defensive behavior in both mice and rats. These defensive behaviors have been divided into fear-like responses, which include escape behavior, and anxiety-like behavior, which include risk assessment (Blanchard and Blanchard 2008). In our animals we used flees to represent escape behavior and stretch attends to represent risk assessment behavior. MDL and mCPP failed to significantly alter the frequency of flees or stretch attends. Because there were no significant changes in flee or stretch attend behavior our study was unable to differentiate the effect of 5-HT₂ receptors on this aspect of fear and anxiety. Future research will require us to modify our ethogram to more carefully address fearful and anxious types of behavior.

Several brain regions may underlie the effect of mCPP on the expression of conditioned defeat. Brain regions such as the bed nucleus of the stria terminalis (BNST) and central nucleus of the amygdala (CeA), have been implicated in the link between 5-HT_{2C} receptors and the expression of anxiety and fear-like responses. 5-HT_{2C} receptor knock-out mice show reduced c-Fos immunoreactivity in the BNST and CeA following exposure to an anxiety-

provoking stimulus (Heisler et al., 2007). Systemic mCPP administration has been shown to increase the expression of c-Fos in the anterolateral BNST (Singewald et al., 2003). Also, the anxiogenic effects of mCPP have been linked to 5-HT_{2A/2C} receptors expressed by BNST projection neurons (Hammack et al., 2009). The BLA is another key brain region because it contains 5-HT_{2C} receptor protein (Pompeiano et al., 1994) and plays a critical role in the expression of conditioned defeat. Others have found that 5-HT_{2C} receptor activation within the BLA causes acute fear-like responses in an open-field test (Campbell and Merchant, 2003). Similarly, 5-HT_{2C} receptor activation in the BLA reduces social exploration in the learned helplessness model (Christianson et al., 2010).

Pharmacological treatments targeting 5-HT_{2A} receptors have been shown to modulate several types of learning including spatial, emotional, and associative learning in several species (Harvey et al., 1982, Alhaider et al., 1993, Williams et al., 2002). Activation of 5-HT_{2A} receptors by lysergic acid diethylamide, LSD (Gimpl et al., 1979, Siegel et al., 1996), 2,5-dimethoxy-4-methylamphetamine, DOM (Harvey et al., 1982), 3,4-methylenedioxyamphetamine, MDA (Romano et al., 1991), and methylenedioxymethamphetamine, MDMA (Romano and Harvey, 1994) enhances eye blink conditioning in rabbits. Also blockade of 5-HT_{2A} receptors with ritanserin (Welsh et al., 1998), mianserin (Romano et al., 1991), MDL 11,939 (Welsh et al., 1998), and pizotifen (Ginn and Powell, 1986) has been shown to impair eye blink conditioning in rabbits. These studies suggest that 5-HT_{2A}

receptor activation facilitates, while 5-HT_{2A} receptor blockade disrupts eye blink conditioning. The 5-HT_{2A} receptors' role in modulating aversive learning is not limited to eye blink conditioning in rabbits; other animal and human studies have shown that activation or blockade of 5-HT_{2A} receptors modulates the formation of memories for aversive events. For example, the acquisition of active avoidance was enhanced in rats using quipazine, a 5-HT agonist, and was blocked by ketanserin, a 5-HT_{2A/2C} antagonist, suggesting that the enhanced formation of active avoidance was facilitated by 5-HT_{2A} receptors (Alhaider et al., 1993). Similarly, cyproheptadine, a 5-HT_{2A/2C} receptor antagonist, impaired the acquisition of active avoidance (Titov et al., 1983, Ma and Yu, 1993). In humans, injection of ritanserin has been shown to impair learning in an aversive classical conditioning test (Hensman et al., 1991).

Consistent with the research on classical conditioning and active avoidance, our results support a role for 5-HT_{2A} receptors in the acquisition of stress-related memories. Our results indicate that blockade of 5-HT_{2A} receptors prior to social defeat impairs the acquisition of conditioned defeat behavior. MDL 11,939 may impair the acquisition of conditioned defeat by acting in several brain regions that have been implicated in emotional memories. 5-HT_{2A} receptors in the hippocampus and frontal cortex have been implicated in eye blink conditioning (Takehara et al., 2003). Importantly, neural transmission in the hippocampus and prefrontal cortex are necessary for the development of conditioned defeat. Previous research has shown that inactivation of the hippocampus using muscimol disrupted the acquisition of conditioned defeat

(Markham et al., 2010) and inactivation of medial prefrontal cortex impairs conditioned defeat resistance in dominant hamsters (Morrison and Cooper, 2010. Online). The BLA is another candidate brain region for mediating the effect of MDL 11,939 on the development of conditioned defeat. We have previously shown that Syrian hamsters have 5-HT_{2A} receptors in the BLA (Morrison et al., 2011), and neural plasticity in the BLA is critical for the development of conditioned defeat (Jasnow et al., 2005, Markham et al., 2010, Day et al., 2011). Also, 5-HT_{2A} receptors are present on GABAergic interneurons and glutamatergic pyramidal cells in the BLA of rats (McDonald and Mascagni, 2007). One possible explanation for our results is that MDL 11,939 may preferentially block 5-HT_{2A} receptors on BLA glutamatergic cells and thereby impair the development of conditioned defeat. Interesting, serotonergic input can desensitize 5-HT_{2A} receptors in vitro (Roth et al., 1995). Thus, another possibility is that MDL 11,939 might prevent the desensitization at 5-HT_{2A} receptors on GABAergic neurons in the BLA and thus enable serotonergic inhibition of the BLA pyramidal neurons at testing (Fig. 5).

These data extend our understanding the role of 5-HT in the acquisition and expression of conditioned defeat. We have previously shown that enhancing 5-HT signaling in the DRN increases conditioned defeat (Cooper et al., 2008). It was unclear which post-synaptic receptors mediated this increase in conditioned defeat. The current study indicates that the 5-HT₂ receptors play a key role in facilitating conditioned defeat. Our data are consistent with previous research suggesting that activation of 5-HT_{2C} receptors is important for the expression of

anxiety-like behavior. While the role of 5-HT_{2A} receptors in the acquisition of anxiety-like behavior is unclear and our findings provide a novel example of the role of 5-HT_{2A} receptors in the formation of anxiety-like behavior. This study builds upon our working model of mechanisms by which 5-HT can modulate the acquisition and expression of conditioned defeat (see Fig. 5). In sum, our results indicate that conditioned defeat is an elegant model for investigating 5-HT's role in anxiety disorders.

REFERENCES

References

- Alhaider AA, Ageel AM, Ginawi OT (1993) The quipazine- and TFMPP-increased conditioned avoidance response in rats: role of 5HT_{1C}/5-HT₂ receptors. *Neuropharmacology* 32:1427-1432.
- Bagdy G, Graf M, Anheuer ZE, Modos EA, Kantor S (2001) Anxiety-like effects induced by acute fluoxetine, sertraline or m-CPP treatment are reversed by pretreatment with the 5-HT_{2C} receptor antagonist SB-242084 but not the 5-HT_{1A} receptor antagonist WAY-100635. *Int J Neuropsychopharmacol* 4:399-408.
- Ballenger JC (1999) Current treatments of the anxiety disorders in adults. *Biol Psychiatry* 46:1579-1594.
- Bartolomucci A, Pederzani T, Sacerdote P, Panerai AE, Parmigiani S, Palanza P (2004) Behavioral and physiological characterization of male mice under chronic psychosocial stress. *Psychoneuroendocrinology* 29:899-910.
- Berton O, Durand M, Aguerre S, Mormede P, Chaouloff F (1999) Behavioral, neuroendocrine and serotonergic consequences of single social defeat and repeated fluoxetine pretreatment in the Lewis rat strain. *Neuroscience* 92:327-341.
- Bilkei-Gorzo A, Gyertyan I, Levay G (1998) mCPP-induced anxiety in the light-dark box in rats--a new method for screening anxiolytic activity. *Psychopharmacology (Berl)* 136:291-298.
- Bjorkqvist K (2001) Social defeat as a stressor in humans. *Physiol Behav* 73:435-442.
- Blanchard DC, Blanchard RJ (1988) Ethoexperimental approaches to the biology of emotion. *Annu Rev Psychol* 39:43-68.
- Blanchard DC, Blanchard RJ, Tom P, Rodgers RJ (1990) Diazepam changes risk assessment in an anxiety/defense test battery. *Psychopharmacology (Berl)* 101:511-518.
- Blanchard DC, Spencer RL, Weiss SM, Blanchard RJ, McEwen B, Sakai RR (1995) Visible burrow system as a model of chronic social stress: behavioral and neuroendocrine correlates. *Psychoneuroendocrinology* 20:117-134.
- Brown GW, Prudo R (1981) Psychiatric disorder in a rural and an urban population: 1. Aetiology of depression. *Psychol Med* 11:581-599.
- Campbell BM, Merchant KM (2003) Serotonin 2C receptors within the basolateral amygdala induce acute fear-like responses in an open-field environment. *Brain Res* 993:1-9.
- Canteras NS, Blanchard DC (2008) A behavioral and neural systems comparison of unconditioned and conditioned defensive behavior. In: *Handbook of Anxiety and Fear*, vol. 17 (Blanchard, D. C. et al., eds), pp 141 - 156 Amsterdam: Elsevier.
- Christianson JP, Ragole T, Amat J, Greenwood BN, Strong PV, Paul ED, Fleshner M, Watkins LR, Maier SF (2010) 5-hydroxytryptamine 2C receptors in the

- basolateral amygdala are involved in the expression of anxiety after uncontrollable traumatic stress. *Biol Psychiatry* 67:339-345.
- Conn PJ, Sanders-Bush E (1986) Regulation of serotonin-stimulated phosphoinositide hydrolysis: relation to the serotonin 5-HT₂ binding site. *J Neurosci* 6:3669-3675.
- Cooper MA, Grober MS, Nicholas CR, Huhman KL (2009) Aggressive encounters alter the activation of serotonergic neurons and the expression of 5-HT_{1A} mRNA in the hamster dorsal raphe nucleus. *Neuroscience* 161:680-690.
- Cooper MA, McIntyre KE, Huhman KL (2008) Activation of 5-HT_{1A} autoreceptors in the dorsal raphe nucleus reduces the behavioral consequences of social defeat. *Psychoneuroendocrinology* 33:1236-1247.
- Davidson JR (2003) Treatment of posttraumatic stress disorder: the impact of paroxetine. *Psychopharmacol Bull* 37 Suppl 1:76-88.
- Day DE, Cooper MA, Markham CM, Huhman KL (2011) NR2B subunit of the NMDA receptor in the basolateral amygdala is necessary for the acquisition of conditioned defeat in Syrian hamsters. *Behav Brain Res* 217:55-59.
- Foster MT, Solomon MB, Huhman KL, Bartness TJ (2006) Social defeat increases food intake, body mass, and adiposity in Syrian hamsters. *Am J Physiol Regul Integr Comp Physiol* 290:R1284-1293.
- Fuchs E, Kramer M, Hermes B, Netter P, Hiemke C (1996) Psychosocial stress in tree shrews: clomipramine counteracts behavioral and endocrine changes. *Pharmacol Biochem Behav* 54:219-228.
- Gimpl MP, Gormezano I, Harvey JA (1979) Effects of LSD on learning as measured by classical conditioning of the rabbit nictitating membrane response. *J Pharmacol Exp Ther* 208:330-334.
- Ginn SR, Powell DA (1986) Pizotifen (BC-105) attenuates orienting and Pavlovian heart rate conditioning in rabbits. *Pharmacol Biochem Behav* 24:677-685.
- Gordon JA, Hen R (2004) The serotonergic system and anxiety. *Neuromolecular Med* 5:27-40.
- Graeff FG (2002) On serotonin and experimental anxiety. *Psychopharmacology (Berl)* 163:467-476.
- Griebel G, Blanchard DC, Jung A, Lee JC, Masuda CK, Blanchard RJ (1995) Further evidence that the mouse defense test battery is useful for screening anxiolytic and panicolytic drugs: effects of acute and chronic treatment with alprazolam. *Neuropharmacology* 34:1625-1633.
- Hammack SE, Guo JD, Hazra R, Dabrowska J, Myers KM, Rainnie DG (2009) The response of neurons in the bed nucleus of the stria terminalis to serotonin: implications for anxiety. *Prog Neuropsychopharmacol Biol Psychiatry* 33:1309-1320.
- Harvey JA (2003) Role of the serotonin 5-HT_{2A} receptor in learning. *Learn Mem* 10:355-362.
- Harvey JA, Gormezano I, Cool VA (1982) Effects of d-lysergic acid diethylamide, d-2-bromolysergic acid diethylamide, dl-2,5-dimethoxy-4-methylamphetamine and d-amphetamine on classical conditioning of the rabbit nictitating membrane response. *J Pharmacol Exp Ther* 221:289-294.

- Heisler LK, Zhou L, Bajwa P, Hsu J, Tecott LH (2007) Serotonin 5-HT(2C) receptors regulate anxiety-like behavior. *Genes Brain Behav* 6:491-496.
- Hensman R, Guimaraes FS, Wang M, Deakin JF (1991) Effects of ritanserin on aversive classical conditioning in humans. *Psychopharmacology (Berl)* 104:220-224.
- Huhman KL (2006) Social conflict models: can they inform us about human psychopathology? *Horm Behav* 50:640-646.
- Huhman KL, Solomon MB, Janicki M, Harmon AC, Lin SM, Israel JE, Jasnow AM (2003) Conditioned defeat in male and female Syrian hamsters. *Horm Behav* 44:293-299.
- Jasnow AM, Shi C, Israel JE, Davis M, Huhman KL (2005) Memory of social defeat is facilitated by cAMP response element-binding protein overexpression in the amygdala. *Behav Neurosci* 119:1125-1130.
- Kahn RS, Wetzler S, van Praag HM, Asnis GM, Strauman T (1988) Behavioral indications for serotonin receptor hypersensitivity in panic disorder. *Psychiatry Res* 25:101-104.
- Kessler RC (1997) The effects of stressful life events on depression. *Annu Rev Psychol* 48:191-214.
- Kimura A, Stevenson PL, Carter RN, Maccoll G, French KL, Paul Simons J, Al-Shawi R, Kelly V, Chapman KE, Holmes MC (2009) Overexpression of 5-HT2C receptors in forebrain leads to elevated anxiety and hypoactivity. *Eur J Neurosci* 30:299-306.
- Kirby LG, Allen AR, Lucki I (1995) Regional differences in the effects of forced swimming on extracellular levels of 5-hydroxytryptamine and 5-hydroxyindoleacetic acid. *Brain Res* 682:189-196.
- Koolhaas JM, De Boer SF, De Rutter AJ, Meerlo P, Sgoifo A (1997) Social stress in rats and mice. *Acta Physiol Scand Suppl* 640:69-72.
- Ma TC, Yu QH (1993) Effect of 20(S)-ginsenoside-Rg2 and cyproheptadine on two-way active avoidance learning and memory in rats. *Arzneimittelforschung* 43:1049-1052.
- Markham CM, Taylor SL, Huhman KL (2010) Role of amygdala and hippocampus in the neural circuit subserving conditioned defeat in Syrian hamsters. *Learn Mem* 17:109-116.
- McDonald AJ, Mascagni F (2007) Neuronal localization of 5-HT type 2A receptor immunoreactivity in the rat basolateral amygdala. *Neuroscience* 146:306-320.
- Meerlo P, De Boer SF, Koolhaas JM, Daan S, Van den Hoofdakker RH (1996a) Changes in daily rhythms of body temperature and activity after a single social defeat in rats. *Physiol Behav* 59:735-739.
- Meerlo P, Overkamp GJ, Daan S, Van Den Hoofdakker RH, Koolhaas JM (1996b) Changes in Behaviour and Body Weight Following a Single or Double Social Defeat in Rats. *Stress* 1:21-32.
- Meerlo P, Sgoifo A, Turek FW (2002) The effects of social defeat and other stressors on the expression of circadian rhythms. *Stress* 5:15-22.

- Middlemiss DN, Price GW, Watson JM (2002) Serotonergic targets in depression. *Curr Opin Pharmacol* 2:18-22.
- Morrison KE, Cooper MA (2010. Online) A role for the ventromedial prefrontal cortex in resistance to conditioned defeat. Program Number 597.22. In: 2010 Neuroscience Meeting Planner San Diego, CA: Society for Neuroscience.
- Morrison KE, Swallows CL, Cooper MA (2011) Effects of dominance status on conditioned defeat and expression of 5-HT_{1A} and 5-HT_{2A} receptors. *Physiol Behav*.
- Naughton M, Mulrooney JB, Leonard BE (2000) A review of the role of serotonin receptors in psychiatric disorders. *Hum Psychopharmacol* 15:397-415.
- Owens MJ, Nemeroff CB (1998) The serotonin transporter and depression. *Depress Anxiety* 8 Suppl 1:5-12.
- Pompeiano M, Palacios JM, Mengod G (1994) Distribution of the serotonin 5-HT₂ receptor family mRNAs: comparison between 5-HT_{2A} and 5-HT_{2C} receptors. *Brain Res Mol Brain Res* 23:163-178.
- Ramboz S, Oosting R, Amara DA, Kung HF, Blier P, Mendelsohn M, Mann JJ, Brunner D, Hen R (1998) Serotonin receptor 1A knockout: an animal model of anxiety-related disorder. *Proc Natl Acad Sci U S A* 95:14476-14481.
- Romano AG, Bormann NM, Harvey JA (1991) A unique enhancement of associative learning produced by methylenedioxyamphetamine. *Behav Pharmacol* 2:225-231.
- Romano AG, Harvey JA (1994) MDMA enhances associative and nonassociative learning in the rabbit. *Pharmacol Biochem Behav* 47:289-293.
- Roth BL, Palvimaki EP, Berry S, Khan N, Sachs N, Uluer A, Choudhary MS (1995) 5-Hydroxytryptamine_{2A} (5-HT_{2A}) receptor desensitization can occur without down-regulation. *J Pharmacol Exp Ther* 275:1638-1646.
- Rygula R, Abumaria N, Flugge G, Fuchs E, Ruther E, Havemann-Reinecke U (2005) Anhedonia and motivational deficits in rats: impact of chronic social stress. *Behav Brain Res* 162:127-134.
- Siegel BW, Freedman J, Vaal MJ, Baron BM (1996) Activities of novel aryloxyalkylimidazolines on rat 5-HT_{2A} and 5-HT_{2C} receptors. *Eur J Pharmacol* 296:307-318.
- Singewald N, Salchner P, Sharp T (2003) Induction of c-Fos expression in specific areas of the fear circuitry in rat forebrain by anxiogenic drugs. *Biol Psychiatry* 53:275-283.
- Strong PV, Greenwood BN, Fleshner M (2009) The effects of the selective 5-HT_{2C} receptor antagonist SB 242084 on learned helplessness in male Fischer 344 rats. *Psychopharmacology (Berl)* 203:665-675.
- Takehara K, Kawahara S, Kirino Y (2003) Time-dependent reorganization of the brain components underlying memory retention in trace eyeblink conditioning. *J Neurosci* 23:9897-9905.
- Titov SA, Shamakina I, Ashmarin IP (1983) [Effect of lysyl vasopressin and vasotocin on a disorder of the conditioned avoidance reaction by a serotonin receptor blockader]. *Biull Eksp Biol Med* 95:31-33.

- van de Poll NE, Smeets J, van Oyen HG, van der Zwan SM (1982) Behavioral consequences of agonistic experience in rats: sex differences and the effects of testosterone. *J Comp Physiol Psychol* 96:893-903.
- Wade GN, Bartness TJ (1984a) Effects of photoperiod and gonadectomy on food intake, body weight, and body composition in Siberian hamsters. *Am J Physiol* 246:R26-30.
- Wade GN, Bartness TJ (1984b) Seasonal obesity in Syrian hamsters: effects of age, diet, photoperiod, and melatonin. *Am J Physiol* 247:R328-334.
- Welsh SE, Romano AG, Harvey JA (1998) Effects of serotonin 5-HT(2A/2C) antagonists on associative learning in the rabbit. *Psychopharmacology (Berl)* 137:157-163.
- Williams GV, Rao SG, Goldman-Rakic PS (2002) The physiological role of 5-HT2A receptors in working memory. *J Neurosci* 22:2843-2854.
- Yoshioka M, Matsumoto M, Togashi H, Saito H (1995) Effects of conditioned fear stress on 5-HT release in the rat prefrontal cortex. *Pharmacol Biochem Behav* 51:515-519.

APPENDICES

APPENDIX A

Table 1.

Behavior	0.0 mg/kg	0.3 mg/kg	1.0 mg/kg	3.0 mg/kg	p value
Flee	.455 ± .366	.091 ± .091	1.091 ± .732	.182 ± .122	> .05
Stretch Attend	.364 ± .203	.364 ± .152	.273 ± .141	.545 ± .287	> .05
Attack	.000 ± .000	.000 ± .000	.000 ± .000	.545 ± .455	> .05

The frequencies of flee, stretch attend, and attack (mean ± SE) during conditioned defeat testing are shown. All subjects received social defeat and were treated with 0.0 mg/kg, 0.3 mg/kg, 1.0 mg/kg, and 3.0 mg/kg of mCPP prior to social defeat.

Table 2.

Behavior	D 0.0 mg/kg	D 0.3 mg/kg	D 1.0 mg/kg	D 3.0 mg/kg	ND 0.0 mg/kg	ND 1.0 mg/kg	p value
Flee	1.800 ± .853	1.222 ± 1.102	1.091 ± .995	.182 ± .122	.000 ± .000	.000 ± .000	> .05
Stretch Attend	.000 ± .000	.111 ± .111	.091 ± .091	.167 ± .167	.000 ± .000	.250 ± .250	> .05
Attack	.500 ± .500	.000 ± .000	.000 ± .000	.455 ± .282	.500 ± .500	.000 ± .000	> .05

The frequencies of flee, stretch attend, and attack (mean ± SE) during conditioned defeat testing are shown. Defeated (D) animals were treated with 0.0 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg of mCPP and No Defeat (ND) animals treated with 0.0 mg/kg and 2.0 mg/kg of mCPP did not significantly differ in any category of behavior. Subjects received i.p. injection prior to conditioned defeat testing.

Table 3.

Behavior	D 0.0 mg/kg	D 0.5 mg/kg	D 2.0 mg/kg	ND 0.0 mg/kg	ND 2.0 mg/kg	p value
Flee	.400 ± .267	.091 ± .091	.200 ± .267	.200 ± .133	.000 ± .000	> .05
Stretch Attend	.400 ± .163	.182 ± .182	.200 ± .267	.000 ± .000	.000 ± .000	> .05
Attack	.000 ± .000	.273 ± .273	.200 ± .200	.556 ± .444	.375 ± .263	> .05

The frequencies of flee, stretch attend, and attack (mean ± SE) during conditioned defeat testing are shown. Defeated (D) animals were treated with 0.0 mg/kg 0.5 mg/kg, or 2.0 mg/kg of MDL 11,939 prior to social defeat. No Defeat (ND) animals were treated with 0.0 mg/kg or 2.0 mg/kg of MDL 11,939 before exposure to an aggressor's empty cage. Subjects did not significantly differ in any category of behavior.

Table 4.

Behavior	D 0.0 mg/kg	D 0.5 mg/kg	D 2.0 mg/kg	ND 0.0 mg/kg	ND 2.0 mg/kg	p value
Flee	1.182 ± 1.086	1.900 ± .824	1.455 ± .824	.000 ± .000	.625 ± .625	> .05
Stretch Attend	.000 ± .000	.300 ± .300	.273 ± .195	.000 ± .000	.000 ± .000	> .05
Attack	.000 ± .000	.000 ± .000	.000 ± .000	2.125 ± 1.716	.250 ± .250	= .027

The frequencies of flee, stretch attend, and attack (mean ± SE) during conditioned defeat testing are shown. Defeated (D) animals were treated with 0.0 mg/kg, 0.5 mg/kg, or 2.0 mg/kg of MDL 11,939 prior to conditioned defeat testing. No Defeat (ND) animals were treated with 0.0 mg/kg or 2.0 mg/kg of MDL 11,939 before conditioned defeat testing. No defeat controls attacked more often than did defeat subjects.

APPENDIX B




Figure 1. Durations (mean \pm S.E.) of submissive and defensive, aggressive, non-agonistic social, and nonsocial behavior are shown for a 5-minute test with a novel, non-aggressive opponent. Subjects received an injection of mCPP (0.3 mg/kg, N=11; 1.0 mg/kg, N=11; or 3.0 mg/kg, N=11) or vehicle (N=11) 15 min before social defeat training.




Figure 2. Durations (mean \pm S.E.) of submissive and defensive, aggressive, non-agonistic social, and nonsocial behavior are shown for a 5-minute test with a novel, non-aggressive opponent. Defeated animals received an injection of mCPP (0.3 mg/kg, N=10; 1.0 mg/kg, N=11; or 3.0 mg/kg, N=10) or vehicle (N=11) 15 minutes before behavioral testing. Likewise controls received an injection of mCPP (1.0 mg/kg, N=8) or vehicle (N=8) 15 minutes before behavioral testing. * indicates significantly different than defeated, vehicle controls. ** indicates significantly different than defeated subjects.




Figure 3. Durations (mean \pm S.E.) of submissive and defensive, aggressive, non-agonistic social, and nonsocial behavior are shown for a 5-minute test with a novel, non-aggressive opponent. Defeated animals received an injection of MDL 11,939 (0.5 mg/kg, N=11 or 2.0 mg/kg, N=10) or vehicle 30 minutes before social defeat training. Likewise, controls received an injection of MDL 11,939 (2.0 mg/kg, N=8) or vehicle (N=9) 30 minutes before exposure to a resident aggressor's empty cage. * indicates significantly different than defeated, vehicle controls. ** indicates significantly different than defeated subjects.




Figure 4. Durations (mean \pm S.E.) of submissive and defensive, aggressive, non-agonistic social, and nonsocial behavior are shown for a 5-minute test with a novel, non-aggressive opponent. Defeated animals received an injection of MDL 11,939 (0.5 mg/kg, N=11 or 2.0 mg/kg, N=10) or vehicle (N=10) 30 minutes before behavioral testing. Likewise controls received an injection of MDL 11,939 (2.0 mg/kg, N=8) or vehicle (N=8) 30 minutes before behavioral testing. * indicates significantly different than defeated, vehicle controls. ** indicates significantly different than defeated subjects.

APPENDIX C




Figure 5. Proposed neural circuit underlying 5-HT_{1A}, 5-HT_{2A}, and 5-HT_{2C} receptors role in conditioned defeat behavior. Social defeat activates 5-HT neurons in the dorsal raphe nucleus (DRN), in which in turn increases 5-HT release into the basolateral amygdala (BLA). 5-HT_{1A} receptor activation in the BLA inhibits glutamatergic projection cells causing a reduction in the acquisition of conditioned defeat behavior. 5-HT_{2A} receptors may facilitate the acquisition of conditioned defeat in two separate ways. 5-HT_{2A} receptor activation in the BLA may enhance activity of the glutamatergic cells projecting to the central amygdala, causing an increase in the acquisition of conditioned defeat behavior. Also, 5-HT_{2A} receptor activation in the BLA may cause desensitization of 5-HT_{2A} receptors on GABAergic interneurons and disinhibit glutamatergic projection cell causing a reduction in conditioned defeat behavior. 5-HT_{2C} receptor activation in the central amygdala on glutamatergic projection cell may an increase the expression of conditioned defeat behavior.

VITA

Marquinta Lee Harvey was born in Nashville, Tennessee on March 27, 1981. She obtained a B.S. degree in Microbiology at the University of Tennessee, Knoxville in May 2004. She entered the graduate program in the Psychology Department in August 2009. After obtaining a M.A. degree in Psychology in May 2011, she left to pursue a Doctorate of Philosophy with a focus in Behavioral Neuroscience. She plans to pursue a career in neuroscience research.