



5-2015

Characterization of head involution defective (hid) as a pro-apoptotic gene in *Megasalia scalaris*

Hakyong Kwak
hkwak1@vols.utk.edu

Follow this and additional works at: https://trace.tennessee.edu/utk_chanhonoproj

 Part of the [Molecular Genetics Commons](#)

Recommended Citation

Kwak, Hakyong, "Characterization of head involution defective (hid) as a pro-apoptotic gene in *Megasalia scalaris*" (2015). *University of Tennessee Honors Thesis Projects*.
https://trace.tennessee.edu/utk_chanhonoproj/1887

This Dissertation/Thesis is brought to you for free and open access by the University of Tennessee Honors Program at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in University of Tennessee Honors Thesis Projects by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Characterization of *head involution defective (hid)* as a pro-apoptotic gene in *Megaselia scalaris*

Hakyong Gloria Kwak

Department of Biochemistry and Cellular and Molecular Biology

The University of Tennessee, Knoxville

Faculty Advisor: Jae H. Park, PhD

Abstract

Apoptosis, a specific form of programmed cell death, is an evolutionary conserved mechanism that unwanted cells undergo as a way for organisms to maintain homeostasis. *Drosophila melanogaster* is known to be a widely studied model organism for the molecular pathways of apoptosis, typically the cell death genes *reaper*, *head involution defective (hid)*, *grim*, and *sickle*. Since the *hid* gene in both *D. melanogaster* and *Megaselia scalaris* were found to be homologous, experiments using *hid* were done to possibly characterize the gene as a pro-apoptotic gene. Transgenic lines of the *schid* gene located on different chromosomes were made and crossed to different UAS-gal4 lines for targeted expression. Most, if not all, of the neurons of all the different lines, underwent apoptosis. The results indicate that the functions of the *hid* gene are conserved and that the *hid* gene is a pro-apoptotic gene in *M. scalaris*. Also, when a caspase inhibitor was introduced, the killing nature of *schid* was mitigated, as some of the neurons were rescued, indicating that caspases were successfully inhibited.

Introduction

Overview of apoptosis

Programmed cell death (PCD) is crucial for normal development of animals. PCD helps maintain homeostasis by eliminating unwanted cells (Lee and Baehrecke, 2000). PCD is also called apoptosis and is a highly regulated genetic program of cell death. The term apoptosis was first coined in 1972 in a paper by Kerr, Wylie, and Currie in order to distinguish this form of cell death, which has distinct morphological characteristics (Elmore, 2007). During apoptosis, the cells shrink, the chromatin condense, the DNA fragments, all the while the integrity of the membrane is retained, and they are quickly eliminated by phagocytosis (Fuchs, 2011). Apoptosis helps maintain cellular proofreading and defend organisms from dangerous cells, such as those infected with pathogens (Hay et al., 2004). Defective apoptosis can lead to neurodegenerative diseases, autoimmune diseases, and many types of cancers (Elmore, 2007).

Although the cell death of certain neurons can cause neurodegenerative diseases, apoptosis is necessary for the proper development of the nervous system. In vertebrates, about 50% of the neurons undergo apoptosis at some point during development (Yeo et al., 2004). Apoptosis can occur in both proliferating (early PCD) and differentiated neurons (neurotrophic cell death) (Yeo et al., 2004). Early PCD is observed in a variety of different organisms. Although the fact that early PCD is seen in different organisms (*C. elegans*, mouse, *D. melanogaster*) suggests that this form of apoptosis is conserved evolutionarily (Yeo et al., 2004), there is little known about the trigger for early PCD (Yeo et al., 2004). In neurotrophic cell death, neurons compete for a finite number of nerve growth factors that is released by the neurons' target cells (Yeo et al., 2004). The neurons that do not receive the nerve growth factor eventually undergo apoptosis.

Much of what is known about the mechanisms of apoptosis was elucidated from studying PCD in the development of the nematode *C. elegans*. Studying this organism led to the discovery of the core cell death machinery called caspases (Fuchs, 2011). Caspases are a family of cysteine proteases that act as the ultimate executioners of apoptosis (Fuchs, 2011; Lee et al., 2013). In general, an apoptotic stimulus (or stimuli) leads to an interaction between signaling proteins. This causes an apoptotic protease factor to induce an initiator caspase that stimulates effector caspases that commit the cells to apoptosis.

Comparative mechanisms of apoptosis

In *C. elegans*, 131 of the 1090 somatic cells undergo apoptosis during the process of forming an adult worm. The start of apoptosis is controlled by *ced-9*, *ced-4*, and *ced-3*. Two of the genes, *ced-9* and *ced-3*, have mammalian homologs that have similar roles during apoptosis; *ced-9* is part of the *Bcl-2* family and *ced-3* is homologous to a family of cysteine proteases called caspases (Grether et al., 1995). Also, expression of human *Bcl-2* can partially make-up for the loss of *ced-9* function in *C. elegans* and suppress some PCD (Grether et al., 1995). Cell death in mammals and flies also show many similar morphological and biochemical hallmarks (Hay et al., 2011). This shows that many of the components of cell death have been conserved throughout evolution. However, despite many studies done on apoptosis regulation and mechanisms in *C. elegans*, *D. melanogaster*, and mammals, the specifics of this mechanism are still largely unknown.

D. melanogaster is a useful model system to study apoptosis because it occurs throughout the development of the fly and has been extensively used as a genetic model system to study the molecular pathways leading to apoptosis (Hay et al., 2004). The adaptor protein ARK interacts with the initiator caspase DRONC, which activates subsequent effector caspases. Cells survive because they express DIAP1, an inhibitor of apoptotic proteins (IAP) that inhibits Dronc and its downstream effects (Hay et al., 2004). Reaper, Grim, and Hid, together known as RHG proteins, bind to DIAP-1, interfering with the DIAP1-caspase complex, which disrupts the anti-caspase function of DIAP1 (Hay et al., 2004).

Molecular functions of the *hid* gene

The *head involution defect* (*hid*) death gene plays an important role in cell death in *D. melanogaster*. The *hid* gene is located in the H99 interval, a locus on the third *Drosophila* chromosome (Grether et al., 1995). Deletion of this genomic region blocked *hid*-induced cell death (Grether et al., 1995). The *hid* mRNA is expressed in the embryo where cell death occurs, and the gene encodes for a 410-amino-acid protein (Grether et al., 1995). Initially, in order to know the role *hid* has in PCD, the effect of expressing *hid* cDNA was studied. A construct in which *hid* cDNA was controlled by the *hsp70* heat shock promoter was made. The heat shocked embryos containing this construct were seen to have much higher levels of PCD, which occurred by apoptosis (Grether et al., 1995). The patterns of *hid* mRNA expression and cell death were also

observed to be correlated (Grether et al., 1995). Interestingly, the *hid* gene seems to work cooperatively, yet with distinct functions, with the other death genes, *rpr* and *grim*, to promote PCD as a way to prevent unintended cell death in corazonin neurons (Sandu et al., 2010).

Biology of Scuttle flies

To understand how the cell death mechanisms are conserved throughout evolution, Park laboratory recently cloned and characterized a cell death gene, *hid*, in Scuttle Fly (*Megaselia scalaris*), a species that has diverged from the *Drosophila* lineage around 150 million years ago. Scuttle flies are so named because they walk in rapid bursts of movement with short pauses in between (Harrison et al., 2003). As compared to *D. melanogaster*, scuttle flies mature at a slower rate and their movements, such as the rate of larval body wall contractions and mouth hook movements, are slower as well (Harrison et al., 2003). The Scuttle *hid* gene encodes a protein of 197 amino acids, which is much shorter than *D. melanogaster hid* protein of 410 amino acids. Despite the size difference, several regions of the gene were found to be highly conserved. This project aims to characterize the role of *head involution defective (hid)* as a potential pro-apoptotic gene in *Megasalia scalaris*.

Methods

Transgenic line

To see if the function of the *hid* gene was evolutionary conserved, the Scuttle *hid* gene (*schid*) was injected into the *Drosophila* embryo using a P-element mediated transformation to produce UAS-*schid* transgenic flies. We established two independent insertions, one on the second and on the third chromosome. We overexpressed *schid* using UAS-GAL4 system in *Drosophila* with five different GAL4 lines that were combined with UAS-*lacZ* to drive the reporter expression in Pigment dispersion factor (Pdf), buriscon (Burs), partner of buriscon receptor (pburs), corazonin (Crz), and neuropeptide F (Npf) expressing neurons. It was evident from the X-gal staining that most of the peptidergic neurons underwent apoptosis. In order to confirm that the apoptotic function of the transgene was because of the nature of the transgene itself and not because of the location of the gene in the genome, *schid* on the third chromosome, *yw;;schid^{III}* was expressed in the GAL4 lines indicated in the table below. We also made a double transgenic line with *schid* and *p35*. The *p35* gene is a caspase inhibitor that inhibits

caspase induced cell death. If the apoptotic function was due to the nature of the transgene itself, then that the *p35* gene should inhibit the activity of the caspase induced by the *U-schid* gene, and the peptidgeric neurons of the larvae ideally should not undergo apoptosis or will have partial apoptosis depending on the neuron. The presence of neurons was observed by staining the tissues using the X-gal staining protocol and viewing the mounted samples using a light microscope.

X-gal staining

X-gal staining was typically used to monitor the expression of genes using a light microscope. X-gal, which is an analog of lactose, is used to test for the presence of Beta-galactosidase, and enzyme produce by the *LacZ* gene. In the presence of an active Beta-galactosidase enzyme, X-gal is cleaved to yield galactose and 5,5'-dibromo-4,4'-dichloro-indigo, which is the blue precipitate that is easily detected with a microscope (Levitsky et al., 2013).

X-gal staining was performed to stain the neurons. The central nervous system (CNS) from third instar stage larvae from the F1 generation of the UAS/GAL4 crosses (indicated in the table below) was dissected in phosphate buffered saline (PBS) on ice. The tissues were then fixed with 8% glutaraldehyde for ten min. and then washed in PBS (3 X 5 min.). The tissues were then stained with x-gal staining solution (0.2 M NaPi, 5M NaCl, 1M MgCl₂, K-Ferro-CN (P-3289/3667, Sigma), 10% Triton X-100) and kept in a dark, damp container at 37°C until the blue staining was clearly visible (1.5-2 h). The tissues were then rinsed in PBS and subsequently in 70% ethanol (2 X 30 min). The tissues were then kept in 90% glycerol and the neurons were viewed using a light microscope.

Fly crossings

Below are all the crosses that were done and dissected.

Females (virgin)	Males
yw; Crz-gal4 ^{S8} ; UAS-lacZ	w ¹¹¹⁸
	UAS-schid ^{S3}
	UAS-schid ^{S3} ; UAS-p35
	UAS-schid ^{III} /TM6B
Burs-gal4 ^x ; UAS-lacZ; +(TM6B)	w ¹¹¹⁸
	UAS-schid ^{S3}
	UAS-schid ^{S3} ; UAS-p35


Pdf-gal4 ^M ; UAS-lacZ; +/- (TM6B)	w ¹¹¹⁸
	UAS-schid ^{S3}
	UAS-schid ^{S3} ; UAS-p35
	UAS-schid ^{III} /TM6B
yw; UAS-lacZ ^{II} ; npf-gal4 ^{III}	w ¹¹¹⁸
	UAS-schid ^{S3}
	UAS-schid ^{S3} ; p-35
UAS-schid ^{III} /TM6B	yw; Crz-gal4 ^{S8} ; UAS-lacZ
	Pdf-gal4 ^M ; UAS X-lacZ; +/- (TM6B)
UAS-schid ^{II} ; UAS-lacZ	::crz-gal4 ^{t2a}
	::crz-gal4 ^{t2a} ; UAS-p35 ^{III}
Pburs-gal4 ^{S3}	yw; Bl/cyo,y+; UAS-lacZ
Pburs-gal4 ^{S3}	yw; UAS-schid; UAS-lacZ
Pburs-gal4 ^{S6}	yw; Bl/cyo,y+; UAS-lacZ
Pburs-gal4 ^{S6}	yw; UAS-schid; UAS-lacZ

Results and Discussion

X-gal Staining

In order to visualize the effects *schid* and *p35* have on the neurons expressing the neuropeptides Pigment dispersion factor (Pdf), buriscon (Burs), partner of buriscon receptor (pburs), corazonin (Crz), and neuropeptide F (Npf), lacZ staining was performed on the brains of third instar larvae.

In larvae, Pdf is expressed in the lateral and abdominal neurons. Using the UAS-GAL4 system, when *schid*^{S3} was overexpressed in Pdf containing lateral neurons, not all of the neurons underwent apoptosis (Fig. 1B). However, when *schid*^{III} was overexpressed in the lateral neurons, all of the neurons underwent apoptosis (Fig. 1C).



Bursicon (BURS) is typically expressed in the ventral nerve cord –four pairs of two neurons followed by four single neurons (Fig. 2A). BURS plays an important role in wing expansion, as well as in tanning and plasticization of the wings (Peabody et al., 2008). When both fly strains of *schid* was overexpressed in the bursicon expressing neurons, all the neurons underwent apoptosis (Fig. 2B, C). A similar result is seen in the pburs expressing neurons (Fig. 2F, H). Like BURS, pBURS also plays a role in wing expansion and the phenotypic effects of the expression of *schid* are visible (Luo et al., 2005). Because of the role in wing expansion pburs has, the hypothesis was that the wings of the F1 generation of the pburs strains crossed with *schid* would be abnormal. Two different pburs strains were tested: Pburs-gal4^{S3} and Pburs-gal4^{S6} with UAS-*schid*; UAS-*lacZ* as the variable and *yw*; *Bl/cyo,y+*;UAS-*lacZ* as the control. As expected, the neuroptides of the larvae crossed with UAS-*schid*; UAS-*lacZ* underwent apoptosis, and the wings of the F1 generation adults did not expand (Fig.2I). The adults of both of the controls had the normal wing phenotype (Fig. 2J).




Figure 2: Burs and pburs neurons detected with X-gal staining. A: control larval CNS (4 pairs of neurons followed by four single neurons in the ventral nerve cord (VNC); n=10. B,C: complete cell death of Burs neurons; n=10. D. Partial rescue (50%) of PDF neurons by p35 expression in both the VNC; n=8. E: control larval CNS expressing pburs (s3) neurons; n=10. F: complete cell death of pburs neurons; n=10. G: control larval CNS expressing pburs (s6) neurons; n=10. H: complete cell death of pburs neurons; n=9. I. image of F1 generation of Pburs X yw; Bl/cyo,y+; UAS-lacZ. J: image of F1 generation of Pburs X yw; UAS-schID; UAS-lacZ. Pictures taken at 10X.

The corzanonin neuropeptide (Crz) is expressed in the dorso-lateral (DL) neurons, the dorso-medial neurons (DM), and in the ventral nerve cord (vCrz) (Fig. 3A). When *schID* was overexpressed using UAS-*schid*^{S3} in the Crz expressing neurons using *Crz-GAL4*^{S8}, there were a small number of neurons that survived (Fig. 3B). However, when overexpressed using UAS-*schid*^{III}, all the Crz expressing neurons underwent complete apoptosis (Fig. 3C).




Figure 3: Crz expressing neurons detected with X-gal staining. A: control larval CNS (are two sets of dorsal neurons, DM (arrowhead) and DL (arrow), and 8 pairs of neurons in the ventral nerve cord); n=10. B: An average of 1.7 neurons remaining; n=10. C: Shows complete cell death of neurons; n=10. D: Partial rescue (50.6%) of Crz neurons by p35 expression in both the lateral and abdominal regions; n=10. Pictures taken at 10X.

In the larval stages, NPF is expressed in four neurons in the lateral region (Fig. 4A). When *schid*^{S3} was ectopically expressed in the four lateral neurons expressing Npf, an average of 0.33 neurons did undergo apoptosis (Fig. 4B). However, when *schid*^{III} was ectopically expressed in the same neurons, all of the neurons underwent apoptosis (Fig. 4C).




Figure 4: Npf expressing neurons detected with X-gal staining. A: control larval CNS expressing Npf neurons (four lateral neurons); n=8. B: An average of 0.33 neurons remaining; n=9. C: complete cell death of neurons; n=9. D: Partial rescue (75%) of Pdf neurons by p35 expression in the lobes; n=9. Pictures taken at 10X.

When the above mentioned genotypes were crossed with *UAS-schid^{S3}*; *UAS-p35*, there was partial recovery of the neurons, although the amount of recovery varied from genotype to genotype, it is clear that the *p35* gene was able to recover some of the neurons. 88% of the Pdf neurons were rescued, 50% of the Burs neurons were rescued, 50.6 % of the Crz neurons were rescued, and 75% of the Npf neurons were rescued (Fig. 1D, 2D, 3D, 4D). As previously mentioned, the *p35* gene is a caspase inhibitor. In this case, *p35* was combined with the *hid* gene. The partial recovery of neurons suggests that it was successful in blocking the caspase activated by the *hid*. This also reinforces the idea that *p35* acts downstream of *hid* (Grether et al., 1995). On the other hand, the partial recovery is also indicative of the fact that other cell death genes, *grim* and *rpr*, also play a role in apoptosis. In a species related to *Drosophila*, *Anastrepha suspensa*, the cell death activity was higher when *Anastrepha-hid* and *Anastrepha-rpr* were expressed together (Schetelig et al., 2011). On the other hand, induced cell death by *rpr* and *hid* was blocked in *Drosophila* when *p35* was introduced (Zhou et al., 1997).

This cooperative interaction between death genes *hid*, *grim*, and *rpr* to induce programmed cell death may also explain why the Pdf, Crz, and Npf expressing neurons did not undergo complete cell death (figures 1B, 3B, 4B). The interaction between *schid* and the other death genes may not have been strong enough to promote complete PCD in the samples. Another reason may be because the apoptotic function of the *grim* death gene is sufficient enough to carry

out PCD even in the absence of *hid* and *rpr* (Cho et al., 1996). Thirdly, the effect of the GAL-4 drivers on their target gene may vary depending on the location of the *schid* gene. Although there were a small number of samples still present when *schid*^{S3} was expressed, when *schid*^{III} was expressed, the neurons underwent complete apoptosis.

The results reveal that the function of the *hid* gene is not specific to one neuropeptide producing neuron, but affects BURS, NPF, CRZ, and PDF producing neurons. The fact that the killing nature of the *hid* gene is seen in all of the neurons tested indicates that the function of the *hid* gene in PCD is conserved between *D. melanogaster* and *M. scalaris*. These results bring us one step closer to exploring PCD in *M. scalaris*.

Acknowledgements

I would like to thank Dr. Jae H. Park for all of his guidance and encouragement during this project and over the past three years. I would also like to thank the other members of the Park lab for their advice, especially Dr. Geni Lee for her work in producing the UAS-scHID transgenic lines

References

- Cho Po, Nordstrom W, Gish B, and Abrams JM (1996). *Grim*, a novel death gene in *Drosophila*. *Genes Dev* 10(14):1773-82.
- Elmore, Susan (2007). Apoptosis: A Review of Programmed Cell Death. *Toxicol Path* 35(4): 495-516.
- Fuchs Y, Steller H (2011). Programmed cell death in animal development and disease. *Cell* 147(4):742-58.
- Grether ME, Abrams JM, Agapite J, White K, and Steller H (1995). The head involution Defective gene of *Drosophila melanogaster* functions in programmed cell death. *Genes*

Dev 9: 1694-1708.

- Harrison DA and Cooper RL (2003). Characterization of development, behavior and neuromuscular physiology in the phorid fly, *Megaselia scalaris*. *Comp Biochem Physiol A Mol Integr Physiol* 136: 427-439.
- Hay BA, Huh JR, and Guo M (2004). The genetics of cell death: approaches, insights and opportunities in *Drosophila*. *Nat Rev Genet* 5:911-922.
- Lee G, Sehgal R, Wang Z, Nair S, Kikuno K, Chen C-H, Hay B, and Park JH (2013). Essential role of *grim*-led programmed cell death for the establishment of corazonin-producing peptidergic nervous system during embryogenesis and metamorphosis in *Drosophila melanogaster*. *Biology Open* 2: 283-94.
- Levitsky KL, Toledo-Aral JJ, Barneo JL, and Villadiego J (2013). Direct confocal acquisition of fluorescence from X-gal staining on thick tissue sections. *Sci Rep* 14;3:2937.
- Luo CW, Dewey EM, Sudo S, Ewer J, Hsu SY, Honegger HW, Hsueh AJ (2005). Bursicon, the insect cuticle-hardening hormone, is a heterodimeric cysteine knot protein that activates G protein-coupled receptor LGR2. *Proc Natl Acad Sci U.S.A* 102(8):2820-5.
- Peabody NC, Diao F, Luan H, Wang H, Dewey EM, Honegger HW, and White BH (2008). Bursicon Functions within the *Drosophila* Central Nervous System to Modulate Wing Expansion Behavior, Hormone Secretion, and Cell Death. *J Neurosci* 28(53):14379-14391.
- Sandu C., Ryoo H. D., Steller H. (2010). *Drosophila* IAP antagonists form multimeric complexes to promote cell death. *J. Cell Biol* 190:1039–1052.
- Schetelig MF, Nirmala X, and Xavier N (2011). Pro-apoptotic cell death genes, *hid* and *reaper*, from the tephritid pest species, *Anastrepha suspensa*. *Apoptosis* 16:759-68.
- Yeo W, Gautier J (2004). Early neural cell death: dying to become neurons. *Dev Bio.* 15;274(2):233-44.
- Zhou L, Schnitzler A, Schwartz LM, Steller H, and Nambu JR (1997). Cooperative functions of the *reaper* and *head involution defective* genes in the programmed cell death of *Drosophila* central nervous system midline cells. *PNAS.* 13; 94(10): 5131–5136.