



7-23-2010

# Developing measures for microbial genome assembly quality control

Rachel M. Adams

*University of Tennessee - Knoxville, radams29@utk.edu*

Jason B. Harris

*University of Tennessee - Knoxville, jharri43@utk.edu*

Jeremy J. Jay

*University of Tennessee - Knoxville, jay@eeecs.utk.edu*

Beth G. Johnson

*University of Tennessee - Knoxville*

Miriam L. Land

*Oak Ridge National Laboratory*

*See next page for additional authors*

Follow this and additional works at: [http://trace.tennessee.edu/utk\\_elecpubs](http://trace.tennessee.edu/utk_elecpubs)

 Part of the [Electrical and Computer Engineering Commons](#)

---

## Recommended Citation

BMC Bioinformatics 2010, 11(Suppl 4):P14 doi:10.1186/1471-2105-11-S4-P14

This Article is brought to you for free and open access by the Electrical Engineering and Computer Science at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in EECS Publications and Other Works by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact [trace@utk.edu](mailto:trace@utk.edu).

---

**Authors**

Rachel M. Adams, Jason B. Harris, Jeremy J. Jay, Beth G. Johnson, Miriam L. Land, and Loren J. Hauser

POSTER PRESENTATION

Open Access

# Developing measures for microbial genome assembly quality control

Rachel M Adams<sup>1</sup>, Jason B Harris<sup>1</sup>, Jeremy J Jay<sup>2\*</sup>, Beth G Johnson<sup>3</sup>, Miriam L Land<sup>4</sup>, Loren J Hauser<sup>4</sup>

From UT-ORNL-KBRIN Bioinformatics Summit 2010  
Cadiz, KY, USA. 19-21 March 2010

## Background

Advances in sequencing technologies are outpacing the rate at which genomes can be thoroughly finished and analyzed. Over the next year, genome sequencing will increase many-fold, but high quality and high-throughput annotation methods have yet to be developed to handle the need. As more microbial genomes are sequenced, whole-genome annotation methods identify many putative genes which need further verification. By analyzing a broad range of annotated genomes we can identify patterns and statistics useful in determining the annotation quality and spurious gene outliers. Our work is attempting to identify quality control measures based on a full inter-genomic comparison instead of individual sequence-level or database-specific statistics. Using these methods to compare and filter, it is possible to narrow the scope of manual gene curation and allow greater scrutiny on putative genes before publication, making higher quality genome annotation possible. Our results plainly show the quality of well-studied genomes, the weaknesses of draft genome builds, and illustrate the need for further high-throughput quality control measures.

## Author details

<sup>1</sup>Genome Science and Technology Graduate School, University of Tennessee, Knoxville, TN 37996, USA. <sup>2</sup>Department of Electrical Engineering and Computer Science, University of Tennessee, Knoxville, TN 37996, USA. <sup>3</sup>Department of Mathematics, University of Tennessee, Knoxville, TN 37996, USA. <sup>4</sup>Computational Biology and Bioinformatics Group, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA.

Published: 23 July 2010

\* Correspondence: jay@eecs.utk.edu

<sup>2</sup>Department of Electrical Engineering and Computer Science, University of Tennessee, Knoxville, TN 37996, USA

doi:10.1186/1471-2105-11-S4-P14

Cite this article as: Adams et al.: Developing measures for microbial genome assembly quality control. *BMC Bioinformatics* 2010 11(Suppl 4):P14.

Submit your next manuscript to BioMed Central  
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

BioMed Central