



April 2010

W197 Dectes Stem Borer

The University of Tennessee Agricultural Extension Service

Follow this and additional works at: http://trace.tennessee.edu/utk_agexcrop



Part of the [Entomology Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

"W197 Dectes Stem Borer," The University of Tennessee Agricultural Extension Service, W197 09-0095, http://trace.tennessee.edu/utk_agexcrop/96

The publications in this collection represent the historical publishing record of the UT Agricultural Experiment Station and do not necessarily reflect current scientific knowledge or recommendations. Current information about UT Ag Research can be found at the [UT Ag Research website](#).

This Insect, Pest and Disease Control - Soybeans is brought to you for free and open access by the UT Extension Publications at Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Field & Commercial Crops by an authorized administrator of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Soybean Insects

Dectes Stem Borer

Scott Stewart, Professor, Entomology and Plant Pathology
 Angela Thompson McClure, Associate Professor, Plant Sciences
 and Russ Patrick, Professor, Entomology and Plant Pathology

Classification and Description: The Dectes stem borer, *Dectes texanus texanus*, is a long-horned beetle (Coleoptera: Cerambycidae) whose larva tunnels in the stem of soybean and wild host plants. This insect was first reported in soybeans in Tennessee during the early 1970s. The adult beetle is gray and approximately 3/8 inch in length. Eggs are yellowish, shiny, elongated and darken to an amber color prior to hatching. The legless larva is creamy white or yellowish with brown mouthparts. It is less than 1/16 inch long when it hatches from the egg, but reaches a length of 1/2 to 5/8 inch when full grown. The pupa is yellowish-white, turning to dark brown before the adult emerges.



Hosts, Life History and Distribution: Dectes stem borer are native to the United States east of the Rocky Mountains. Besides soybean, giant ragweed and cocklebur are known hosts. It is believed that Dectes

stem borer has only one generation annually. Mature larvae overwinter in the stem at the base of infested plants, near or just below the soil level. Pupation occurs in the spring. The pupal stage lasts 10-15 days, with emergence of adults beginning in late June and continuing through August. Mating takes place approximately five days after adults emerge from the pupal stage. The female beetle chews a small hole in the leaf petiole, or less commonly in stems, where she lays a single egg. An adult female can live 4-8 weeks and lay as many as 50 eggs.

Pest Status and Injury: It is not clear whether the Dectes stem borer causes direct yield loss. Fields can have 90 percent or more of plants infested by larvae. Some defoliation occurs from larval tunneling in leaf petioles that eventually causes leaf abscission. As it grows, a larva moves into the main stem from an infested leaf petiole where it tunnels within the pith of the main stem until the plant matures. More than one larva may infest a soybean plant, but only one individual survives to maturity. In preparation for overwintering, older larvae girdle the stem from the inside at 2-5 inches above the soil. Recent evidence suggests that tunneling in leaf petioles or



in the main stem has little effect on yield. However, girdling in preparation for overwintering can cause lodging, particularly during high winds. Lodging has been reported to cause yield loss or slow harvesting operations.

Management Considerations: *Dectes* stem borer has become increasingly common in Tennessee. Widespread adoption of reduced tillage systems has undoubtedly increased the occurrence of this insect. Foliar insecticides are not recommended for control of *Dectes*, in part because they are not very effective, and also because recent data suggest that even severe infestations cause little economic damage. However, infested fields should be harvested as soon as possible

to reduce any lodging associated with girdling activity. *Dectes* appear to prefer reproductive-stage plants for oviposition. Late-planted soybeans may sometimes be less infested by missing the peak oviposition activity of adult females. However, late-planted soybeans are more likely to have other pest problems such as stink bugs and loopers.

References:

Handbook of Soybean Insect Pests. L. G. Higley and D. J. Boethel (eds.), Entomological Society of America, 1994.

Dectes Stem Borer in Tennessee, C. Patrick, G. Lentz, S. Stewart and A. Thompson. University of Tennessee Extension, SP503-F.



Larva in stem



Girdle caused by overwintering larva



Wilted leaf petiole caused by larva tunneling

This publication contains pesticide recommendations that are subject to change at any time. The recommendations in this publication are provided only as a guide. It is always the pesticide applicator's responsibility, by law, to read and follow all current label directions for the specific pesticide being used. The label always takes precedence over the recommendations found in this publication.

Use of trade or brand names in this publication is for clarity and information; it does not imply approval of the product to the exclusion of others that may be of similar, suitable composition, nor does it guarantee or warrant the standard of the product. The author(s), the University of Tennessee Institute of Agriculture and University of Tennessee Extension assume no liability resulting from the use of these recommendations.