



National Quail Symposium Proceedings

Volume 6

Article 55

2009

Index

Follow this and additional works at: <http://trace.tennessee.edu/nqsp>

Recommended Citation

(2009) "Index," *National Quail Symposium Proceedings*: Vol. 6 , Article 55.
Available at: <http://trace.tennessee.edu/nqsp/vol6/iss1/55>

This Index is brought to you for free and open access by Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

Index

A

Alectoris rufa, 267
 abundance, 178, 220, 251
 action plan, 291
 active management, 370
 adaptive management, 2
 age, 135
 ageing, 505
 agricultural intensification, 111
 agriculture, 172, 232
 Amdro[®], 78
 Ames Plantation, 121
 area curves, 489
 Arizona, 370
 Arkansas, 146, 184
 Austria, 410
 autumn, 178

B

Bonasa umbellus, 480, 499
 barriers, 322
 basal area, 178
 Belize, 99
 biological, 322
 black-throated bobwhite, 99
 breeding success, 267
 brood, 88
 brood size, 267
 brood survival, 410
 broods, 302
 buffer strips, 232

C

Callipepla gambelii, 106
Callipepla squamata, 210, 456
Colinus nigrogularis, 99
Colinus virginianus, 16, 46, 54, 66, 78, 88, 121, 135, 146, 160, 178, 184, 193, 220, 251, 256, 271, 282, 315, 328, 338, 348, 370, 396, 432, 447, 467, 518
Cyrtonyx montezumae, 322, 426
 cactus, 256
 call counts, 46
 carcass fat, 499
 Chihuahuan Desert, 456
 Chihuahuan desert, 210
 climate trends, 328
 closed-canopy pine, 135
 compositional analysis, 410
 conservation, 111

Conservation Reserve Program, 302
 crippling loss, 420

D

data collection, 370
 decision making, 2
 density, 46
 discriminant function analysis, 505
 dispersal, 135
 distance sampling, 46
 disturbance, 391

E

east Texas, 16, 66
 edge, 184

F

farming, 172
 farmland, 348
 fat condition, 499
 field borders, 232
 fire, 193, 271, 315
 Florida, 88, 193, 271, 282
 food availability, 160
 food habits, 160
 food plots, 16, 66
 food preferences, 160
 food quantity, 146
 forestland, 348
 formal inference-based recursive modeling (FIRM), 302
 French Mediterranean region, 267

G

Gambel's quail, 106
 game crop, 267
 genetic variability, 121
 Georgia, 46, 135, 282, 338, 348
 GIS, 106, 302
 gray partridge, 24, 27, 291, 380, 391
 guzzlers, 210

H

habitat, 106, 135, 172, 193, 251, 302, 410
 habitat enhancement, 391
 habitat loss, 370
 habitat management, 260, 267
 habitat modeling, 220
 habitat quality, 184
 habitat selection, 88

- habitat use, 232, 391
 half-cutting, 251
 harvest, 260, 420
 herbicide, 315
 home range, 232, 410, 489
 hunter recruitment, 370
 hunter retention, 370
 hunting, 338
 hunting management, 267
 hunting season, 322
 hybridization, 121
- I
- incentives, 348
 insecticide, 78
 invertebrate biomass, 16, 66
 invertebrate richness, 16, 66
- J
- Johnson's rank method, 160
- K
- kernel estimator, 489
- L
- landowners, 348
 landscape composition, 432
 landscape structure, 432
- M
- management, 11, 251, 456
 management philosophies, 146
 mesomammal abundance, 447
 metapopulation, 135
 microsattellites, 121
 milo, 282
 minimum convex polygon, 489
 Minnesota, 302
 Mississippi, 220
 modern quail management, 338
 Montana, 260
 Montezuma quail, 322, 426
 morning covey call, 38
 mortality, 54, 467
- N
- Nothura maculosa*, 111
 necklace-type collar transmitter, 480
 nest, 256
 nesting success, 447
 New Mexico, 456
 northern bobwhite, 16, 38, 46, 54, 66, 78, 88, 121, 135, 146, 160, 178, 184, 193, 220, 232, 251, 256, 271, 282, 315, 328, 338, 348, 396, 420, 432, 447, 467, 489, 518
- O
- Ouachita Highlands, 146
- P
- Perdix perdix*, 24, 27, 291, 380, 391
Phasianus colchicus, 302, 410, 505
Procyon lotor, 256
 pairs, 172
 parent fostering, 391
 Partridge Count Scheme, 27
 passive management, 370
 pen-raised quail, 396
 pen-reared, 121
 pheasant, 260, 505
 pine-bluestem, 184
 pine-grassland restoration, 146
 plantation, 338
 population, 420
 population dynamics, 267
 population estimate, 99
 population recovery, 291
 pre-season release, 396
 predation, 24, 256, 456
 predation control, 24
 predator control, 447
 PREFER, 160
 prescribed burning, 184, 271
 prescribed fire, 78, 146
 PROC NL MIXED, 232
 productivity, 271
 Program MARK, 232
 proximal primary feather, 505
- Q
- quail, 88, 178, 282, 370
- R
- raccoon, 256
 radio telemetry, 426
 radio-handicapping, 518
 radio-marked, 426
 radio-telemetry, 338
 radiotelemetry, 456
 radius of audibility, 38
 re-introduction, 391
 red imported fire ant, 78
 red-legged partridge, 267

- reintroduction, 380
 - releasing, 380
 - reproduction, 232
 - research, 370
 - research and monitoring, 2
 - restocking, 380
 - ring-necked pheasant, 302
 - riparian vegetation, 106
 - Rolling Plains, 447
 - ruffed grouse, 499
 - ruffed grouse chicks, 480
 - rural abandonment, 267
- S
- Solenopsis invicta*, 78
 - Sorghum vulgare*, 282
 - salt cedar, 106
 - sample size, 46
 - SAS, 232
 - scale, 135
 - scale of management, 271
 - scaled quail, 210, 456
 - September burn, 315
 - sex, 135
 - simulation, 46
 - sociological, 322
 - South Carolina, 396
 - south Texas, 328
 - southeastern USA, 54, 467, 518
 - southern African gamebirds, 11
 - space use, 220
 - spotted tinamou, 111
 - suitability, 220
 - supplemental feed, 282
 - supplemental feeding, 210
 - surveillance camera, 210
 - survey, 38
 - survival, 54, 111, 121, 172, 232, 271, 391, 426, 432, 456, 467, 518
 - sweetgum, 315
 - Switzerland, 391
- T
- Texas, 420
 - telemetry, 54, 467, 518
 - Tennessee, 121
 - Texas, 78, 106, 251, 322, 447, 456
 - threat, 11
 - timber, 178
 - TOBEC, 499
 - total body electrical conductivity, 499
- Trans-Pecos, 106, 322
- U
- UK, 24, 27, 291
 - Upper Coastal Plain, 348
 - urbanization, 370
 - usable space, 146, 184
 - utilization, 11
- W
- water developments, 210
 - weather, 193
 - West Virginia, 480
 - wild pheasant, 410
 - wildlife stand improvement, 184
 - winter habitat, 432
 - woody control, 315